

AN EXTENSION OF THE EXTENDED HURWITZ-LERCH ZETA FUNCTIONS OF TWO VARIABLES

JUNESANG CHOI, RAKESH K. PARMAR, AND RAM K. SAXENA

ABSTRACT. We aim to introduce a further extension of a family of the extended Hurwitz-Lerch Zeta functions of two variables. We then systematically investigate several interesting properties of the extended function such as its integral representations which provide extensions of various earlier corresponding results of two and one variables, its summation formula, its Mellin-Barnes type contour integral representations, its computational representation and fractional derivative formulas. A multi-parameter extension of the extended Hurwitz-Lerch Zeta function of two variables is also introduced. Relevant connections of certain special cases of the main results presented here with some known identities are pointed out.

1. Introduction and preliminaries

Throughout this paper, let \mathbb{C} , \mathbb{R}^+ , \mathbb{Z}_0^- and \mathbb{N}_0 be the sets of complex numbers, positive real numbers, non-positive and non-negative integers, respectively, and $\mathbb{N} := \mathbb{N}_0 \setminus \{0\}$. The *Hurwitz-Lerch Zeta function* $\Phi(z, s, a)$ is defined by (see, e.g., [5, p. 27, Eq. 1.11(1)]; see also [19, p. 121] and [20, p. 194]):

$$(1) \quad \Phi(z, s, a) := \sum_{n=0}^{\infty} \frac{z^n}{(n+a)^s}$$

$$(a \in \mathbb{C} \setminus \mathbb{Z}_0^-; s \in \mathbb{C} \text{ when } |z| < 1; \Re(s) > 1 \text{ when } |z| = 1).$$

For various properties and the special cases of the Hurwitz-Lerch Zeta function $\Phi(z, s, a)$, one may refer to [19] and [20]. Many generalizations of the Hurwitz-Lerch Zeta function $\Phi(z, s, a)$ have been investigated (see, e.g., [1–3, 5–7, 9, 11, 18, 21, 23, 25]). In particular, Srivastava et al. [25, p. 491, Eq. (1.20)] introduced

Received July 5, 2016; Revised February 3, 2017; Accepted February 20, 2017.

2010 *Mathematics Subject Classification.* Primary 11M25, 11M99, 33B15, 33C60; Secondary 11M35, 11B68, 33C05, 33C90.

Key words and phrases. Hurwitz-Lerch Zeta function, extended Hurwitz-Lerch Zeta functions, Gauss hypergeometric function, Fox-Wright hypergeometric function, Mellin-Barnes contour integral representations, analytic continuation.

and investigated the following extended Hurwitz-Lerch Zeta function:

$$(2) \quad \Phi_{\lambda, \mu; \nu}^{(\rho, \sigma; \kappa)}(z, s, a) := \sum_{n=0}^{\infty} \frac{(\lambda)_{\rho n} (\mu)_{\sigma n}}{(\nu)_{\kappa n} n!} \frac{z^n}{(n+a)^s}$$

($\lambda, \mu \in \mathbb{C}$; $a, \nu \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $\rho, \sigma, \kappa \in \mathbb{R}^+$; $\kappa - \rho - \sigma > -1$ when $s, z \in \mathbb{C}$; $\kappa - \rho - \sigma = -1$ and $s \in \mathbb{C}$ when $|z| < \delta^* := \rho^{-\rho} \sigma^{-\sigma} \kappa^{\kappa}$; $\kappa - \rho - \sigma = -1$ and $\Re(s + \nu - \lambda - \mu) > 1$ when $|z| = \delta^*$), where $(\lambda)_{\nu}$ is the well-known and useful Pochhammer symbol defined (for $\lambda, \nu \in \mathbb{C}$) by

$$(3) \quad (\lambda)_{\nu} := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1 & (\nu = 0) \\ \lambda(\lambda+1) \cdots (\lambda+n-1) & (\nu = n \in \mathbb{N}), \end{cases}$$

it being understood conventionally that $(0)_0 = 1$, and Γ is the familiar Gamma function.

Motivated mainly by various extensions of the Hurwitz-Lerch Zeta function in one and two variables, we aim to give a further extension of the generalized Hurwitz-Lerch Zeta function of two variables as in (4) and then systematically investigate its several interesting properties such as various integral representations, summation formula, Mellin-Barnes contour integral representations (see [15]), fractional derivatives, and analytic continuation formula. A multi-parameter extension of the extended Hurwitz-Lerch Zeta function of two variables is also introduced. Relevant connections of certain special cases of the main results presented here with some known identities are further pointed out.

2. Extended Hurwitz-Lerch Zeta function of two variables

Here we introduce a further extension of the generalized Hurwitz-Lerch Zeta function of two variables as in (4) and consider some of its special and limiting cases.

An extension

$$(4) \quad \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a) := \sum_{m, n=0}^{\infty} \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (\lambda)_{\rho m} (\mu)_{\sigma m}}{(\gamma)_{\tau m} (\nu)_{\kappa m} m! n!} \frac{z^m t^n}{(m+n+a)^s}$$

($\alpha, \beta, \lambda, \mu \in \mathbb{C}$; $a, \gamma, \nu \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $\delta, \eta, \tau, \rho, \sigma, \kappa \in \mathbb{R}^+$; $\tau - \delta - \eta > -1$, $\kappa - \rho - \sigma > -1$ when $s, z, t \in \mathbb{C}$; $\tau - \delta - \eta = -1$, $\kappa - \rho - \sigma = -1$ and $s \in \mathbb{C}$ when $|z| < \Delta := \delta^{-\delta} \eta^{-\eta} \tau^{\tau}$, $|t| < \Delta^* := \rho^{-\rho} \sigma^{-\sigma} \kappa^{\kappa}$; $\tau - \delta - \eta = -1$, $\kappa - \rho - \sigma = -1$ and $\Re(s + \gamma + \nu - \alpha - \beta - \mu - \lambda) > 0$ when $|z| = \Delta$, $|t| = \Delta^*$).

Special and limiting cases

Case 1. If we set $\alpha = \lambda = \delta = \rho = 1$ in (4), we obtain an extended Hurwitz-Lerch Zeta function of two variables:

$$(5) \quad \begin{aligned} \Phi_{1,\beta;\gamma;1,\mu;\nu}^{(1,\eta,\tau;1,\sigma,\kappa)}(z,t,s,a) &:= \Phi_{\beta;\gamma;\mu;\nu}^{(\eta,\tau;\sigma,\kappa)}(z,t,s,a) \\ &= \sum_{m,n=0}^{\infty} \frac{(\beta)_{\eta m}(\mu)_{\sigma n}}{(\gamma)_{\tau m}(\nu)_{\kappa n}} \frac{z^m t^n}{(m+n+a)^s} \end{aligned}$$

($\beta, \mu \in \mathbb{C}$; $a, \gamma, \nu \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $\eta, \tau, \sigma, \kappa \in \mathbb{R}^+$; $\eta < \tau$, $\sigma < \kappa$ when $s, z, t \in \mathbb{C}$; $\eta = \tau$, $\sigma = \kappa$ and $s \in \mathbb{C}$ when $|z| < \eta^{-\eta} \tau^\tau$, $|t| < \sigma^{-\sigma} \kappa^\kappa$; $\Re(s + \gamma + \nu - \beta - \mu) > 0$ when $|z| = \eta^{-\eta} \tau^\tau$, $|t| = \sigma^{-\sigma} \kappa^\kappa$).

Case 2. The special case of (4) when $\delta = \rho = \eta = \tau = \sigma = \kappa = 1$ yields the known generalized Hurwitz-Lerch Zeta function introduced by Daman and Pathan [4, p. 253, Eq. (10)]:

$$(6) \quad \begin{aligned} \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(1,1,1;1,1,1)}(z,t,s,a) &:= \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}(z,t,s,a) \\ &= \sum_{m,n=0}^{\infty} \frac{(\alpha)_m(\beta)_m(\lambda)_n(\mu)_n}{(\gamma)_m(\nu)_n m! n!} \frac{z^m t^n}{(m+n+a)^s} \end{aligned}$$

($\alpha, \beta, \lambda, \mu \in \mathbb{C}$; $\gamma, \nu, a \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $s \in \mathbb{C}$ when $|z| < 1$; $\Re(s + \gamma + \nu - \alpha - \beta - \mu - \lambda) > 0$ when $|z| = 1$ and $|t| = 1$).

Case 3. The special case of (4) when $\eta = \tau$, $\sigma = \kappa$ and $\delta = \rho = 1$ with $\gamma = \beta$ and $\mu = \nu$ reduces to yield another known generalized Hurwitz-Lerch Zeta function studied by Pathan and Daman [16]:

$$(7) \quad \Phi_{\alpha,\beta;\beta;\lambda,\nu;\nu}^{(1,1,1;1,1,1)}(z,t,s,a) := \Phi_{\alpha;\lambda}^*(z,t,s,a) = \sum_{m,n=0}^{\infty} \frac{(\alpha)_m(\lambda)_n}{m! n!} \frac{z^m t^n}{(m+n+a)^s}$$

($\mu, \lambda \in \mathbb{C}$; $a \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $s \in \mathbb{C}$ when $|z| < 1$ and $|t| < 1$; $\Re(s - \mu - \lambda) > 0$ when $|z| = 1$ and $|t| = 1$).

Case 4. We use the following limiting case of the extended Hurwitz-Lerch Zeta function (4):

$$(8) \quad \begin{aligned} \Phi_{\beta;\gamma;\mu;\nu}^*(\eta,\sigma;\tau,\kappa)(z,s,a) &= \lim_{\min\{|\alpha|,|\lambda|\} \rightarrow \infty} \left\{ \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)} \left(\frac{z}{\alpha^\delta}, \frac{t}{\lambda^\rho}, s, a \right) \right\} \\ &:= \sum_{m,n=0}^{\infty} \frac{(\beta)_{\eta m}(\mu)_{\sigma n}}{(\gamma)_{\tau m}(\nu)_{\kappa n} m! n!} \frac{z^m t^n}{(m+n+a)^s} \end{aligned}$$

($\beta, \mu \in \mathbb{C}$; $a, \gamma, \nu \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $\eta, \tau, \sigma, \kappa \in \mathbb{R}^+$; $s \in \mathbb{C}$ when $|z| < \eta^{-\eta} \tau^\tau$, $|t| < \sigma^{-\sigma} \kappa^\kappa$; $\Re(s + \gamma + \nu - \beta - \mu) > 0$ when $|z| = \eta^{-\eta} \tau^\tau$, $|t| = \sigma^{-\sigma} \kappa^\kappa$).

Case 5. Another limiting case of the extended Hurwitz-Lerch Zeta function (4) is given as follows:

$$\Phi_{\beta;\mu}^*(\eta;\sigma)(z,t,s,a) := \lim_{\min\{|\alpha|,|\gamma|,|\lambda|,|\nu|\} \rightarrow \infty} \left\{ \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)} \left(\frac{z\gamma^\tau}{\beta^\eta}, \frac{t\nu^\kappa}{\lambda^\rho}, s, a \right) \right\}$$

$$(9) \quad := \sum_{m,n=0}^{\infty} \frac{(\beta)_{\eta m} (\mu)_{\sigma n}}{m!n!} \frac{z^m t^n}{(m+n+a)^s}$$

($\beta, \mu \in \mathbb{C}$; $a \in \mathbb{C} \setminus \mathbb{Z}_0^-$; $0 < \eta < 1$, $0 < \sigma < 1$ and $s, z, t \in \mathbb{C}$; $\eta = 1$, $\sigma = 1$ and $s \in \mathbb{C}$ when $|z| < \eta^{-\eta}$, $|t| < \sigma^{-\sigma}$ and $\Re(s - \beta - \mu) > 0$ when $|z| = \eta^{-\eta}$, $|t| = \sigma^{-\sigma}$).

3. Integral Representations of $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a)$

We first recall the Fox-Wright function ${}_p\Psi_q(z)$ ($p, q \in \mathbb{N}_0$) or its normalization ${}_p\Psi_q^*$ ($p, q \in \mathbb{N}_0$) with p numerator and q denominator parameters defined for $\alpha_1, \dots, \alpha_p \in \mathbb{C}$ and $\beta_1, \dots, \beta_q \in \mathbb{C} \setminus \mathbb{Z}_0^-$ by (see, for details, [5, 10, 12–14]; see also [17, 22, 24]):

$$(10) \quad \begin{aligned} & {}_p\Psi_q^* \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_p, A_p); \\ (\beta_1, B_1), \dots, (\beta_q, B_q); \end{matrix} z \right] \\ & := \sum_{n=0}^{\infty} \frac{(\alpha_1)_{A_1 n} \cdots (\alpha_p)_{A_p n}}{(\beta_1)_{B_1 n} \cdots (\beta_q)_{B_q n}} \frac{z^n}{n!} \\ & = \frac{\Gamma(\beta_1) \cdots \Gamma(\beta_q)}{\Gamma(\alpha_1) \cdots \Gamma(\alpha_p)} {}_p\Psi_q \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_p, A_p); \\ (\beta_1, B_1), \dots, (\beta_q, B_q); \end{matrix} z \right] \\ & \left(A_j \in \mathbb{R}^+ (j = 1, \dots, p); B_j \in \mathbb{R}^+ (j = 1, \dots, q); 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j \geq 0 \right), \end{aligned}$$

where the equality in the convergence condition holds true for

$$|z| < \nabla := \left(\prod_{j=1}^p A_j^{-A_j} \right) \cdot \left(\prod_{j=1}^q B_j^{B_j} \right).$$

In particular, when $A_j = B_k = 1$ ($j = 1, \dots, p$; $k = 1, \dots, q$), (10) reduces immediately to the generalized hypergeometric function ${}_pF_q$ ($p, q \in \mathbb{N}_0$) (see, e.g., [22]):

$$(11) \quad \begin{aligned} & {}_p\Psi_q^* \left[\begin{matrix} (\alpha_1, 1), \dots, (\alpha_p, 1); \\ (\beta_1, 1), \dots, (\beta_q, 1); \end{matrix} z \right] = {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} z \right] \\ & = \frac{\Gamma(\beta_1) \cdots \Gamma(\beta_q)}{\Gamma(\alpha_1) \cdots \Gamma(\alpha_p)} {}_p\Psi_q \left[\begin{matrix} (\alpha_1, 1), \dots, (\alpha_p, 1); \\ (\beta_1, 1), \dots, (\beta_q, 1); \end{matrix} z \right]. \end{aligned}$$

Theorem 3.1. *The following integral representation for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a)$ in (4) holds true:*

$$(12) \quad \begin{aligned} \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a) & = \frac{1}{\Gamma(s)} \int_0^{\infty} x^{s-1} e^{-ax} {}_2\Psi_1^* \left[\begin{matrix} (\alpha, \delta), (\beta, \eta); \\ (\gamma, \tau); \end{matrix} ze^{-x} \right] \\ & \quad \times {}_2\Psi_1^* \left[\begin{matrix} (\lambda, \rho), (\mu, \sigma); \\ (\nu, \kappa); \end{matrix} te^{-x} \right] dx \end{aligned}$$

($\min\{\Re(s), \Re(a)\} > 0$, $\delta, \eta, \tau, \rho, \sigma, \kappa \in \mathbb{R}^+$ and $\tau - \delta - \eta \geq -1$, $\kappa - \rho - \sigma \geq -1$ when $|z| < \Delta := \delta^{-\delta} \eta^{-\eta} \tau^\tau$, $|t| < \Delta^* := \rho^{-\rho} \sigma^{-\sigma} \kappa^\kappa$).

Proof. Using the following Eulerian integral

$$\frac{1}{(m+n+a)^s} := \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-(m+n+a)t} dt$$

($\min\{\Re(s), \Re(a)\} > 0$; $m, n \in \mathbb{N}_0$)

in (4), and interchanging the order of summation and integration which is guaranteed under the condition stated in Theorem 3.1, and using (10), we are led to the desired integral representation. \square

If, in Theorem 3.1, we set $\alpha = \lambda = \delta = \rho = 1$ and apply the relationship (5), we obtain an integral representation for the function $\Phi_{\beta; \gamma; \mu; \nu}^{(\eta; \tau; \sigma; \kappa)}(z, t, s, a)$ asserted by Corollary 3.2.

Corollary 3.2. *The following integral representation for the function $\Phi_{\beta; \gamma; \mu; \nu}^{(\eta; \tau; \sigma; \kappa)}(z, t, s, a)$ in (5) holds true:*

$$(13) \quad \Phi_{\beta; \gamma; \mu; \nu}^{(\eta; \tau; \sigma; \kappa)}(z, t, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_2\Psi_1^* \left[\begin{matrix} (1, 1), (\beta, \eta); \\ (\gamma, \tau); \end{matrix} ; ze^{-x} \right] \\ \times {}_2\Psi_1^* \left[\begin{matrix} (1, 1), (\mu, \sigma); \\ (\nu, \kappa); \end{matrix} ; te^{-x} \right] dx$$

($\min\{\Re(s), \Re(a)\} > 0$, $\tau > \eta > 0$, $\kappa > \sigma > 0$ when $z, t \in \mathbb{C}$; $\tau \geq \eta > 0$, $\kappa \geq \sigma > 0$ when $|z| = \eta^{-\eta} \tau^\tau$, $|t| < \rho^{-\rho} \kappa^\kappa$).

Remark 3.3. The special case $\delta = \rho = \eta = \tau = \sigma = \kappa = 1$ of the result in Theorem 3.1 yields the following known integral representation [4, p. 253, Eq. (11)]:

$$(14) \quad \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_2F_1(\alpha, \beta; \gamma; ze^{-x}) {}_2F_1(\lambda, \mu; \nu; te^{-x}) dx$$

($\Re(a) > 0$; $\Re(s) > 0$ when $|z| \leq 1$ ($z \neq 1$), $|t| \leq 1$ ($t \neq 1$); $\Re(s) > 1$ when $z = 1, t = 1$), which, upon applying Euler's transformation formula (see, e.g., [5, p. 64, Eq. 2.1.4 (23)]):

$${}_2F_1(a, b; c; z) = (1-z)^{c-a-b} {}_2F_1(c-a, c-b; c; z) \quad (|\arg(1-z)| < \pi),$$

gives another integral representation for the function (6):

$$(15) \quad \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{(1-ze^{-x})^{\alpha+\beta-\gamma} (1-te^{-x})^{\lambda+\mu-\nu}} \\ {}_2F_1(\gamma-\alpha, \gamma-\beta; \gamma; ze^{-x}) {}_2F_1(\nu-\lambda, \nu-\mu; \nu; te^{-x}) dx$$

($\Re(a) > 0$; $\Re(s) > 0$ when $|z| \leq 1$ ($z \neq 1$), $|t| \leq 1$ ($t \neq 1$); $\Re(s) > 1$ when $z = 1$, $t = 1$). The special case of (15) when $\beta = \gamma$ and $\mu = \nu$ immediately reduces to the following known result for the function (7) (see [4, p. 252, Eq. (7)]):

$$(16) \quad \Phi_{\alpha;\lambda}^*(z, t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{(1 - ze^{-x})^\alpha (1 - te^{-x})^\lambda} dx$$

($\Re(a) > 0$; $\Re(s) > 0$ when $|z| \leq 1$ ($z \neq 1$), $|t| \leq 1$ ($t \neq 1$); $\Re(s) > 1$ when $z = 1$, $t = 1$), which, upon setting $\alpha = \lambda = 1$, yields the following integral representation for the function:

$$(17) \quad \Phi(z, t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{(1 - ze^{-x})(1 - te^{-x})} dx$$

($\Re(a) > 0$; $\Re(s) > 0$ when $|z| \leq 1$ ($z \neq 1$), $|t| \leq 1$ ($t \neq 1$); $\Re(s) > 1$ when $z = 1$, $t = 1$).

Remark 3.4. It is interesting to see that the special cases for $z = 0$ or $t = 0$ in (12) to (17) yield the known integral representations earlier studied by Srivastava et al. [25, p. 494, Eq. (2.4)], Lin and Srivastava [11, p. 728, Eq. (14)], Garg et al. [6, p. 313, Eq. (2.1)], Goyal and Laddha [7, p. 100, Eq. (1.6)] and the familiar Hurwitz-Lerch Zeta function [5, p. 27, Eq. 1.11(3)] which are, respectively, given as follows:

$$(18) \quad \Phi_{\lambda,\mu;\nu}^{(\rho,\sigma,\kappa)}(t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_2\Psi_1^* \left[\begin{matrix} (\lambda, \rho), (\mu, \sigma); \\ (\nu, \kappa); \end{matrix} te^{-x} \right] dx$$

$$(\min\{\Re(s), \Re(a)\} > 0, \rho, \sigma, \kappa \in \mathbb{R}^+ \text{ and } \kappa - \rho - \sigma \geq -1 \\ \text{when } |t| < \delta^* := \rho^{-\rho} \sigma^{-\sigma} \kappa^\kappa);$$

$$(19) \quad \Phi_{(\mu;\nu)}^{(\rho,\kappa)}(t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_2\Psi_1^* \left[\begin{matrix} (1, 1), (\mu, \rho); \\ (\nu, \kappa); \end{matrix} te^{-x} \right] dx$$

$$(\min\{\Re(s), \Re(a)\} > 0, \kappa > \rho > 0 \text{ when } t \in \mathbb{C}; \kappa \geq \rho > 0 \text{ when } |t| < \rho^{-\rho} \kappa^\kappa);$$

$$(20) \quad \Phi_{\lambda,\mu;\nu}(t, s, a) := \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_2F_1(\lambda, \mu; \nu; te^{-x}) dx$$

$$(\Re(a) > 0; \Re(s) > 0 \text{ when } |t| \leq 1 \text{ } (t \neq 1); \Re(s) > 1 \text{ when } t = 1);$$

$$(21) \quad \Phi_\mu^*(t, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{(1 - te^{-x})^\mu} dx$$

$$(\Re(a) > 0; \Re(s) > 0 \text{ when } |t| \leq 1 \text{ } (t \neq 1); \Re(s) > 1 \text{ when } t = 1);$$

$$(22) \quad \Phi(t, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-ax}}{1 - te^{-x}} dx$$

$$(\Re(a) > 0; \Re(s) > 0 \text{ when } |t| \leq 1 \text{ } (t \neq 1); \Re(s) > 1 \text{ when } t = 1).$$

Theorem 3.5. *The following double and triple integral representations for $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$ in (4) hold true:*

$$(23) \quad \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a) = \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \frac{w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\ \times \Phi_{\beta,\mu;\gamma-\alpha,\nu-\lambda}^{*(\eta,\sigma;\tau-\delta,\kappa-\rho)}\left(\frac{zw^\delta}{(1+w)^\tau}, \frac{ty^\rho}{(1+y)^\kappa}, s, a\right) dw dy$$

($\Re(\gamma) > \Re(\alpha) > 0$, $\Re(\nu) > \Re(\lambda) > 0$; $\tau \geq \delta > 0$, $\kappa \geq \rho > 0$; $\eta, \sigma \in \mathbb{R}^+$) and

$$(24) \quad \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a) \\ = \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(s)\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \int_0^\infty \frac{x^{s-1}e^{-ax}w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\ \times {}_1\Psi_1^*\left[\begin{matrix} (\beta, \eta); \\ (\gamma-\alpha, \tau-\delta); \end{matrix} \frac{ze^{-x}w^\delta}{(1+w)^\tau}\right] \\ \times {}_1\Psi_1^*\left[\begin{matrix} (\mu, \sigma); \\ (\nu-\lambda, \kappa-\rho); \end{matrix} \frac{te^{-x}y^\rho}{(1+y)^\kappa}\right] dx dw dy$$

($\Re(\gamma) > \Re(\alpha) > 0$, $\Re(\nu) > \Re(\lambda) > 0$; $\tau \geq \delta > 0$, $\kappa \geq \rho > 0$; $\eta, \sigma \in \mathbb{R}^+$; $\min\{\Re(s), \Re(a)\} > 0$).

Proof. Setting $a = \alpha + \delta m$ and $b = \gamma + \tau m$, and $a = \lambda + \rho n$ and $b = \nu + \kappa n$, respectively, in the Eulerian Beta-function formula [5, p. 9, Eq. 1.5(2)]:

$$(25) \quad B(a, b-a) = \frac{\Gamma(a)\Gamma(b-a)}{\Gamma(b)} = \int_0^\infty \frac{y^{a-1}}{(1+y)^b} dy \quad (\Re(b) > \Re(a) > 0),$$

we find

$$(26) \quad \frac{(\alpha)_{\delta m}}{(\gamma)_{\tau m}} = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \frac{1}{(\gamma-\alpha)_{(\tau-\delta)m}} \int_0^\infty \frac{w^{\alpha+\delta m-1}}{(1+w)^{\gamma+\tau m}} dw \\ (\Re(\gamma) > \Re(\alpha) > 0; \tau \geq \delta; m \in \mathbb{N}_0)$$

and

$$(27) \quad \frac{(\lambda)_{\rho n}}{(\nu)_{\kappa n}} = \frac{\Gamma(\nu)}{\Gamma(\lambda)\Gamma(\nu-\lambda)} \frac{1}{(\nu-\lambda)_{(\kappa-\rho)n}} \int_0^\infty \frac{y^{\lambda+\rho n-1}}{(1+y)^{\nu+\kappa n}} dy \\ (\Re(\nu) > \Re(\lambda) > 0; \kappa \geq \rho; n \in \mathbb{N}_0),$$

which, by appealing to the definition (8), immediately yields the assertion (23).

Moreover, by (12), (25) and (26), we also obtain

$$\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a) \\ := \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} \sum_{m=0}^\infty \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (ze^{-x})^m}{(\gamma)_{\tau m} m!} \sum_{n=0}^\infty \frac{(\lambda)_{\rho n} (\mu)_{\sigma n} (te^{-x})^n}{(\nu)_{\kappa n} n!} dx$$

$$\begin{aligned}
&= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(s)\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \int_0^\infty \frac{x^{s-1}e^{-ax}w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\
&\quad \times \sum_{m=0}^\infty \frac{(\beta)_{\eta m}}{(\gamma-\alpha)_{(\tau-\delta)m}m!} \left(\frac{ze^{-x}w^\delta}{(1+w)^\tau} \right)^m \\
&\quad \times \sum_{n=0}^\infty \frac{(\mu)_{\sigma n}}{(\nu-\lambda)_{(\kappa-\rho)n}n!} \left(\frac{te^{-x}y^\rho}{(1+y)^\kappa} \right)^n dx dw dy,
\end{aligned}$$

which, in view of (10), leads us to the second assertion (24) of Theorem 3.5. \square

If, in Theorem 3.5, we set $\tau = \delta$ and $\kappa = \rho$ and using the definition (9) and (10), we obtain the following special cases for the function $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\delta;\rho,\sigma,\rho)}(z,t,s,a)$, respectively, asserted by Corollary 3.6 below.

Corollary 3.6. *Each of the following integral representations for $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\delta;\rho,\sigma,\rho)}(z,t,s,a)$ holds true:*

$$\begin{aligned}
(28) \quad &\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\delta;\rho,\sigma,\rho)}(z,t,s,a) \\
&= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \frac{w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\
&\quad \times \Phi_{\beta,\mu}^{*(\eta,\sigma)} \left(\frac{zx^\delta}{(1+y)^\delta}, \frac{ty^\rho}{(1+y)^\rho}, s, a \right) dw dy \\
&\quad (\Re(\gamma) > \Re(\alpha) > 0, \Re(\nu) > \Re(\lambda) > 0; \eta, \sigma \in \mathbb{R}^+)
\end{aligned}$$

and

$$\begin{aligned}
(29) \quad &\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\delta;\rho,\sigma,\rho)}(z,t,s,a) \\
&= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(s)\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \int_0^\infty \frac{x^{s-1}e^{-ax}w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\
&\quad \times {}_1\Psi_0^* \left[\begin{matrix} (\beta, \eta) \\ - \end{matrix}; \frac{ze^{-x}w^\delta}{(1+w)^\delta} \right] {}_1\Psi_0^* \left[\begin{matrix} (\mu, \sigma) \\ - \end{matrix}; \frac{te^{-x}y^\rho}{(1+y)^\rho} \right] dx dw dy \\
&\quad (\Re(\gamma) > \Re(\alpha) > 0, \Re(\nu) > \Re(\lambda) > 0; \eta, \sigma \in \mathbb{R}^+; \min\{\Re(s), \Re(a)\} > 0).
\end{aligned}$$

Further, if we set $\delta = \eta = \rho = \sigma = 1$ in (29), we obtain a new integral representation for $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}(z,t,s,a)$ in (6):

$$\begin{aligned}
(30) \quad &\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}(z,t,s,a) \\
&= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(s)\Gamma(\alpha)\Gamma(\gamma-\alpha)\Gamma(\lambda)\Gamma(\nu-\lambda)} \int_0^\infty \int_0^\infty \int_0^\infty \frac{x^{s-1}e^{-ax}w^{\alpha-1}y^{\lambda-1}}{(1+w)^\gamma(1+y)^\nu} \\
&\quad \times \left(1 - \frac{zwe^{-x}}{1+w} \right)^{-\beta} \left(1 - \frac{tye^{-x}}{1+y} \right)^{-\mu} dx dw dy \\
&\quad (\Re(\gamma) > \Re(\alpha) > 0, \Re(\nu) > \Re(\lambda) > 0; \min\{\Re(s), \Re(a)\} > 0).
\end{aligned}$$

Theorem 3.7. *The following summation formula for $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$ in (4) holds true:*

$$(31) \quad \sum_{k=0}^{\infty} \frac{(s)_k}{k!} \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s+k, a) x^n = \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a-x)$$

$$(\lambda \in \mathbb{C}; |x| < |a|; s \neq 1).$$

Proof. Using (4) in the right-hand side of the assertion (31), we have

$$(32) \quad \begin{aligned} & \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a-x) \\ &= \sum_{m,n=0}^{\infty} \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (\lambda)_{\rho n} (\mu)_{\sigma n}}{(\gamma)_{\tau m} (\nu)_{\kappa n}} \frac{z^m t^n}{m!n!(m+n+a-x)^s} \\ &= \sum_{m,n=0}^{\infty} \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (\lambda)_{\rho n} (\mu)_{\sigma n}}{(\gamma)_{\tau m} (\nu)_{\kappa n}} \frac{z^m t^n}{m!n!(m+n+a)^s} \left(1 - \frac{x}{m+n+a}\right)^{-s} \\ &= \sum_{m,n=0}^{\infty} \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (\lambda)_{\rho n} (\mu)_{\sigma n}}{(\gamma)_{\tau m} (\nu)_{\kappa n}} \frac{z^m t^n}{m!n!(m+n+a)^s} \left\{ \sum_{k=0}^{\infty} \frac{(s)_k}{k!} \frac{x^k}{(m+n+a)^k} \right\} \\ &= \sum_{k=0}^{\infty} \frac{(s)_k}{k!} \left\{ \sum_{m,n=0}^{\infty} \frac{(\alpha)_{\delta m} (\beta)_{\eta m} (\lambda)_{\rho n} (\mu)_{\sigma n}}{(\gamma)_{\tau m} (\nu)_{\kappa n}} \frac{z^m t^n}{m!n!(m+n+a)^{s+k}} \right\} x^k, \end{aligned}$$

which, upon using (4), leads to the right-hand side of (31). \square

4. Mellin-Barnes contour integral representations

We begin by presenting the following Mellin-Barnes contour integral representation of the extended Hurwitz-Lerch Zeta function $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$ of two variables as in Theorem 4.1.

Theorem 4.1. *The following Mellin-Barnes integral representation for $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$ in (4) holds true:*

$$(33) \quad \begin{aligned} & \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a) \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \frac{1}{(2\pi i)^2} \int_{-i\infty}^{+i\infty} \int_{-i\infty}^{+i\infty} \Gamma(-\xi)\Gamma(-\zeta) \\ & \quad \times \frac{\Gamma(\alpha + \delta\xi)\Gamma(\beta + \eta\xi)\Gamma(\lambda + \rho\zeta)\Gamma(\mu + \sigma\zeta)(-z)^\xi(-t)^\zeta}{\Gamma(\gamma + \tau\xi)\Gamma(\nu + \kappa\zeta)(\xi + \zeta + a)^s} d\xi d\zeta \end{aligned}$$

$$(|\arg(-z)| < \pi, |\arg(-t)| < \pi, \min\{\Re(a), \Re(s), \Re(\gamma), \Re(\nu)\} > 0),$$

where it is assumed that the poles of the integrand in (33) are simple. The contours of integration are so described that the poles of $\Gamma(-\xi)$, $\Gamma(-\zeta)$ are

separated from the poles of $\Gamma(\alpha + \delta\xi)$, $\Gamma(\beta + \eta\xi)$, $\Gamma(\lambda + \rho\zeta)$ and $\Gamma(\mu + \sigma\zeta)$ with indentations, if necessary.

Proof. If we calculate the integral (33) as sum of the residues at the simple poles of $\Gamma(-\xi)$ at the points $\xi = m$ ($m \in \mathbb{N}_0$) and of the $\Gamma(-\zeta)$ ($n \in \mathbb{N}_0$) at the points $\zeta = n$, respectively, we immediately obtain the following series expansion:

$$(34) \quad \begin{aligned} & \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a) \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \\ & \times \sum_{m, n=0}^{\infty} \frac{\Gamma(\alpha + \delta m)\Gamma(\beta + \eta m)\Gamma(\lambda + \rho n)\Gamma(\mu + \sigma n)}{\Gamma(\gamma + \tau m)\Gamma(\nu + \kappa n)m!n!} \frac{z^m t^n}{(m + n + a)^s} \end{aligned}$$

for the extended Hurwitz-Lerch Zeta function of two variables. This completes the proof. \square

Suitably specializing the Mellin-Barnes integral representation in (33), we can obtain Mellin-Barnes contour integral representations of the extended Hurwitz-Lerch Zeta functions in (5), (7), (8) and (9), whose explicit expressions are left to the interested reader. Here we consider the following special case (33) when $\delta = \rho = \eta = \tau = \sigma = \kappa = 1$ which is a Mellin-Barnes contour integral representation of the Hurwitz-Lerch Zeta function of two variables in (6) as in Corollary 4.2.

Corollary 4.2. *The following contour integral representations for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a)$ in (6) holds true:*

$$(35) \quad \begin{aligned} \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a) &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} \frac{1}{(2\pi i)^2} \int_{-i\infty}^{+i\infty} \int_{-i\infty}^{+i\infty} \Gamma(-\xi)\Gamma(-\zeta) \\ & \times \frac{\Gamma(\alpha + \xi)\Gamma(\beta + \xi)\Gamma(\lambda + \zeta)\Gamma(\mu + \zeta)(-z)^\xi (-t)^\zeta}{\Gamma(\gamma + \xi)\Gamma(\nu + \zeta)(\xi + \zeta + a)^s} d\xi d\zeta. \end{aligned}$$

5. Computational Representation for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a)$

Here we investigate an analytic continuation for the extended Hurwitz-Lerch function of two variables in (4). This result will enable us to derive corresponding continuation formulas for most of members of the extended Hurwitz-Lerch Zeta functions of two variables and its related functions.

Theorem 5.1. *The following computational representation for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a)$ holds true:*

$$\begin{aligned} \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a) &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} (-z)^{-\frac{\alpha}{\delta}} (-t)^{-\frac{\lambda}{\rho}} \\ & \times \sum_{m, n=0}^{\infty} \frac{\Gamma(\frac{\alpha+m}{\delta})\Gamma(\frac{\lambda+n}{\rho})\Gamma(\beta-\eta(\frac{\alpha+m}{\delta}))\Gamma(\mu-\sigma(\frac{\lambda+n}{\rho}))(-z)^{-\frac{1}{\delta}m}(-t)^{-\frac{1}{\rho}n}}{\Gamma(\gamma-\tau(\frac{\alpha+m}{\delta}))\Gamma(\nu-\kappa(\frac{\lambda+n}{\rho}))\Gamma(a-(\frac{\alpha+m}{\delta})-(\frac{\lambda+n}{\rho}))^s m!n!} \end{aligned}$$

$$(36) \quad + \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} (-z)^{-\frac{\beta}{\eta}} (-t)^{-\frac{\mu}{\sigma}}$$

$$\times \sum_{m,n=0}^{\infty} \frac{\Gamma(\frac{\beta+m}{\eta})\Gamma(\frac{\mu+n}{\sigma})\Gamma(\alpha-\delta(\frac{\beta+m}{\eta}))\Gamma(\lambda-\rho(\frac{\mu+n}{\sigma}))(-(-z)^{-\frac{1}{\eta}})^m (-(-t)^{-\frac{1}{\sigma}})^n}{\Gamma(\gamma-\tau(\frac{\beta+m}{\eta}))\Gamma(\nu-\kappa(\frac{\mu+n}{\sigma}))\left(a-\left(\frac{\beta+m}{\eta}\right)-\left(\frac{\mu+n}{\sigma}\right)\right)^s m!n!}$$

$$(\min\{\Re(a), \Re(s)\} > 0, \max\{|\arg(-z)|, |\arg(-t)|\} < \pi, |x| \rightarrow \infty, |t| \rightarrow \infty).$$

Proof. To obtain the analytic continuation formula for the extended Hurwitz-Lerch Zeta function in (33), we first calculate the residue of the $\Gamma(\alpha + \delta\xi)$ at the simple poles:

$$\xi = -\frac{\alpha + m}{\delta} \quad (m \in \mathbb{N}_0)$$

and the residues of $\Gamma(\lambda + \rho\zeta)$ at the simple poles:

$$\zeta = -\frac{\lambda + n}{\rho} \quad (n \in \mathbb{N}_0)$$

as (say I_1)

$$(37) \quad I_1 = \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)}$$

$$\times \sum_{m,n=0}^{\infty} \frac{\Gamma(\frac{\alpha+m}{\delta})\Gamma(\frac{\lambda+n}{\rho})\Gamma(\beta-\eta(\frac{\alpha+m}{\delta}))\Gamma(\mu-\sigma(\frac{\lambda+n}{\rho}))(-z)^{-\left(\frac{\alpha+m}{\delta}\right)} (-t)^{-\left(\frac{\lambda+n}{\rho}\right)} (-1)^{m+n}}{\Gamma(\gamma-\tau(\frac{\alpha+m}{\delta}))\Gamma(\nu-\kappa(\frac{\lambda+n}{\rho}))\left(a-\left(\frac{\alpha+m}{\delta}\right)-\left(\frac{\lambda+n}{\rho}\right)\right)^s m!n!}$$

$$= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} (-z)^{-\frac{\alpha}{\delta}} (-t)^{-\frac{\lambda}{\rho}}$$

$$\times \sum_{m,n=0}^{\infty} \frac{\Gamma(\frac{\alpha+m}{\delta})\Gamma(\frac{\lambda+n}{\rho})\Gamma(\beta-\eta(\frac{\alpha+m}{\delta}))\Gamma(\mu-\sigma(\frac{\lambda+n}{\rho}))(-(-z)^{-\frac{1}{\delta}})^m (-(-t)^{-\frac{1}{\rho}})^n}{\Gamma(\gamma-\tau(\frac{\alpha+m}{\delta}))\Gamma(\nu-\kappa(\frac{\lambda+n}{\rho}))\left(a-\left(\frac{\alpha+m}{\delta}\right)-\left(\frac{\lambda+n}{\rho}\right)\right)^s m!n!}.$$

Similarly, the residues of the $\Gamma(\beta + \eta\xi)$ at the simple poles:

$$\xi = -\frac{\beta + m}{\eta} \quad (m \in \mathbb{N}_0)$$

and the residues of $\Gamma(\mu + \sigma\zeta)$ at the simple poles:

$$\zeta = -\frac{\mu + n}{\sigma} \quad (n \in \mathbb{N}_0)$$

as (say I_2)

$$(38) \quad I_2 = \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)} (-z)^{-\frac{\beta}{\eta}} (-t)^{-\frac{\mu}{\sigma}}$$

$$\times \sum_{m,n=0}^{\infty} \frac{\Gamma(\frac{\beta+m}{\eta})\Gamma(\frac{\mu+n}{\sigma})\Gamma(\alpha-\delta(\frac{\beta+m}{\eta}))\Gamma(\lambda-\rho(\frac{\mu+n}{\sigma}))\left(-(-z)^{-\frac{1}{\eta}}\right)^m \left(-(-t)^{-\frac{1}{\sigma}}\right)^n}{\Gamma(\gamma-\tau(\frac{\beta+m}{\eta}))\Gamma(\nu-\kappa(\frac{\mu+n}{\sigma}))m!n! \left(a-\left(\frac{\beta+m}{\eta}\right)-\left(\frac{\mu+n}{\sigma}\right)\right)^s}.$$

Thus, by combining the residues (37) and (38), we are led to the representation (36). \square

Corollary 5.2. *Under the hypothesis of Theorem 5.1, the following asymptotic formula holds true:*

$$(39) \quad \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a) \sim A(-z)^{-\frac{\alpha}{\delta}}(-t)^{-\frac{\lambda}{\rho}} + B(-z)^{-\frac{\beta}{\eta}}(-t)^{-\frac{\mu}{\sigma}},$$

where

$$A := \frac{\Gamma(\gamma)\Gamma(\nu)\Gamma(\frac{\alpha}{\delta})\Gamma(\frac{\lambda}{\rho})\Gamma(\beta - \eta(\frac{\alpha}{\delta}))\Gamma(\mu - \sigma(\frac{\lambda}{\rho}))}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)\Gamma(\gamma - \tau(\frac{\alpha}{\delta}))\Gamma(\nu - \kappa(\frac{\lambda}{\rho}))\left(a - (\frac{\alpha}{\delta}) - (\frac{\lambda}{\rho})\right)^s}$$

and

$$B := \frac{\Gamma(\gamma)\Gamma(\nu)\Gamma(\frac{\beta}{\eta})\Gamma(\frac{\mu}{\sigma})\Gamma\left(\alpha - \delta(\frac{\beta}{\eta})\right)\Gamma(\lambda - \rho(\frac{\mu}{\sigma}))}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)\Gamma(\gamma - \tau(\frac{\beta}{\eta}))\Gamma(\nu - \kappa(\frac{\mu}{\sigma}))\left(a - (\frac{\beta}{\eta}) - (\frac{\mu}{\sigma})\right)^s}$$

$$(\min\{\Re(\alpha), \Re(\beta), \Re(\lambda), \Re(\mu), \Re(a), \Re(s)\} > 0,$$

$$\max\{|\arg(-z)|, |\arg(-t)|\} < \pi; |x| \rightarrow \infty, |t| \rightarrow \infty).$$

Analytic continuation formulas of the extended Hurwitz-Lerch Zeta functions in (5), (7), (8) and (9) can be easily obtained as suitable special cases of the analytic continuation formula in (36). In particular, the special case of (36) when $\eta = \tau = \sigma = \kappa = 1$ reduces to yield the following representation for the Hurwitz-Lerch Zeta function of two variable due to Daman and Pathan [4, p. 253, Eq. (11)] as in Corollary 5.3.

Corollary 5.3. *The following computational representation for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a)$ holds true:*

$$(40) \quad \begin{aligned} & \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}(z, t, s, a) \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)}(-z)^{-\alpha}(-t)^{-\lambda} \\ & \quad \times \sum_{m, n=0}^{\infty} \frac{\Gamma(\alpha+m)\Gamma(\lambda+n)\Gamma(\beta-(\alpha+m))\Gamma(\mu-(\lambda+n))(-z)^{-m}(-t)^{-n}}{\Gamma(\gamma-(\alpha+m))\Gamma(\nu-(\lambda+n))(a-(\alpha+m)-(\lambda+n))^s m! n!} \\ & \quad + \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\alpha)\Gamma(\beta)\Gamma(\lambda)\Gamma(\mu)}(-z)^{-\beta}(-t)^{-\mu} \\ & \quad \times \sum_{m, n=0}^{\infty} \frac{\Gamma(\beta+m)\Gamma(\mu+n)\Gamma(\alpha-(\beta+m))\Gamma(\lambda-(\mu+n))(-z)^{-m}(-t)^{-n}}{\Gamma(\gamma-(\beta+m))\Gamma(\nu-(\mu+n))(a-(\beta+m)-(\mu+n))^s m! n!} \end{aligned}$$

$$(\min\{\Re(a), \Re(s)\} > 0, \max\{|\arg(-z)|, |\arg(-t)|\} < \pi; |x| \rightarrow \infty, |t| \rightarrow \infty).$$

Furthermore, by letting $s \rightarrow 0$ in (40), and using the formula:

$$(\alpha)_{-n} = \frac{(-1)^n}{(1-\alpha)_n} \quad (\alpha \in \mathbb{C} \setminus \mathbb{N}; n \in \mathbb{N}_0),$$

we obtain the following interesting continuation formula for the product of two Gauss hypergeometric functions:

$$(41) \quad \begin{aligned} & {}_2F_1(\alpha, \beta; \gamma; z) {}_2F_1(\lambda, \mu; \nu; t) \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)\Gamma(\beta-\alpha)\Gamma(\mu-\lambda)}{\Gamma(\beta)\Gamma(\mu)\Gamma(\gamma-\alpha)\Gamma(\nu-\lambda)} (-z)^{-\alpha} (-t)^{-\lambda} \\ & \quad \times {}_2F_1\left(\alpha, 1+\alpha-\gamma; 1+\alpha-\beta; \frac{1}{z}\right) {}_2F_1\left(\lambda, 1+\lambda-\nu; 1+\lambda-\mu; \frac{1}{t}\right) \\ & \quad + \frac{\Gamma(\gamma)\Gamma(\nu)\Gamma(\alpha-\beta)\Gamma(\lambda-\mu)}{\Gamma(\beta)\Gamma(\mu)\Gamma(\gamma-\beta)\Gamma(\nu-\mu)} (-z)^{-\beta} (-t)^{-\mu} \\ & \quad \times {}_2F_1\left(\beta, 1+\beta-\gamma; 1+\beta-\alpha; \frac{1}{z}\right) {}_2F_1\left(\mu, 1+\mu-\nu; 1+\mu-\lambda; \frac{1}{t}\right) \\ & \quad (\max\{|\arg(-z)|, |\arg(-t)|\} < \pi; \min\{|z|, |t|\} > 1 \text{ when} \\ & \quad \min\{\Re(\gamma-\alpha-\beta), \Re(\nu-\lambda-\mu)\} > 0). \end{aligned}$$

6. Fractional Derivatives for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, t, s, a)$

Recall the *Riemann-Liouville fractional derivative operator* \mathcal{D}_z^μ defined by (see, e.g. [10] and [17, p. 70 et seq.]):

$$(42) \quad \mathcal{D}_z^\mu \{f(z)\} := \begin{cases} \frac{1}{\Gamma(-\mu)} \int_0^z (z-t)^{-\mu-1} f(t) dt & (\Re(\mu) < 0), \\ \frac{d^m}{dz^m} \left\{ \mathcal{D}_z^{\mu-m} \{f(z)\} \right\} & (m-1 \leq \Re(\mu) < m \text{ } (m \in \mathbb{N})). \end{cases}$$

It is easy to find the following formula:

$$(43) \quad \mathcal{D}_z^\mu \{z^\lambda\} = \frac{\Gamma(\lambda+1)}{\Gamma(\lambda-\mu+1)} z^{\lambda-\mu} \quad (\Re(\lambda) > -1).$$

Theorem 6.1. *The following fractional derivative formula for $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, s, a)$ in (4) holds true:*

$$(44) \quad \begin{aligned} & \mathcal{D}_z^{\gamma-\xi} \mathcal{D}_t^{\nu-\zeta} \left\{ z^{\gamma-1} t^{\nu-1} \Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z^\tau, t^\kappa, s, a) \right\} \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\xi)\Gamma(\zeta)} z^{\xi-1} t^{\zeta-1} \Phi_{\alpha, \beta; \xi; \lambda, \mu; \zeta}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z^\tau, t^\kappa, s, a) \\ & \quad (\min\{\Re(\gamma), \Re(\nu)\} > 0; \min\{\tau, \kappa\} > 0). \end{aligned}$$

Proof. By virtue of the definition (4) of $\Phi_{\alpha, \beta; \gamma; \lambda, \mu; \nu}^{(\delta, \eta, \tau; \rho, \sigma, \kappa)}(z, s, a)$ and the formula (43), the assertion (44) follows easily. \square

Corollary 6.2. *Each of the following fractional derivative formulas holds true:*

$$(45) \quad \begin{aligned} & \mathcal{D}_z^{\gamma-\xi} \mathcal{D}_t^{\nu-\zeta} \left\{ z^{\gamma-1} t^{\nu-1} \Phi_{\beta;\gamma;\mu;\nu}^{(\eta,\tau;\sigma,\kappa)}(z^\tau, t^\kappa, s, a) \right\} \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\xi)\Gamma(\zeta)} z^{\xi-1} t^{\zeta-1} \Phi_{\beta;\xi;\mu;\zeta}^{(\eta,\tau;\sigma,\kappa)}(z^\tau, t^\kappa, s, a) \\ & \quad (\min\{\Re(\gamma), \Re(\nu)\} > 0; \min\{\tau, \kappa\} > 0); \end{aligned}$$

$$(46) \quad \begin{aligned} & \mathcal{D}_z^{\gamma-\xi} \mathcal{D}_t^{\nu-\zeta} \left\{ z^{\gamma-1} t^{\nu-1} \Phi_{\beta;\gamma;\mu;\nu}(z, t, s, a) \right\} \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\xi)\Gamma(\zeta)} z^{\xi-1} t^{\zeta-1} \Phi_{\beta;\xi;\mu;\zeta}(z, t, s, a) \\ & \quad (\min\{\Re(\gamma), \Re(\nu)\} > 0). \end{aligned}$$

Theorem 6.3. *The following fractional derivative formula holds true:*

$$(47) \quad \begin{aligned} & \mathcal{D}_z^{\alpha-\gamma} \mathcal{D}_t^{\lambda-\nu} \left\{ z^{\alpha-1} t^{\lambda-1} \Phi_{\beta;\mu}^*(z, t, s, a) \right\} \\ &= \frac{\Gamma(\alpha)\Gamma(\lambda)}{\Gamma(\gamma)\Gamma(\nu)} z^{\gamma-1} t^{\nu-1} \Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}(z, t, s, a) \\ & \quad (\Re(\nu) > \Re(\mu) > 0). \end{aligned}$$

Proof. Using the result (43) and (6), we can easily deduce fractional derivative formulas for (7) as in (47). \square

7. Multi-Parametric Extension of $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$

Here we present a multi-parametric extension of the $\Phi_{\alpha,\beta;\gamma;\lambda,\mu;\nu}^{(\delta,\eta,\tau;\rho,\sigma,\kappa)}(z, t, s, a)$ by introducing arbitrary number of numerator and denominator parameters as follows:

$$(48) \quad \begin{aligned} & \Phi_{\lambda_1, \dots, \lambda_p; \gamma_1, \dots, \gamma_q; \mu_1, \dots, \mu_r; \nu_1, \dots, \nu_w}^{(\rho_1, \dots, \rho_p, \tau_1, \dots, \tau_q; \sigma_1, \dots, \sigma_r, \kappa_1, \dots, \kappa_w)}(z, t, s, a) \\ &:= \sum_{m, n=0}^{\infty} \frac{\prod_{j=1}^p (\lambda_j)_{m\rho_j} \prod_{j=1}^r (\mu_j)_{n\sigma_j}}{\prod_{j=1}^q (\gamma_j)_{m\tau_j} \prod_{j=1}^w (\nu_j)_{n\kappa_j}} \frac{z^m t^n}{m!n!(m+n+a)^s} \end{aligned}$$

$$(\lambda_i, \mu_j \in \mathbb{C} \ (i = 1, \dots, p; \ j = 1, \dots, r);$$

$$a, \gamma_i, \nu_j \in \mathbb{C} \setminus \mathbb{Z}_0^- \ (i = 1, \dots, q; \ j = 1, \dots, w);$$

$$\rho_i, \tau_j, \sigma_k, \kappa_l \in \mathbb{R}^+ \ (i = 1, \dots, p; \ j = 1, \dots, q; \ k = 1, \dots, r; \ l = 1, \dots, w);$$

$$\Delta > -1, \delta > -1 \text{ when } s, z, t \in \mathbb{C};$$

$$\Delta = -1, \delta = -1 \text{ and } s \in \mathbb{C} \text{ when } |z| < \Delta^*, |t| < \delta^*;$$

$$\Delta = -1 \text{ and } \Re(\Xi) > 0 \text{ when } |z| = \Delta^*, |t| = \delta^* \text{ where}$$

$$\begin{aligned}\Delta^* &:= \left(\prod_{j=0}^p \rho_j^{-\rho_j} \right) \cdot \left(\prod_{j=0}^q \tau_j^{\tau_j} \right), \quad \Delta = \sum_{j=0}^q \tau_j - \sum_{j=0}^p \rho_j \quad \text{and} \\ \delta^* &:= \left(\prod_{j=0}^r \sigma_j^{-\sigma_j} \right) \cdot \left(\prod_{j=0}^w \kappa_j^{\kappa_j} \right), \quad \delta = \sum_{j=0}^w \kappa_j - \sum_{j=0}^r \sigma_j \quad \text{and} \\ \Xi &= s + \sum_{j=0}^q \gamma_j + \sum_{j=0}^w \nu_j - \sum_{j=0}^p \lambda_j - \sum_{j=0}^r \mu_j.\end{aligned}$$

Theorem 7.1. *The following integral representation in (48) holds true:*

$$\begin{aligned}(49) \quad & \Phi_{\lambda_1, \dots, \lambda_p; \gamma_1, \dots, \gamma_q; \mu_1, \dots, \mu_r; \nu_1, \dots, \nu_w}^{(\rho_1, \dots, \rho_p, \tau_1, \dots, \tau_q; \sigma_1, \dots, \sigma_r, \kappa_1, \dots, \kappa_w)}(z, t, s, a) \\ &:= \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-ax} {}_p\Psi_q^* \left[\begin{matrix} (\lambda_1, \rho_1), \dots, (\lambda_p, \rho_p); \\ (\gamma_1, \tau_1), \dots, (\gamma_p, \tau_p); \end{matrix} ; ze^{-x} \right] \\ & \quad \times {}_r\Psi_w^* \left[\begin{matrix} (\mu_1, \sigma_1), \dots, (\mu_r, \sigma_r); \\ (\nu_1, \kappa_1), \dots, (\nu_w, \kappa_w); \end{matrix} ; te^{-x} \right] dx \\ & (\rho_j, \tau_k, \sigma_l, \kappa_m \in \mathbb{R}^+ (j = 1, \dots, p; k = 1, \dots, q; l = 1, \dots, r; m = 1, \dots, w) \\ & \quad \text{and } \min\{\Re(s), \Re(a)\} > 0; |z| < \Delta^*, |t| < \delta^*).\end{aligned}$$

Theorem 7.2. *The following Mellin-Barnes contour integral representation in (48) holds true:*

$$\begin{aligned}(50) \quad & \Phi_{\lambda_1, \dots, \lambda_p; \gamma_1, \dots, \gamma_q; \mu_1, \dots, \mu_r; \nu_1, \dots, \nu_w}^{(\rho_1, \dots, \rho_p, \tau_1, \dots, \tau_q; \sigma_1, \dots, \sigma_r, \kappa_1, \dots, \kappa_w)}(z, t, s, a) \\ &= \frac{\prod_{j=1}^q \Gamma(\gamma_j) \prod_{j=1}^w \Gamma(\nu_j)}{\prod_{j=1}^p \Gamma(\lambda_j) \prod_{j=1}^r \Gamma(\mu_j)} \frac{1}{(2\pi i)^2} \int_{-i\infty}^{+i\infty} \int_{-i\infty}^{+i\infty} \Gamma(-\xi) \Gamma(-\zeta) \\ & \quad \times \frac{\prod_{j=1}^p \Gamma(\lambda_j + \rho_j \xi) \prod_{j=1}^r \Gamma(\mu_j + \sigma_j \zeta) (-z)^\xi (-t)^\zeta}{(\xi + \zeta + a)^s \prod_{j=1}^q \Gamma(\gamma_j + \tau_j \xi) \prod_{j=1}^w \Gamma(\nu_j + \kappa_j \zeta)} d\xi d\zeta \\ & (\max\{|\arg(-z)|, |\arg(-t)|\} < \pi).\end{aligned}$$

Theorem 7.3. *The following fractional derivative formula in (48) holds true:*

$$\begin{aligned}(51) \quad & \mathcal{D}_z^{\gamma-\xi} \mathcal{D}_t^{\nu-\zeta} \left\{ z^{\gamma-1} t^{\nu-1} \Phi_{\lambda_1, \dots, \lambda_p; \gamma_1, \dots, \gamma_q; \mu_1, \dots, \mu_r; \nu_1, \dots, \nu_w}^{(\rho_1, \dots, \rho_p, \tau_1, \dots, \tau_q; \sigma_1, \dots, \sigma_r, \kappa_1, \dots, \kappa_w)}(z^\tau, t^\kappa, s, a) \right\} \\ &= \frac{\Gamma(\gamma)\Gamma(\nu)}{\Gamma(\xi)\Gamma(\zeta)} z^{\xi-1} t^{\zeta-1} \Phi_{\lambda_1, \dots, \lambda_p; \xi; \gamma_1, \dots, \gamma_q; \gamma; \mu_1, \dots, \mu_r; \zeta; \nu_1, \dots, \nu_w, \nu}^{(\rho_1, \dots, \rho_p, \tau, \tau_1, \dots, \tau_q, \tau; \sigma_1, \dots, \sigma_r, \kappa, \kappa_1, \dots, \kappa_w, \kappa)}(z^\tau, t^\kappa, s, a) \\ & (\min\{\Re(\gamma), \Re(\nu)\} > 0; \min\{\tau, \kappa\} > 0).\end{aligned}$$

Acknowledgments. The authors would like to express their deep-felt thanks for the reviewer's detailed and helpful comments.

References

- [1] E. W. Barnes, *The asymptotic expansion of integral functions defined by Taylor series*, Philos. Trans. Roy. Soc. London Ser. A **206** (1906), 249–297.
- [2] M. A. Chaudhry and S. M. Zubair, *On a Class of Incomplete Gamma Functions with Applications*, Chapman and Hall, (CRC Press Company), Boca Raton, London, New York and Washington, D. C., 2001.
- [3] J. Choi, D. S. Jang, and H. M. Srivastava, *A generalization of the Hurwitz-Lerch Zeta function*, Integral Transforms Spec. Funct. **19** (2008), no. 1-2, 65–79.
- [4] O. Daman and M. A. Pathan, *A further generalization of the Hurwitz Zeta function*, Math. Sci. Res. J. **16** (2012), no. 10, 251–259.
- [5] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions. Vol. I*, McGraw-Hill Book Company, New York, Toronto and London, 1953.
- [6] M. Garg, K. Jain, and S. L. Kalla, *A further study of general Hurwitz-Lerch zeta function*, Algebras Groups Geom. **25** (2008), 311–319.
- [7] S. P. Goyal and R. K. Laddha, *On the generalized Zeta function and the generalized Lambert function*, Ganita Sandesh **11** (1997), no. 2, 99–108.
- [8] N. T. Hai and S. B. Yakubovich, *The Double Mellin-Barnes Type Integrals and Their Applications to Convolution Theory*, World Scientific, Singapore, 1992.
- [9] D. Jankov, T. K. Pogany, and R. K. Saxena, *An extended general Hurwitz-Lerch Zeta function as a Mathieu (a, λ) -series*, Appl. Math. Lett. **24** (2011), no. 8, 1473–1476.
- [10] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematical Studies, Vol. **204**, Elsevier (North-Holland) Science Publishers, Amsterdam, London and New York, 2006.
- [11] S. D. Lin and H. M. Srivastava, *Some families of the Hurwitz-Lerch Zeta functions and associated fractional derivative and other integral representations*, Appl. Math. Comput. **154** (2004), no. 3, 725–733.
- [12] A. M. Mathai and R. K. Saxena, *Generalized Hypergeometric Functions with Applications in Statistics and Physical Sciences*, Lecture Notes Series No. 348, Springer-Verlag, Berlin, New York, Heidelberg, Germany, 1973.
- [13] ———, *The H-function with Applications in Statistics and Other Disciplines*, Wiley Eastern Ltd. New Delhi and John Wiley and Sons, Inc. New York, 1978.
- [14] A. M. Mathai, R. K. Saxena, and H. J. Haubold, *The H-Functions: Theory and Applications*, Springer, New York, 2010.
- [15] R. B. Paris and D. Kaminski, *Asymptotic and Mellin-Barnes Integrals*, Cambridge University Press, Cambridge, 2001.
- [16] M. A. Pathan and O. Daman, *On generalization of Hurwitz zeta function*, Submitted.
- [17] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Translated from the Russian: Integrals and Derivatives of Fractional Order and Some of Their Applications (“Nauka i Tekhnika”, Minsk, 1987); Gordon and Breach Science Publishers: Reading, UK, 1993.
- [18] H. M. Srivastava, *A new family of the λ -generalized Hurwitz-Lerch Zeta functions with applications*, Appl. Math. Inf. Sci. **8** (2014), no. 4, 1485–1500.
- [19] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer, Academic Publishers, Dordrecht, Boston and London, 2001.
- [20] ———, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier Science, Publishers, Amsterdam, London and New York, 2012.
- [21] H. M. Srivastava, D. Jankov, T. K. Pogany, and R. K. Saxena, *Two-sided inequalities for the extended Hurwitz-Lerch Zeta function*, Comput. Math. Appl. **62** (2011), no. 1, 516–522.
- [22] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1985.

- [23] H. M. Srivastava, M.-J. Luo and R. K. Raina, *New results involving a class of generalized Hurwitz-Lerch Zeta functions and their applications*, Turkish J. Anal. Number Theory **1** (2013), no. 1, 26–35.
- [24] H. M. Srivastava and H. L. Manocha, *A Treatise on Generating Functions*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, 1984.
- [25] H. M. Srivastava, R. K. Saxena, T. K. Pogány, and R. Saxena, *Integral and computational representations of the extended Hurwitz-Lerch Zeta function*, Integral Transforms Spec. Funct. **22** (2011), no. 7, 487–506.

JUNESANG CHOI
DEPARTMENT OF MATHEMATICS
DONGGUK UNIVERSITY
GYEONGJU 38066, KOREA
E-mail address: junesang@mail.dongguk.ac.kr

RAKESH K. PARMAR
DEPARTMENT OF MATHEMATICS
GOVERNMENT COLLEGE OF ENGINEERING AND TECHNOLOGY
BIKANER-334004, RAJASTHAN STATE, INDIA
E-mail address: rakeshparmar27@gmail.com

RAM K. SAXENA
DEPARTMENT OF MATHEMATICS AND STATISTICS
JAI NARAIN VYAS UNIVERSITY
JODHPUR-342004, RAJASTHAN STATE, INDIA
E-mail address: ram.saxena@yahoo.com