

## SECOND MAIN THEOREM AND UNIQUENESS PROBLEM OF ZERO-ORDER MEROMORPHIC MAPPINGS FOR HYPERPLANES IN SUBGENERAL POSITION

THI TUYET LUONG, DANG TUYEN NGUYEN, AND DUC THOAN PHAM

**ABSTRACT.** In this paper, we show the Second Main Theorems for zero-order meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  intersecting hyperplanes in subgeneral position without truncated multiplicity by considering the  $p$ -Casorati determinant with  $p \in \mathbb{C}^m$  instead of its Wronskian determinant. As an application, we give some unicity theorems for meromorphic mapping under the growth condition “order=0”. The results obtained include  $p$ -shift analogues of the Second Main Theorem of Nevanlinna theory and Picard’s theorem.

### 1. Introduction

In 2006, R. Halburd-R. Korhonen [7] considered the Second Main Theorem for complex difference operator with finite order in complex plane. Later, in [8] and [18], difference analogues of the Second Main Theorem for holomorphic curves in  $\mathbb{P}^n(\mathbb{C})$  were obtained independently, and in [2] and [12], difference analogues of the Second Main Theorem for meromorphic functions on  $\mathbb{C}^m$  were obtained. In particular, Nevanlinna theory for the  $p$ -difference operator can be found in [1, 11, 15–17, 19].

Recently, T. B. Cao-R. Korhonen [3] obtained a new natural difference analogue of H. Cartan’s Theorem for meromorphic mapping  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$ . In which, the counting function  $N(r, \nu_{W(f)}^0)$  of the Wronskian determinant of  $f$  is replaced by the counting function  $N(r, \nu_{C^c(f)}^0)$  of the Casorati determinant of  $f$  (it was called the finite difference Wronskian determinant in [18]) and in addition, the hyper-order of  $f$  is strictly less than one.

Our first aim in this paper is to prove a new natural  $p$ -difference analogue Second Main Theorem for zero-order meromorphic mapping by considering  $p$ -Casorati determinant. For our purpose, we now recall some notations.

---

Received November 20, 2016; Revised June 18, 2017; Accepted August 11, 2017.

2010 *Mathematics Subject Classification.* Primary 53A10; Secondary 53C42, 30D35, 32H30.

*Key words and phrases.* second main theorem, Nevanlinna theory, Casorati determinant, zero-order meromorphic mapping, hyperplanes.

Let  $p \in \mathbb{C}^m$ , denote by  $\mathcal{M}_m$  the set of all meromorphic functions on  $\mathbb{C}^m$ , denote by  $\phi_p$  the set of all meromorphic functions of  $\mathcal{M}_m$  satisfying  $f(z) = f(pz)$  and denote by  $\phi_p^0$  the set of all meromorphic functions of  $\phi_p$  having their zero-orders. Obviously,  $\phi_p^0 \subset \phi_p \subset \mathcal{M}_m$ .

**Definition 1.** Let  $f$  be a meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  with reduced representation  $f = (f_0 : \cdots : f_n)$ . Then the map  $f$  is said to be linearly nondegenerate over field  $\phi_p^0$  if the entire functions  $f_0, \dots, f_n$  are linearly independent over field  $\phi_p^0$ .

For  $c = (c_1, \dots, c_m)$  and  $p = (p_1, \dots, p_m)$  with  $p_i \neq 0$  ( $1 \leq i \leq m$ ) and  $z = (z_1, \dots, z_m)$ , we write  $c+z = (c_1+z_1, \dots, c_m+z_m)$  and  $pz = (p_1z_1, \dots, p_mz_m)$ . Denote

$$f(z) \equiv f := \bar{f}^{[0]}, f(z+c) \equiv \bar{f} := \bar{f}^{[1]}, f(z+2c) \equiv \bar{f} := \bar{f}^{[2]}, \dots, f(z+kc) \equiv \bar{f}^{[k]}$$

and

$$f(z) \equiv f := \hat{f}^{[0]}, f(pz) \equiv \hat{f} := \hat{f}^{[1]}, f(p^2z) \equiv \hat{f} := \hat{f}^{[2]}, \dots, f(p^kz) \equiv \hat{f}^{[k]}.$$

Let

$$D^{(j)} = \left( \frac{\partial}{\partial z_1} \right)^{\alpha_1(j)} \cdots \left( \frac{\partial}{\partial z_m} \right)^{\alpha_m(j)}$$

be a partial differentiation operator of order at most  $j = \sum_{k=1}^m \alpha_k(j)$ . Similarly as the Wronskian determinant

$$W(f) = W(f_0, \dots, f_n) = \begin{vmatrix} f_0 & f_1 & \cdots & f_n \\ D^{(1)}f_0 & D^{(1)}f_1 & \cdots & D^{(1)}f_n \\ \vdots & \vdots & \vdots & \vdots \\ D^{(n)}f_0 & D^{(n)}f_1 & \cdots & D^{(n)}f_n \end{vmatrix},$$

the Casorati determinant is defined by

$$C^c(f) = C^c(f_0, \dots, f_n) = \begin{vmatrix} f_0 & f_1 & \cdots & f_n \\ \bar{f}_0 & \bar{f}_1 & \cdots & \bar{f}_n \\ \vdots & \vdots & \vdots & \vdots \\ \bar{f}_0^{[n]} & \bar{f}_1^{[n]} & \cdots & \bar{f}_n^{[n]} \end{vmatrix}$$

and the  $p$ -Casorati determinant is defined by

$$C_p(f) = C_p(f_0, \dots, f_n) = \begin{vmatrix} f_0 & f_1 & \cdots & f_n \\ \hat{f}_0 & \hat{f}_1 & \cdots & \hat{f}_n \\ \vdots & \vdots & \vdots & \vdots \\ \hat{f}_0^{[n]} & \hat{f}_1^{[n]} & \cdots & \hat{f}_n^{[n]} \end{vmatrix}.$$

**Definition 2.** Let  $\{H_j\}_{j=1}^q$  be the hyperplanes in  $\mathbb{P}^n(\mathbb{C})$ . Let  $N \geq n$  and  $q \geq N+1$ . The family  $\{H_j\}_{j=1}^q$  is said to be in  $N$ -subgeneral position in  $\mathbb{P}^n(\mathbb{C})$

if for every subset  $R \subset \{1, \dots, q\}$  with the cardinality  $|R| = N + 1$ , then

$$\bigcap_{j \in R} H_j = \emptyset.$$

If they are in  $n$ -subgeneral position, we simply say that they are in general position.

Consider  $f$  be a meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  with reduced representation  $f = (f_0 : \dots : f_n)$  and a hyperplane  $H : a_0\omega_0 + \dots + a_n\omega_n = 0$ . We define

$$(f, H) = H(f) := a_0f_0 + \dots + a_nf_n,$$

which is a holomorphic function on  $\mathbb{C}^m$ .

Using above notations, we have the  $p$ -difference analogue of H. Cartan's Theorem [4] as follows.

**Theorem 1.** *Let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j \in \{1, \dots, m\}$  and let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be a linearly nondegenerate meromorphic mapping over the field  $\phi_p^0$ . Let  $H_j$  ( $1 \leq j \leq q$ ) be  $q$  hyperplanes in  $\mathbb{P}^n(\mathbb{C})$ , located in  $N$ -subgeneral position. Assume that  $f$  has the zero-order. Then we have*

$$(q - 2N + n - 1)T(r, f) \leq \sum_{j=1}^q N(r, \nu_{H_j(f)}^0) - \frac{N}{n}N(r, \nu_{C_p(f)}^0) + o(T(r, f))$$

for all  $r$  on a set of logarithmic density 1.

Here, by  $\nu_\varphi^0$  we denote the zero-divisor of holomorphic function  $\varphi$  from  $\mathbb{C}^m$  into  $\mathbb{C}$ .

**Definition 3.** Let  $k \in \mathbb{N}$ ,  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j \in \{1, \dots, m\}$  and  $a \in \mathbb{C}$ . An  $a$ -point  $z_0$  of meromorphic function  $h(z)$  is said to be  $k$ -successive with separated  $p$  respect to the rescaling  $\tau_p(z) = pz$ , if the  $k$  functions  $h(p^l z)$ , ( $l = 1, \dots, k$ ) take the value  $a$  at  $z = z_0$  with multiplicity not less than that of  $h(z)$  there. All the other  $a$ -points of  $h(z)$  are called  $k$ -aperiodic of pace  $p$  respect to the rescaling  $\tau_p(z) = pz$ .

Consider  $H$  be a hyperplane. By  $\hat{N}^{[k,p]}(r, H(f))$ , we denote the counting function of  $k$ -aperiodic zeros of the function  $H(f)$  of pace  $p$  respect to the rescaling  $\tau_p(z) = pz$ .

Note that  $\hat{N}^{[k,p]}(r, H(f)) \equiv 0$  when all zeros of  $H(f)$  with taking their multiplicities into account are located periodically with period  $p$  respect to the rescaling  $\tau_p(z) = pz$ . This is also the case when the hyperplane  $H$  is forward invariant by  $f$  with respect to the rescaling  $\tau_p(z) = pz$ , i.e.,  $\tau_p(f^{-1}(H)) \subset f^{-1}(H)$  and  $f^{-1}(H)$  is considered to be multi-sets in which each point is repeated according to its multiplicity. Then we have the result as follows.

**Theorem 2.** *Let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j \in \{1, \dots, m\}$  and let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be a linearly nondegenerate meromorphic mapping*

over the field  $\phi_p^0$ . Let  $H_j$  ( $1 \leq j \leq q$ ) be  $q$  hyperplanes in  $\mathbb{P}^n(\mathbb{C})$ , located in  $N$ -subgeneral position. Assume that  $f$  has the zero-order. Then we have

$$(q - 2N + n - 1)T(r, f) \leq \sum_{j=1}^q \hat{N}^{[n,p]}(r, H_j(f)) + o(T(r, f))$$

for all  $r$  on a set of logarithmic density 1.

The uniqueness problem for meromorphic mappings was first investigated by R. Nevanlinna. In 1975, H. Fujimoto [5] generalized Nevanlinna's five-value theorem to the case of higher dimension by showing that if two linearly non-degenerate meromorphic mappings  $f, g : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  have the same inverse images counted with multiplicities for  $q \geq 3n+2$  hyperplanes in general position in  $\mathbb{P}^n(\mathbb{C})$ , then  $f \equiv g$ .

By considering the uniqueness problem for holomorphic curves  $f(z)$  and  $f(z+c)$  also for holomorphic curves  $f(z)$  and  $f(pz)$  intersecting hyperplanes in general position, R. Halburd, R. Korhonen, K. Tohge [8, Theorem 1.1 and Theorem 6.1] obtained a difference analogue of Picard's theorem. Recently, T. B. Cao, R. Korhonen [3] generalized this result [8, Theorem 1.1] for the case of meromorphic mappings  $f(z)$  and  $f(z+c)$  intersecting hyperplanes in subgeneral position.

Our final aim in this paper is to extend the result in [8, Theorem 6.1] to meromorphic mappings  $f(z)$  and  $f(pz)$  of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  intersecting hyperplanes in  $N$ -subgeneral position. Our result is a difference analogue of Picard's theorem. Namely, we will prove the following theorem.

**Theorem 3.** *Let  $f$  be a zero-order meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0, 1$  for all  $j \in \{1, \dots, m\}$ . Assume that  $f$  is forward invariant over  $q$  hyperplanes in  $N$ -subgeneral position in  $\mathbb{P}^n(\mathbb{C})$  respect to the rescaling  $\tau_p(z) = pz$ . Then the image of  $f$  is contained in a projective linear subspace over  $\phi_p^0$  of dimension  $\leq \left\lfloor \frac{N}{q-N} \right\rfloor$ . Special, if  $q \geq 2N + 1$  then  $f(z) = f(pz)$ .*

Note that when  $|p_i| \neq 1$  for all  $i \in \{1, \dots, m\}$ , then  $f(z) = f(pz)$  implies that  $f$  must be a constant mapping. Immediately, we have the following corollary.

**Corollary 4.** *Let  $f$  be a zero-order meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  satisfying  $|p_j| \neq 0, 1$  for all  $j \in \{1, \dots, m\}$ . Assume that  $f$  is forward invariant over  $q$  hyperplanes in general position in  $\mathbb{P}^n(\mathbb{C})$  respect to the rescaling  $\tau_p(z) = pz$ . If  $q \geq 2n + 1$  then  $f$  is constant.*

## 2. Preliminaries and auxiliary lemmas

**2.1.** We set  $\|z\| = (|z_1|^2 + \dots + |z_n|^2)^{1/2}$  for  $z = (z_1, \dots, z_n) \in \mathbb{C}^n$  and define

$$B_m(r) := \{z \in \mathbb{C}^m : \|z\| < r\}, \quad S_m(r) := \{z \in \mathbb{C}^m : \|z\| = r\} \quad (0 < r < \infty).$$

Define

$$\sigma_m(z) := (dd^c \|z\|^2)^{m-1} \quad \text{and}$$

$$\eta_m(z) := d^c \log \|z\|^2 \wedge (dd^c \log \|z\|^2)^{m-1} \text{ on } \mathbb{C}^m \setminus \{0\}.$$

**2.2.** Let  $F$  be a nonzero holomorphic function on a domain  $\Omega$  in  $\mathbb{C}^m$ . For a set  $\alpha = (\alpha_1, \dots, \alpha_m)$  of nonnegative integers, we set  $|\alpha| = \alpha_1 + \dots + \alpha_m$  and  $\mathcal{D}^\alpha F = \frac{\partial^{|\alpha|} F}{\partial^{\alpha_1} z_1 \dots \partial^{\alpha_m} z_m}$ . We define the map  $\nu_F : \Omega \rightarrow \mathbb{Z}$  by

$$\nu_F(z) := \max \{n : \mathcal{D}^\alpha F(z) = 0 \text{ for all } \alpha \text{ with } |\alpha| < n\} \quad (z \in \Omega).$$

We mean by a divisor on a domain  $\Omega$  in  $\mathbb{C}^m$  a map  $\nu : \Omega \rightarrow \mathbb{Z}$  such that, for each  $a \in \Omega$ , there are nonzero holomorphic functions  $F$  and  $G$  on a connected neighbourhood  $U \subset \Omega$  of  $a$  such that  $\nu(z) = \nu_F(z) - \nu_G(z)$  for each  $z \in U$  outside an analytic set of dimension  $\leq m-2$ . Two divisors are regarded as the same if they are identical outside an analytic set of dimension  $\leq m-2$ . For a divisor  $\nu$  on  $\Omega$  we set  $|\nu| := \overline{\{z : \nu(z) \neq 0\}}$ , which is a purely  $(m-1)$ -dimensional analytic subset of  $\Omega$  or empty.

Take a nonzero meromorphic function  $\varphi$  on a domain  $\Omega$  in  $\mathbb{C}^n$ . For each  $a \in \Omega$ , we choose nonzero holomorphic functions  $F$  and  $G$  on a neighbourhood  $U \subset \Omega$  such that  $\varphi = \frac{F}{G}$  on  $U$  and  $\dim(F^{-1}(0) \cap G^{-1}(0)) \leq m-2$ , and we define the divisors  $\nu_\varphi^0, \nu_\varphi^\infty$  by  $\nu_\varphi^0 := \nu_F, \nu_\varphi^\infty := \nu_G$ , which are independent of choices of  $F$  and  $G$  and so globally well-defined on  $\Omega$ .

**2.3.** For a divisor  $\nu$  on  $\mathbb{C}^m$ , we define the counting functions of  $\nu$  by

$$n(t) = \begin{cases} \int_{|\nu| \cap B(t)} \nu(z) \sigma_{m-1} & \text{if } m \geq 2 \\ \sum_{|z| \leq t} \nu(z) & \text{if } m = 1 \end{cases}$$

and

$$N(r, \nu) = \int_1^r \frac{n(t)}{t^{2m-1}} dt \quad (1 < r < \infty).$$

Let  $\varphi : \mathbb{C}^m \rightarrow \mathbb{C}$  be a meromorphic function. Define

$$N_\varphi(r) = N(r, \nu_\varphi).$$

**2.4.** Let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be a meromorphic mapping. For arbitrarily fixed homogeneous coordinates  $(w_0 : \dots : w_n)$  on  $\mathbb{P}^n(\mathbb{C})$ , we take a reduced representation  $f = (f_0 : \dots : f_n)$ , which means that each  $f_i$  is a holomorphic function on  $\mathbb{C}^m$  and  $f(z) = (f_0(z) : \dots : f_n(z))$  outside the analytic set  $I(f) = \{z \in \mathbb{C}^m : f_0(z) = \dots = f_n(z) = 0\}$  of codimension  $\geq 2$ . Set  $\|f\| = (\sum_{j=0}^n |f_j|^2)^{1/2}$ . The characteristic function of  $f$  is defined by

$$T(r, f) = \int_{r_0}^r \frac{dt}{2^{m-1}} \int_{B_m(r)} dd^c \log \|f\|^2 \wedge \sigma_m(z)$$

$$= \int_{S_m(r)} \log \|f\| \eta_m - \int_{S_m(r_0)} \log \|f\| \eta_m(z).$$

Note that  $T(r, f)$  is independent of the choice of the representation of  $f$ . The order and hyperorder of  $f$  are respectively defined by

$$\sigma(f) := \limsup_{r \rightarrow \infty} \frac{\log^+ T(r, f)}{\log r} \quad \text{and} \quad \zeta(f) := \limsup_{r \rightarrow \infty} \frac{\log^+ \log^+ T(r, f)}{\log r},$$

where  $\log^+ x := \max\{\log x, 0\}$  for any  $x > 0$ .

**2.5.** Let  $f$  be a meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^n(\mathbb{C})$  with reduced representation  $f = (f_0 : \cdots : f_n)$  and a hyperplane  $H : a_0\omega_0 + \cdots + a_n\omega_n = 0$  satisfies

$$(f, H) = a_0f_0 + \cdots + a_nf_n \neq 0.$$

The proximity function is defined as

$$m_{f,H}(r) := \int_{S_m(r)} \log^+ \frac{\|f\| \cdot \|H\|}{|(f, H)|} \eta_m(z) + \int_{S_m(1)} \log^+ \frac{\|f\| \cdot \|H\|}{|(f, H)|} \eta_m(z).$$

We have the First Main Theorem of Nevanlinna theory

$$m_{f,H}(r) + N(r, \nu_{H(f)}^0) = T(r, f) + O(1),$$

where  $O(1)$  is a constant independent of  $r$ .

**2.6.** Let  $\varphi$  be a nonzero meromorphic function on  $\mathbb{C}^m$ , which is occasionally regarded as a meromorphic map into  $\mathbb{P}^1(\mathbb{C})$ . The proximity function of  $\varphi$  is defined by

$$m(r, \varphi) := \int_{S_m(r)} \log^+ |\varphi| \eta_m.$$

**Lemma 5** ([1, Lemmas 5.1, 5.2, and 5.3]). *Let  $f$  be a non-constant zero-order meromorphic function of  $\mathbb{C}$  into  $\mathbb{C}$  and let  $p \in \mathbb{C} \setminus \{0\}$ . Then*

$$m\left(r, \frac{f(pz)}{f(z)}\right) < \frac{4D_1 + 2D_2}{2^n} T(r, f(z))$$

*on a set of logarithmic density 1 for all  $n \in \mathbb{N}$ , where  $D_1, D_2$  are positive constants.*

**Lemma 6** ([9, Lemma 4]). *If  $T : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is an increasing function such that order*

$$\sigma(T) = \lim_{r \rightarrow \infty} \frac{\log T(r)}{\log r} = 0,$$

*then the set*

$$E := \{r \in \mathbb{R}^+ : T(C_1r) \geq C_2T(r)\}$$

*has logarithmic density 0 for all  $C_1 > 1$  and  $C_2 > 1$ .*

**Lemma 7** ([1, Lemma 5.4]). *Let  $T : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be an increasing function and  $U : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ . If there exists a decreasing sequence  $\{c_n\}_{n \in \mathbb{N}}$  such that  $c_n \rightarrow 0$  as  $n \rightarrow \infty$  and for all  $n \in \mathbb{N}$ , the set*

$$F_n = \{r \geq 1 : U(r) < c_n T(r)\}$$

*has logarithmic density 1, then  $U(r) = o(T(r))$  on a set of logarithmic density 1.*

**Lemma 8.** *Let  $T$  be a function as in Lemma 6 and let  $p \in \mathbb{R}^+$ . Then we have*

$$T(pr) = T(r) + o(T(r))$$

*on a set of logarithmic density 1.*

*Proof. Case 1:  $p \leq 1$ .* Since  $T(r)$  is an increasing function, we have  $T(pr) \leq T(r)$  for all  $r > 0$ . Obviously, the conclusion holds.

*Case 2:  $p > 1$ .* By Lemma 6, for each  $n \in \mathbb{N}$ , we have

$$E_n := \left\{ r \geq 1 : T(pr) < \left(1 + \frac{1}{n}\right) T(r) \right\}$$

has logarithmic density 1. Put  $U(r) = T(pr) - T(r)$ , we deduce that

$$0 < U(r) < \frac{1}{n} T(r)$$

on a set of logarithmic density 1. It follows from Lemma 7 that  $U(r) = o(T(r))$  on a set of logarithmic density 1. Therefore, we get

$$(2.1) \quad T(pr) = T(r) + o(T(r))$$

on a set of logarithmic density 1. Therefore, the proof of the Lemma 8 is finished.  $\square$

For each  $\omega \in \overline{B}_{m-1}(r)$ , we define a function  $p_r(\omega) = \sqrt{r^2 - |\omega|^2}$ . We need the following lemma from W. Stoll.

**Lemma 9** ([10]). *Let  $r > 0$  and let  $h$  be a function on  $S_m(r)$  such that  $h\eta_m$  is integrable over  $S_m(r)$ . Then*

$$\int_{S_m(r)} h(z)\eta_m(z) = \frac{1}{r^{2m-2}} \int_{\overline{B}_{m-1}(r)} \sigma_{m-1}(\omega) \int_{S_1(P_r(\omega))} h(\omega, \zeta)\eta_1(\zeta).$$

Consider a non-constant meromorphic function  $f$  on  $\mathbb{C}^m$ , take  $\omega \in \mathbb{C}^{m-1}$  and define  $f_\omega(z) := f(\omega, z)$  on  $\mathbb{C}$ . We will prove the following lemma.

**Lemma 10.** *Let  $f$  be a meromorphic function on  $\mathbb{C}^m$  of zero-order such that  $f(0) \neq 0, \infty$  and let  $\tilde{p}_j := (1, \dots, p_j, \dots, 1)$  with  $p_j \neq 0$ . Then*

$$m \left( r, \frac{f(\tilde{p}_j z)}{f(z)} \right) = \int_{S_m(r)} \log^+ \left| \frac{f(\tilde{p}_j z)}{f(z)} \right| \eta_m(z) = o(T(r, f(z)))$$

*on a set of logarithmic density 1.*

*Proof.* By applying Lemma 9 for  $h(z) = \log^+ \left| \frac{f(\tilde{p}_j z)}{f(z)} \right|$ , we have

$$\begin{aligned} m\left(r, \frac{f(\tilde{p}_j z)}{f(z)}\right) &= \int_{S_m(r)} \log^+ \left| \frac{f(\tilde{p}_j z)}{f(z)} \right| \eta_m(z) \\ &= \frac{1}{r^{2m-2}} \int_{\bar{B}_{m-1}(r)} \sigma_{m-1}(\omega) \int_{S_1(P_r(\omega))} \log^+ \left| \frac{f_\omega(p_j z_j)}{f_\omega(z_j)} \right| \eta_1(\zeta) \\ &= \frac{1}{r^{2m-2}} \int_{\bar{B}_{m-1}(r)} m\left(P_r(\omega), \frac{f_\omega(p_j z_j)}{f_\omega(z_j)}\right) \sigma_{m-1}(\omega). \end{aligned}$$

By Lemma 5, there exist two positive constants  $D_1$  and  $D_2$  which are independent of  $P_r(\omega)$  such that for all  $n \in \mathbb{N}$ , we have

$$\begin{aligned} &m\left(r, \frac{f(\tilde{p}_j z)}{f(z)}\right) \\ &< \frac{1}{r^{2m-2}} \int_{\bar{B}_{m-1}(r)} \frac{4D_1 + 2D_2}{2^n} T(P_r(\omega), f_\omega(z_j)) \sigma_{m-1}(\omega) \\ &= \frac{4D_1 + 2D_2}{2^n} \cdot \frac{1}{r^{2m-2}} \int_{\bar{B}_{m-1}(r)} \sigma_{m-1}(\omega) \int_{S_1(P_r(\omega))} \log \|f_\omega(z_j)\| \eta_1(z_j) + O(1) \\ &= \frac{4D_1 + 2D_2}{2^n} \int_{S_m(r)} \log \|f(\omega, z_j)\| \eta_m(z) + O(1) \\ &= \frac{4D_1 + 2D_2}{2^n} T(r, f(z)) + O(1) \end{aligned}$$

on a set of logarithmic density 1 for all  $n \in \mathbb{N}$ . By applying the Lemma 7, we get

$$m\left(r, \frac{f(\tilde{p}_j z)}{f(z)}\right) = o(T(r, f(z)))$$

on a set of logarithmic density 1. We finish the proof of Lemma 10.  $\square$

The Lemma on the Logarithmic Derivative [4–6, 14] plays an important role in Nevanlinna theory. Here, it is replaced by the following lemma.

**Lemma 11.** *Let  $f$  be a non-constant zero-order meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{C}$  and  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j$ . Then*

$$m\left(r, \frac{f(pz)}{f(z)}\right) = o(T(r, f(z)))$$

*on a set of logarithmic density 1.*



*Proof.* Since  $f$  is a meromorphic function on  $\mathbb{C}^m$  of zero-order, according to Lemma 10, it follows that

$$\begin{aligned} m\left(r, \frac{f(pz)}{f(z)}\right) &= \int_{S_m(r)} \log^+ \left| \frac{f(pz)}{f(z)} \right| \eta_m(z) \\ &= \int_{S_m(r)} \log^+ \prod_{k=1}^n \left| \frac{f\left(\prod_{j=0}^k \tilde{p}_j z\right)}{f\left(\prod_{j=0}^{k-1} \tilde{p}_j z\right)} \right| \eta_m(z) \\ &\leq \sum_{k=1}^n \int_{S_m(r)} \log^+ \left| \frac{f\left(\prod_{j=0}^k \tilde{p}_j z\right)}{f\left(\prod_{j=0}^{k-1} \tilde{p}_j z\right)} \right| \eta_m(z) = o(T(r, f)) \end{aligned}$$

on a set of logarithmic density 1. The proof of Lemma 11 is finished.  $\square$

**Lemma 12.** *Let  $f$  be a meromorphic function on  $\mathbb{C}^m$  of zero-order such that  $f(0) \neq 0, \infty$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j$ . Then we have*

$$T(r, f(pz)) = T(r, f(z)) + o(T(r, f(z)))$$

on a set of logarithmic density 1.

*Proof.* By the First Main Theorem, we have

$$T\left(r, \frac{f(pz)}{f(z)}\right) = m\left(r, \frac{f(pz)}{f(z)}\right) + N\left(r, \frac{f(pz)}{f(z)}\right) + O(1).$$

Therefore, by Lemma 11, we get

$$(2.2) \quad T(r, f(pz)) - T(r, f(z)) = N(r, f(pz)) - N(r, f(z)) + o(T(r, f(z)))$$

on a set of logarithmic density 1. Also by the First Main Theorem, we deduce that

$$\overline{\lim}_{r \rightarrow \infty} \frac{\log(N(r, f))}{\log r} \leq \overline{\lim}_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r} = \sigma(f) = 0.$$

This, by Lemma 8, we have

$$(2.3) \quad N(|p|r, f) = N(r, f) + o(N(r, f)) \leq N(r, f) + o(T(r, f))$$

on a set of logarithmic density 1. Together (2.2) with (2.3), we get

$$T(r, f(pz)) \leq T(r, f(z)) + o(T(r, f(z)))$$

on a set of logarithmic density 1. We have the assertion of Lemma 12.  $\square$

The similar results to Lemmas 10, 11, and 12 can be found in [1, 11, 16, 17, 19].

It is known that holomorphic functions  $f_0, \dots, f_n$  on  $\mathbb{C}^m$  are linearly dependent over  $\mathbb{C}$  if and only if their Wronskian determinants  $W(f_0, \dots, f_n)$  vanish identically [6, 13, 14]. Similarly, holomorphic functions  $f_0, \dots, f_n$  on  $\mathbb{C}^m$  are linearly dependent over  $\mathcal{P}_c^\lambda$  if and only if their Casorati determinants

$C^c(f_0, \dots, f_n)$  vanish identically [3], where  $\mathcal{P}_c^\lambda$  is the field of  $c$ -periodic meromorphic functions having hyper-order of  $\lambda$ .

Here, we introduce a similar result for the case of  $p$ -Casorati determinant by the same method as in [8]. Namely, we have the following.

**Lemma 13.** *Let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be a meromorphic mapping with reduce presentation  $f = (f_0 : \dots : f_n)$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0$  for all  $j$ . Assume that  $\sigma(f) = 0$ . Then  $p$ -Casorati determinant  $C_p(f_0, \dots, f_n) \equiv 0$  if and only if the functions  $f_0, \dots, f_n$  are linear dependent over the field  $\phi_p^0$ .*

*Proof.* Suppose first that  $f_0, \dots, f_n$  are linear dependent over the field  $\phi_p^0$ . Then there exist  $\varphi_0, \dots, \varphi_n \in \phi_p^0$  such that  $\varphi_0 f_0 + \dots + \varphi_n f_n = 0$  and so

$$(2.4) \quad \begin{cases} \varphi_0 f_0 + \dots + \varphi_n f_n = 0 \\ \varphi_0 \hat{f}_0 + \dots + \varphi_n \hat{f}_n = 0 \\ \vdots \\ \varphi_0 \hat{f}_0^{[n]} + \dots + \varphi_n \hat{f}_n^{[n]} = 0. \end{cases}$$

Since (2.4) has a nontrivial solution, we get  $p$ -Casorati determinant

$$C_p(f_0, \dots, f_n) \equiv 0.$$

We apply induction on  $n$  to prove the converse assertion.

In the case when  $n = 1$ , suppose that  $C_p(f_0, f_1) \equiv 0$ . We consider the system of equations

$$(2.5) \quad \begin{cases} \varphi_0 f_0 + \varphi_1 f_1 = 0 \\ \varphi_0 \hat{f}_0 + \varphi_1 \hat{f}_1 = 0. \end{cases}$$

Since  $C_p(f_0, f_1) \equiv 0$ , it is easy to see that  $\varphi_0 = \frac{f_1}{f_0}$ ,  $\varphi_1 = -1$  is a solution of (2.5). Moreover, by assumption  $\sigma(f) = 0$ , we have  $\sigma(\tilde{f}) = 0$  where  $\tilde{f} := (f_0 : f_1)$ . Then the order of  $\varphi_0$  satisfies  $\sigma(\varphi_0) = \sigma\left(\frac{f_1}{f_0}\right) \leq \sigma(\tilde{f}) \leq \sigma(f) = 0$ .

Obviously,  $\varphi_1 = -1 \in \phi_p^0$  and  $\varphi_0 = \frac{f_1}{f_0} = \frac{\hat{f}_1}{\hat{f}_0}$ . Therefore, we also have  $\varphi_0 \in \phi_p^0$ .

This implies that  $f_0, f_1$  are linearly dependent over  $\phi_p^0$ .

Suppose now that  $C_p(f_0, \dots, f_j) \equiv 0$  implies that  $f_0, \dots, f_j$  are linearly dependent over  $\phi_p^0$  for all  $j \in \{1, \dots, k-1\}$ , where  $k \leq n$  and assume that  $C_p(f_0, \dots, f_k) \equiv 0$ . Then the linear system

$$(2.6) \quad \begin{cases} \varphi_0 f_0 + \dots + \varphi_{k-1} f_{k-1} = f_k \\ \varphi_0 \hat{f}_0 + \dots + \varphi_{k-1} \hat{f}_{k-1} = \hat{f}_k \\ \vdots \\ \varphi_0 \hat{f}_0^{[k-1]} + \dots + \varphi_{k-1} \hat{f}_{k-1}^{[k-1]} = \hat{f}_k^{[k-1]} \\ \varphi_0 \hat{f}_0^{[k]} + \dots + \varphi_{k-1} \hat{f}_{k-1}^{[k]} = \hat{f}_k^{[k]}, \end{cases}$$

where we have made the choice  $\varphi_k = -1$ . If  $C_p(f_0, \dots, f_{k-1}) \equiv 0$ , then  $f_0, \dots, f_{k-1}$  are linearly dependent over  $\phi_p^0$  by the induction assumption. Thus

also  $f_0, \dots, f_{k-1}, f_k$  are linearly dependent over  $\phi_p^0$ . If  $C_p(f_0, \dots, f_{k-1}) \neq 0$ , then by Cramer's rule for each  $i = 0, \dots, k-1$ , we have

$$\varphi_i = \frac{C_p(f_0, \dots, f_{i-1}, f_k, f_{i+1}, \dots, f_{k-1})}{C_p(f_0, \dots, f_{k-1})},$$

where  $f_k$  occurs in the  $i^{\text{th}}$  entry of  $p$ -Casorati determinant in the numerator instead of  $f_i$ . By writing

$$\varphi_i = \frac{f_i \hat{f}_i \cdots \hat{f}_i^{[k-1]} \cdot C_p\left(\frac{f_0}{f_i}, \dots, \frac{f_{i-1}}{f_i}, \frac{f_k}{f_i}, \frac{f_{i+1}}{f_i}, \dots, \frac{f_{k-1}}{f_i}\right)}{f_k \hat{f}_k \cdots \hat{f}_k^{[k-1]} \cdot C_p\left(\frac{f_0}{f_k}, \dots, \frac{f_{k-1}}{f_k}\right)},$$

it can be seen that

$$T(r, \varphi_i) = O\left(\sum_{j=0}^k \sum_{l=0}^{k-1} \left(T\left(r, \frac{\hat{f}_j^{[l]}}{\hat{f}_i^{[l]}}\right) + T\left(r, \frac{\hat{f}_j^{[l]}}{\hat{f}_k^{[l]}}\right)\right)\right)$$

for all  $i = 0, \dots, k-1$ . Now by Lemma 12, we have  $T(r, \hat{f}) = T(r, f) + o(T(r, f))$  for all meromorphic mappings  $f(z)$  with  $\sigma(f) = 0$ , and it follows that  $\sigma(\varphi_i) = 0$  for all  $i = 0, \dots, k-1$ .

We still need to prove that  $\varphi_i$  satisfies  $\varphi_i(pz) = \varphi_i(z)$  for all  $i = 0, \dots, k-1$ . By applying the operator  $\hat{\Delta}_p$  to  $k$  equations in the system (2.6), where  $\hat{\Delta}_p f = \hat{f} - f$ , it follows that

$$(2.7) \quad \begin{cases} (\varphi_0 \hat{\Delta}_p f_0 + \cdots + \varphi_{k-1} \hat{\Delta}_p f_{k-1}) + (\hat{f}_0 \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1} \hat{\Delta}_p \varphi_{k-1}) = \hat{\Delta}_p f_k \\ (\varphi_0 \hat{\Delta}_p \hat{f}_0 + \cdots + \varphi_{k-1} \hat{\Delta}_p \hat{f}_{k-1}) + (\hat{f}_0 \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1} \hat{\Delta}_p \varphi_{k-1}) = \hat{\Delta}_p \hat{f}_k \\ \vdots \\ (\varphi_0 \hat{\Delta}_p \hat{f}_0^{[k-1]} + \cdots + \varphi_{k-1} \hat{\Delta}_p \hat{f}_{k-1}^{[k-1]}) + (\hat{f}_0^{[k]} \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1}^{[k]} \hat{\Delta}_p \varphi_{k-1}) = \hat{\Delta}_p \hat{f}_k^{[k-1]}. \end{cases}$$

On the other hand also from (2.6), we have

$$(2.8) \quad \begin{cases} \varphi_0 \hat{\Delta}_p f_0 + \cdots + \varphi_{k-1} \hat{\Delta}_p f_{k-1} = \hat{\Delta}_p f_k \\ \varphi_0 \hat{\Delta}_p \hat{f}_0 + \cdots + \varphi_{k-1} \hat{\Delta}_p \hat{f}_{k-1} = \hat{\Delta}_p \hat{f}_k \\ \vdots \\ \varphi_0 \hat{\Delta}_p \hat{f}_0^{[k-1]} + \cdots + \varphi_{k-1} \hat{\Delta}_p \hat{f}_{k-1}^{[k-1]} = \hat{\Delta}_p \hat{f}_k^{[k-1]}. \end{cases}$$

Together (2.7) with (2.8), we get

$$\begin{cases} \hat{f}_0 \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1} \hat{\Delta}_p \varphi_{k-1} = 0 \\ \hat{f}_0 \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1} \hat{\Delta}_p \varphi_{k-1} = 0 \\ \vdots \\ \hat{f}_0^{[k]} \hat{\Delta}_p \varphi_0 + \cdots + \hat{f}_{k-1}^{[k]} \hat{\Delta}_p \varphi_{k-1} = 0, \end{cases}$$

which has only trivial solution. Therefore,  $\hat{\Delta}_p \varphi_0 \equiv \cdots \equiv \hat{\Delta}_p \varphi_{k-1} \equiv 0$ . It follows that  $\varphi_i(pz) = \varphi_i(z)$  for all  $i = 0, \dots, k-1$ . We finish the proof of the lemma 13.  $\square$

### 3. The proof of Theorem 1

We recall the lemma due to Nochka (see [5, 6, 13, 14]) as follows.

**Lemma 14.** *Let  $H_1, \dots, H_q$  ( $q > 2N - n + 1$ ) be hyperplanes in  $\mathbb{P}^n(\mathbb{C})$  located in  $N$ -subgeneral position. Then there exists a function  $\omega : \{1, \dots, q\} \rightarrow (0, 1]$  called a Nochka weight and a real number  $\tilde{\omega} \geq 1$  called a Nochka constant satisfying the following conditions:*

- (i) *If  $j \in \{1, \dots, q\}$ , then  $0 < \omega(j)\tilde{\omega} \leq 1$ .*
- (ii)  *$q - 2N + n - 1 = \tilde{\omega}(\sum_{j=1}^q \omega(j) - n - 1)$ .*
- (iii) *For  $R \subset \{1, \dots, q\}$  with  $|R| = N + 1$ , then  $\sum_{i \in R} \omega(i) \leq n + 1$ .*
- (iv)  *$\frac{N}{n} \leq \tilde{\omega} \leq \frac{2N-n+1}{n+1}$ .*
- (v) *Given real numbers  $\lambda_1, \dots, \lambda_q$  with  $\lambda_j \geq 1$  for  $1 \leq j \leq q$  and given any  $R \subset \{1, \dots, q\}$  and  $|R| = N + 1$ , there exists a subset  $R^1 \subset R$  such that  $|R^1| = \text{rank}\{H_i\}_{i \in R^1} = n + 1$  and*

$$\prod_{j \in R} \lambda_j^{\omega(j)} \leq \prod_{i \in R^1} \lambda_i.$$

**Lemma 15.** *Let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be an linearly nondegenerate meromorphic mapping over  $\phi_p^0$ , and  $H_j, j \in Q = \{1, \dots, q\}$  are hyperplanes, located in  $N$ -subgeneral position in  $\mathbb{P}^n(\mathbb{C})$ . Let  $\omega(j)$  be the Nochka weights of  $\{H_j\}_{j \in Q}$ . Assume that  $q > 2N - n + 1$ . Then we get*

$$\begin{aligned} & \|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot \prod_{j=0}^n \|\hat{f}^{[j]}\|^{-1} \\ & \leq K \cdot \frac{\prod_{t_j \in R} |H_j(\hat{f}^{[j]})|^{\omega(t_j)} \cdot \prod_{j \in S} |H_j(f)|^{\omega(j)}}{|C_p(f)|} \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |H_{t_j}(\hat{f}^{[j]})|} \end{aligned}$$

for an arbitrarily  $z \in \mathbb{C}^m \setminus \left( \{z \in \mathbb{C}^m : \prod_{t_j \in R} H_{t_j}(\hat{f}^{[j]}) \cdot \prod_{j \in S} H_j(f) = 0\} \cup I(f) \right)$ , where  $I(f) = \{z \in \mathbb{C}^m : f_0(z) = \cdots = f_n(z) = 0\}$  and  $K$  depends on  $\{H_j\}_{j \in Q}$ , and  $R^0, R, S$  are some subsets of  $Q$  such that

$$R^0 = \{t_0, t_1, \dots, t_n\} \subset R = \{t_0, t_1, \dots, t_n, t_{n+1}, \dots, t_N\} \subset Q \setminus S.$$

*Proof.* Since the hyperplanes  $\{H_j\}_{j=1}^q$  are in  $N$ -subgeneral position of  $\mathbb{P}^n(\mathbb{C})$ , we have  $\bigcap_{j \in R} H_j = \emptyset$  for any  $R \subset Q$  with  $|R| = N + 1$ . This implies that there exists a subset  $S \subset Q$  with  $|S| = q - N - 1$  such that  $\prod_{j \in S} H_j(\omega) \neq 0$ .

For each  $j \in S$ , we consider function  $h_j(\omega) = \frac{|H_j(\omega)|}{\|\omega\|}$  with  $\omega \in \mathbb{P}^n(\mathbb{C})$ . It is a positive continuous function on  $\mathbb{P}^n(\mathbb{C})$ . By the compactness of  $\mathbb{P}^n(\mathbb{C})$ , there

exists a positive constant  $K_j$  such that  $\frac{1}{K_j} \leq h_j(\omega) \leq K_j$ . Therefore, we have

$$(3.9) \quad \frac{1}{K_j} \leq \frac{|H_j(\hat{f}^{[k_j]})|}{\|\hat{f}^{[k_j]}\|} \leq K_j$$

for each  $j \in S$ ,  $k_j \in \mathbb{N}^*$ . It is easy to see that for each  $j \in Q \setminus S$  and  $k_j \in \mathbb{N}^*$ , there exists a positive constant  $K_j$  such that

$$\frac{|H_j(\hat{f}^{[k_j]})|}{\|\hat{f}^{[k_j]}\|} \leq K_j.$$

Put  $R = Q \setminus S$ . Then  $|R| = N + 1$ . Choose  $R^0 \subset R$  such that  $|R^0| = n + 1$  and  $R^0$  satisfies Lemma 14(v) with respect to numbers  $\frac{\|\hat{f}^{[k_j]}\|K_j}{|H_j(\hat{f}^{[k_j]})|}$  for arbitrary fixed point  $z \in \mathbb{C}^m \setminus \left( \{z \in \mathbb{C}^m : \prod_{j \in Q} |H_j(\hat{f}^{[k_j]})| = 0\} \cup I(f) \right)$  and  $k_j \in \mathbb{N}$ . We may assume that

$$R = \{t_0, t_1, \dots, t_n, t_{n+1}, \dots, t_N\} \text{ and } R^0 = \{t_0, t_1, \dots, t_n\}.$$

For  $Q$ , we can rewrite its elements as follows.

$$Q = \{t_0, t_1, \dots, t_n, t_{n+1}, \dots, t_N, t_{N+1}, \dots, t_{q-1}\}.$$

Then

$$(3.10) \quad \prod_{t_j \in R} \left( \frac{\|\hat{f}^{[j]}\|K_{t_j}}{|H_{t_j}(\hat{f}^{[j]})|} \right)^{\omega(t_j)} \leq \prod_{t_j \in R^0} \frac{\|\hat{f}^{[j]}\|K_{t_j}}{|H_{t_j}(\hat{f}^{[j]})|}.$$

Since  $f$  is linearly nondegenerate over field  $\phi_p^0$ , it follows from Lemma 13 that the Casorati determinant  $C_p(f) \neq 0$ . By  $\text{rank}\{H_{t_j}\}_{j \in R^0} = n + 1$ , there exists a positive constant  $K_{R^0}$  such that  $|C_p(f)| = K_{R^0} \cdot |C_p(H_j(f) : j \in R^0)|$ . Thus

$$(3.11) \quad \frac{K_{R^0} \cdot |C_p(H_j(f) : j \in R^0)|}{|C^d(f)|} = 1.$$

Since (3.9) and (3.10), for an arbitrarily

$$z \in G := \mathbb{C}^m \setminus \left( \{z \in \mathbb{C}^m : \prod_{t_j \in R} H_j(\hat{f}^{[j]}) \cdot \prod_{j \in S} H_j(f) = 0\} \cup I(f) \right),$$

we have

$$(3.12) \quad \prod_{j \in S} \left( \frac{1}{K_j^2} \right)^{\omega(j)} \leq \prod_{j \in S} \left( \frac{|H_j(f)|}{\|f\|K_j} \right)^{\omega(j)}$$

$$\begin{aligned}
&\leq \prod_{t_j \in R} \left( \frac{\|\hat{f}^{[j]}\| K_{t_j}}{|H_{t_j}(\hat{f}^{[j]})|} \right)^{\omega(t_j)} \cdot \frac{\prod_{t_j \in R} |H_{t_j}(\hat{f}^{[j]})|^{\omega(t_j)} \prod_{j \in S} |H_j(f)|^{\omega(j)}}{\|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot K_0^{\sum_{j=1}^q \omega(j)}} \\
&\leq \prod_{t_j \in R^0} \frac{\|\hat{f}^{[j]}\| K_{t_j}}{|H_{t_j}(\hat{f}^{[j]})|} \cdot \frac{\prod_{t_j \in R} |H_{t_j}(\hat{f}^{[j]})|^{\omega(t_j)} \prod_{j \in S} |H_j(f)|^{\omega(j)}}{\|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot K_0^{\sum_{j=1}^q \omega(j)}} \\
&= \frac{\prod_{t_j \in R^0} K_{t_j}}{K_0^{\sum_{j=1}^q \omega(j)}} \cdot \frac{\prod_{t_j \in R} |H_{t_j}(\hat{f}^{[j]})|^{\omega(t_j)} \prod_{j \in S} |H_j(f)|^{\omega(j)}}{|H_{t_0}(f) \cdot H_{t_1}(\hat{f}) \cdots H_{t_n}(\hat{f}^{[n]})|} \\
&\quad \times \frac{1}{\|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot \prod_{t_j \in R^0} \|\hat{f}^{[j]}\|^{-1}},
\end{aligned}$$

where  $K_0 := \min\{K_1, \dots, K_q\}$ . Together (3.11) with (3.12), for  $z \in G$ , we have

$$\begin{aligned}
&\prod_{j \in S} \left( \frac{1}{K_j^2} \right)^{\omega(j)} \\
&\leq \frac{\prod_{t_j \in R^0} K_{t_j} \cdot K_{R^0}}{K_0^{\sum_{j=1}^q \omega(j)}} \cdot \frac{1}{\|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot \prod_{t_j \in R^0} \|\hat{f}^{[j]}\|^{-1}} \\
&\quad \times \frac{\prod_{t_j \in R} |H_{t_j}(\hat{f}^{[j]})|^{\omega(t_j)} \prod_{j \in S} |H_j(f)|^{\omega(j)}}{|C_p(f)|} \cdot \frac{|C_p(H_j(f) : j \in R^0)|}{|H_{t_0}(f) \cdot H_{t_1}(\hat{f}) \cdots H_{t_n}(\hat{f}^{[n]})|}.
\end{aligned}$$

It implies that

$$\begin{aligned}
&\|f\|^{\sum_{j \in S} \omega(j)} \cdot \prod_{t_j \in R} \|\hat{f}^{[j]}\|^{\omega(t_j)} \cdot \prod_{j=0}^n \|\hat{f}^{[j]}\|^{-1} \\
&\leq \frac{\prod_{t_j \in R^0} K_{t_j} \cdot K_{R^0} \cdot \prod_{j \in S} (K_j)^{2\omega(j)}}{K_0^{\sum_{j=1}^q \omega(j)}} \cdot \frac{\prod_{t_j \in R} |H_{t_j}(\hat{f}^{[j]})|^{\omega(t_j)} \prod_{j \in S} |H_j(f)|^{\omega(j)}}{|C_p(f)|} \\
&\quad \times \frac{|C_p(H_j(f) : j \in R^0)|}{|H_{t_0}(f) \cdot H_{t_1}(\hat{f}) \cdots H_{t_n}(\hat{f}^{[n]})|}
\end{aligned}$$

for an arbitrarily  $z \in G$ . We obtain Lemma 15 by setting

$$K = \frac{\prod_{t_j \in R^0} K_{t_j} \cdot K_{R^0} \cdot \prod_{j \in S} (K_j)^{2\omega(j)}}{K_0^{\sum_{j=1}^q \omega(j)}}$$

which is a positive constant depending on  $\{H_j\}_{j=1}^q$ ,  $R^0$ ,  $R$  and  $S$ . We finish the proof of Lemma 15.  $\square$

*Proof of Theorem 1.* By Lemma 15, for  $r > 1$ , we have

$$\begin{aligned} & \sum_{j \in S} \omega(j) \log \|f\| + \sum_{t_j \in R} \omega(t_j) \log \|\hat{f}^{[j]}\| - \sum_{j=0}^n \log \|\hat{f}^{[j]}\| \\ & \leq \sum_{t_j \in R} \omega(t_j) \log |H_{t_j}(\hat{f}^{[j]})| + \sum_{j \in S} \omega(j) \log |H_j(f)| - \log |C_p(f)| \\ & \quad + \log \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} + O(1). \end{aligned}$$

Integrating both sides of this inequality and using Jensen's theorem and by definition of the characteristic function of  $f$ , we have

$$\begin{aligned} (3.13) \quad & \sum_{j \in S} \omega(j) T_f(r) + \sum_{t_j \in R} \omega(t_j) T_{\hat{f}^{[j]}}(r) - \sum_{j=0}^n T_{\hat{f}^{[j]}}(r) \\ & \leq \sum_{t_j \in R} \omega(t_j) N(r, \nu_{H_{t_j}(f)}^0) + \sum_{j \in S} \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) \\ & \quad + \int_{S_m(r)} \log^+ \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} \eta_m(z) + O(1) \\ & \leq \sum_{t_j \in R} \omega(t_j) N(|p|r, \nu_{H_{t_j}(f)}^0) + \sum_{j \in S} \omega(j) N(r, \nu_{Q_j(f)}^0) - N(r, \nu_{C^d(f)}^0) \\ & \quad + \int_{S_m(r)} \log^+ \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} \eta_m(z) + O(1). \end{aligned}$$

By the First Main Theorem, the order of  $N(r, \nu_{H_j(f)}^0)$  satisfies

$$\limsup_{r \rightarrow \infty} \frac{\log^+ N(r, \nu_{H_{t_j}(f)}^0)}{\log r} \leq \limsup_{r \rightarrow \infty} \frac{\log^+ T(r, f)}{\log r} = \sigma(f) = 0.$$

So, by Lemma 8, the below inequality holds on a set of logarithmic density 1

$$\begin{aligned} N(|p|r, \nu_{H_{t_j}(f)}^0) &= N(r, \nu_{H_{t_j}(f)}^0) + o\left(N(r, \nu_{H_{t_j}(f)}^0)\right) \\ &\leq N(r, \nu_{H_{t_j}(f)}^0) + o(T(r, f)). \end{aligned}$$

It follows from (3.13) that

$$\begin{aligned} (3.14) \quad & \sum_{j \in S} \omega(j) T_f(r) + \sum_{t_j \in R} \omega(t_j) T_{\hat{f}^{[j]}}(r) - \sum_{j=0}^n T_{\hat{f}^{[j]}}(r) \\ & \leq \sum_{t_j \in R} \omega(t_j) N(r, \nu_{H_{t_j}(f)}^0) + \sum_{j \in S} \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) \\ & \quad + \int_{S_m(r)} \log^+ \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} \eta_m(z) + o(T(r, f)) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j \in Q} \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) \\
&\quad + \int_{S_m(r)} \log^+ \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} \eta_m(z) + o(T(r, f)).
\end{aligned}$$

We have

$$\begin{aligned}
\frac{C_p(H_j(f) : j \in R^0)}{\prod_{t_j \in R^0} |(Q_{t_j}(\hat{f}^{[j]}))|} &= \frac{\begin{vmatrix} 1 & \frac{H_{t_1}(f)}{H_{t_0}(f)} & \cdots & \frac{H_{t_n}(f)}{H_{t_0}(f)} \\ 1 & \frac{H_{t_1}(\hat{f})}{H_{t_0}(\hat{f})} & \cdots & \frac{H_{t_n}(\hat{f})}{H_{t_0}(\hat{f})} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \frac{H_{t_1}(\hat{f}^{[n]})}{H_{t_0}(\hat{f}^{[n]})} & \cdots & \frac{H_{t_n}(\hat{f}^{[n]})}{H_{t_0}(\hat{f}^{[n]})} \end{vmatrix}}{\left| \frac{H_{t_1}(\hat{f})}{H_{t_0}(\hat{f})} \cdots \frac{H_{t_n}(\hat{f}^{[n]})}{H_{t_0}(\hat{f}^{[n]})} \right|} \\
&= \frac{\begin{vmatrix} 1 & 1 & \cdots & 1 \\ 1 & \frac{H_{t_1}(\hat{f})/H_{t_1}(f)}{H_{t_0}(\hat{f})/H_{t_0}(f)} & \cdots & \frac{H_{t_n}(\hat{f})/H_{t_n}(f)}{H_{t_0}(\hat{f})/H_{t_0}(f)} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \frac{H_{t_1}(\hat{f}^{[n]})/H_{t_1}(f)}{H_{t_0}(\hat{f}^{[n]})/H_{t_0}(f)} & \cdots & \frac{H_{t_n}(\hat{f}^{[n]})/H_{t_n}(f)}{H_{t_0}(\hat{f}^{[n]})/H_{t_0}(f)} \end{vmatrix}}{\left| \frac{H_{t_1}(\hat{f})/H_{t_1}(f)}{H_{t_0}(\hat{f})/H_{t_0}(f)} \cdots \frac{H_{t_n}(\hat{f}^{[n]})/H_{t_n}(f)}{H_{t_0}(\hat{f}^{[n]})/H_{t_0}(f)} \right|}.
\end{aligned}$$

It is easy to see that  $\sigma\left(\frac{H_i(f)}{H_j(f)}\right) \leq \sigma(f) = 0$  for all  $i, j$ . Therefore, by Lemma 11, we have

$$\begin{aligned}
\int_{S_m(r)} \log^+ \frac{|C_p(H_j(f) : j \in R^0)|}{\prod_{t_j \in R^0} |(H_{t_j}(\hat{f}^{[j]}))|} \eta_m(z) &\leq \sum_{j=1}^n o\left(T\left(r, \frac{H_{t_j}(\hat{f}^{[j]})}{H_{t_0}(\hat{f}^{[j]})}\right)\right) \\
&= o(T(r, f))
\end{aligned}$$

on a set of logarithmic density 1. Hence, together this with (3.14), we get

$$\begin{aligned}
(3.15) \quad &\sum_{j \in S} \omega(j) T_f(r) + \sum_{t_j \in R} \omega(t_j) T_{\hat{f}^{[j]}}(r) - \sum_{j=0}^n T_{\hat{f}^{[j]}}(r) \\
&\leq \sum_{j \in Q} \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) + o(T(r, f))
\end{aligned}$$

on a set of logarithmic density 1. From (3.15) and Lemma 12, we get

$$(3.16) \quad \left( \sum_{j \in Q} \omega(j) - n - 1 \right) T(r, f) \leq \sum_{j \in Q} \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) + o(T(r, f))$$

on a set of logarithmic density 1.



By (i), (ii) and (iv) of Lemma 14, the inequality (3.16) implies that the below inequality holds on a set of logarithmic density 1

$$(q - 2N + n - 1)T(r, f) \leq \sum_{j \in Q} N(r, \nu_{H_j(f)}^0) - \frac{N}{n} N(r, \nu_{C_p(f)}^0) + o(T(r, f)).$$

The proof of Theorem 1 is completed.  $\square$

#### 4. The proof of Theorem 2

Let  $z_0$  be a  $n$ -successive zero with separation  $p$  of  $H_j(f)$  respect to the rescaling  $\tau_p(z) = pz$  for some  $j \in \{1, \dots, q\}$ . Since  $\{H_j\}_{j=1}^q$  is in  $N$ -subgeneral position, there are at most  $N$  functions  $H_j(f)$  vanishing at  $z_0$ . Without loss of generality, we may assume that  $z_0$  is a  $n$ -successive with separation  $p$  zero of  $H_j(f)$  respect to the rescaling  $\tau_p(z) = pz$  with all  $j \in A$  and  $z_0$  is a  $n$ -aperiodic zero with separation  $p$  of  $H_j(f)$  respect to the rescaling  $\tau_p(z) = pz$  with all  $j \in B$  and  $z_0$  is not a zero of  $H_j(f)$  with all  $j \notin A \cup B$ , where  $|A \cup B| = N$ . Take  $R \subset \{1, \dots, q\}$  containing  $A$  such that  $|R| = N + 1$  and  $R \cap B = \emptyset$ . Choose subset  $R^1 \subset R$  with  $|R^1| = \text{rank}\{H_j\}_{j \in R^1} = n + 1$  such that  $R^1$  satisfies (v) of Lemma 14 with respect to numbers  $\{\lambda_j = e^{\nu_{H_j(f)}^0(z_0)}\}_{j=1}^q$ . Then we have

$$\prod_{j \in R} e^{\omega(j)\nu_{H_j(f)}^0(z_0)} \leq \prod_{j \in R^1} e^{\nu_{H_j(f)}^0(z_0)}.$$

Therefore,

$$(4.17) \quad \sum_{j \notin B} \omega(j)\nu_{H_j(f)}^0(z_0) \leq \sum_{j \in A \cap R^1} \nu_{H_j(f)}^0(z_0).$$

By rearrangement index if necessary, we may assume that  $R^1 = \{t_0, \dots, t_n\}$  and  $A \cap R^1 = \{t_0, \dots, t_k\}$  with  $0 \leq k \leq n$ . Since  $\text{rank}\{\{H_{t_j}\}_{j=0}^n\} = n + 1$ , there exists a nonzero constant  $C_{R^1}$  such that

$$C_p(f) = C_{R^1} \cdot C_p(H_{t_0}(f), \dots, H_{t_n}(f)).$$

This deduces that  $\nu_{C_p(f)}^0 = \nu_{C_p(H_{t_0}(f), \dots, H_{t_n}(f))}^0$ . We have

$$\begin{aligned} & C_p(H_{t_0}(f), \dots, H_{t_n}(f)) \\ &= H_{t_0}(f) \cdots H_{t_k}(f) \\ & \times \begin{vmatrix} 1 & \cdots & 1 & H_{t_{k+1}}(f) & \cdots & H_{t_n}(f) \\ \frac{H_{t_0}(\hat{f})}{H_{t_0}(f)} & \cdots & \frac{H_{t_k}(\hat{f})}{H_{t_k}(f)} & H_{t_{k+1}}(\hat{f}) & \cdots & H_{t_n}(\hat{f}) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{H_{t_0}(\hat{f}^{[n]})}{H_{t_0}(f)} & \cdots & \frac{H_{t_k}(\hat{f}^{[n]})}{H_{t_k}(f)} & H_{t_{k+1}}(\hat{f}^{[n]}) & \cdots & H_{t_n}(\hat{f}^{[n]}) \end{vmatrix}. \end{aligned}$$

It follows that

$$\nu_{C_p(f)}^0(z_0) \geq \nu_{H_{t_0}(f)\dots H_{t_k}(f)}^0(z_0) = \sum_{j=0}^k \nu_{H_{t_j}(f)}^0(z_0).$$

Together this inequality with (4.17), we get

$$\nu_{C_p(f)}^0(z_0) \geq \sum_{j \notin B} \omega(j) \nu_{H_j(f)}^0(z_0).$$

This, by going through all points  $z_0 \in \mathbb{C}^m$  and by definitions of  $\hat{N}^{[n,p]}(r, H_j(f))$  implies that

$$\sum_{j=1}^q \omega(j) N(r, \nu_{H_j(f)}^0) - N(r, \nu_{C_p(f)}^0) \leq \sum_{j=1}^q \omega(j) \hat{N}^{[n,p]}(r, H_j(f)).$$

This and (3.16) yield

$$\left( \sum_{j=1}^q \omega(j) - n - 1 \right) T(r, f) \leq \sum_{j=1}^q \omega(j) \hat{N}^{[n,p]}(r, H_j(f)) + o(T(r, f))$$

on a set of logarithmic density 1. By (i), (ii) and (iv) of Lemma 14, the above inequality implies that

$$(q - 2N + n - 1) T(r, f) \leq \sum_{j=1}^q \hat{N}^{[n,p]}(r, H_j(f)) + o(T(r, f))$$

on a set of logarithmic density 1. The proof of Theorem 2 is completed.

### 5. The proof of Theorem 3

**Lemma 16.** *Let  $f : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  be a meromorphic mapping with reduce presentation  $f = (f_0 : \dots : f_n)$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0, 1$  for all  $j$ . Assume that  $\sigma(f) = 0$  and all zeros of  $f_0, \dots, f_n$  are forward invariant with respect to the rescaling  $\tau_p(z) = pz$ . If  $\frac{f_i}{f_j} \notin \phi_p^0$  for all  $i, j \in \{0, \dots, n\}$  such that  $i \neq j$ , then  $f_0, \dots, f_n$  are linearly independent over the field  $\phi_p^0$ .*

*Proof.* Assume that  $f$  is linearly degenerate over  $\phi_p^0$ . Without loss generality we assume that there exist  $\varphi_0, \dots, \varphi_n \in \phi_p^0 \setminus \{0\}$  such that  $\varphi_0 f_0 + \dots + \varphi_{n-1} f_{n-1} = \varphi_n f_n$ . Since all zeros of  $f_0, \dots, f_n$  are forward invariant with respect to the rescaling  $\tau_p(z) = pz$  and since  $\varphi_0, \dots, \varphi_n \in \phi_p^0 \setminus \{0\}$ , we can choose a meromorphic  $h$  such that  $h\varphi_0 f_0, \dots, h\varphi_n f_n$  are holomorphic functions on  $\mathbb{C}^m$  without common zeros and such that preimages of all zeros of  $h\varphi_0 f_0, \dots, h\varphi_n f_n$  are forward invariant with respect to the rescaling  $\tau_p(z) = pz$ . Then we get

$$(5.18) \quad \lim_{r \rightarrow \infty} \frac{\log^+ (N(r, \nu_h^0) + N(r, \nu_h^\infty))}{\log r} = 0$$

and  $h\varphi_0 f_0, \dots, h\varphi_{n-1} f_{n-1}$  can not have any common zeros.

Put  $g_i = h\varphi_i f_i$  for  $0 \leq i \leq n$  and  $G = (g_0 : \cdots : g_{n-1})$  is a holomorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^{n-1}(\mathbb{C})$ . Then by definition of characteristic function, we have

$$\begin{aligned} T(r, G) &= \int_{S_m(r)} \log \|G\| \eta_m(z) + O(1) \\ &\leq \int_{S_m(r)} \log |h| \eta_m(z) + \int_{S_m(r)} \log \left( \sum_{j=0}^{n-1} |f_j|^2 \right)^{\frac{1}{2}} \eta_m(z) \\ &\quad + \sum_{j=0}^{n-1} \int_{S_m(r)} \log |\varphi_j| \eta_m(z) + O(1) \\ &\leq N(r, \nu_h^0) + N(r, \nu_h^\infty) + T_f(r) + \sum_{j=0}^{n-1} T_{\varphi_j}(r) + O(1). \end{aligned}$$

This together (5.18) deduce that  $\sigma(G) = 0$ .

Assume that  $G : \mathbb{C}^m \rightarrow \mathbb{P}^n(\mathbb{C})$  is linearly nondegenerate over  $\phi_p^0$ . Since Lemma 13, it follows that  $C_p(g_0, \dots, g_{n-1}) \neq 0$ . Take  $n+1$  hyperplanes

$$H_0 : \omega_0 = 0, H_1 : \omega_1 = 0, \dots, H_{n-1} : \omega_{n-1} = 0$$

and

$$H_n : \omega_0 + \cdots + \omega_{n-1} = 0,$$

where  $(\omega_0, \dots, \omega_{n-1})$  is homogeneous coordinate system of  $\mathbb{P}^{n-1}(\mathbb{C})$ . So  $(G, H_j) = g_j$  for  $0 \leq j \leq n-1$  and  $(G, H_n) = g_0 + \cdots + g_{n-1} = h\varphi_n f_n = g_n$ . Obviously,  $\{H_j\}_{j=0}^n$  are in general position in  $\mathbb{P}^{n-1}(\mathbb{C})$ . Applying Theorem 2, we have

$$T(r, G) \leq \sum_{j=0}^n \hat{N}^{[n,p]}(r, H_j(G)) + o(T(r, G))$$

on a set of logarithmic density 1. Since all zeros of  $H_j(G) = (G, H_j) = g_j$  ( $0 \leq j \leq n$ ) are forward invariant with respect to the rescaling  $\tau_p(z) = pz$ ,  $\hat{N}^{[n,p]}(r, H_j(G)) \equiv 0$  and therefore,  $T(r, G) \leq o(T(r, G))$  on a set of logarithmic density 1. This is a contradiction. It follows that  $G$  is linearly dependent over  $\phi_p^0$ . Thus there exist  $\psi_0, \dots, \psi_{n-1}$  satisfying

$$\psi_0 f_0 + \cdots + \psi_{n-2} f_{n-2} = \psi_{n-1} f_{n-1}$$

and not all  $\psi_i$  are identically zero. By continuing in this fashion it follows after at most  $n-2$  time, we have  $\frac{f_i}{f_j} \in \phi_p^0$  for some  $i \neq j$ . This is contradiction. Hence,  $f$  is linearly nondegenerate over  $\phi_p^0$ . We finish the proof of Lemma 16.  $\square$

**Lemma 17.** *Let  $f = (f_0 : \cdots : f_n)$  be a meromorphic mapping of  $\mathbb{C}^m$  to  $\mathbb{P}^n(\mathbb{C})$  such that  $\sigma(f) = 0$  and let  $p = (p_1, \dots, p_m) \in \mathbb{C}^m$  with  $p_j \neq 0, 1$  for*

all  $j$ . Assume that all zeros of  $f_0, \dots, f_n$  are forward invariant with respect to the rescaling  $\tau_p(z) = pz$ . Let  $S_1 \cup \dots \cup S_l$  be the partition of  $\{0, \dots, n\}$  formed in such a way that  $i$  and  $j$  are in the same class  $S_k$  if and only if  $\frac{f_i}{f_j} \in \phi_p^0$ . If  $f_0 + \dots + f_n = 0$ , then  $\sum_{j \in S_k} f_j = 0$  for all  $k \in \{1, \dots, l\}$ .

*Proof.* For each  $i \in S_k, k \in \{1, \dots, l\}$  we have  $f_i = \varphi_{i,j_k} f_{j_k}$  for  $\varphi_{i,j_k} \in \phi_p^0$  whenever the  $i, j_k \in S_k$ . It implies that

$$0 = \sum_{k=0}^n f_k = \sum_{k=1}^l \sum_{i \in S_k} \varphi_{i,j_k} f_{j_k} = \sum_{k=1}^l B_k f_{j_k},$$

where  $B_k = \sum_{i \in S_k} \varphi_{i,j_k} \in \phi_p^0$ . This deduces that  $f_{j_1}, \dots, f_{j_l}$  are linearly dependent over  $\phi_p^0$  if not all  $B_k$  are identically zeros. This contradicts to the Lemma 16. Then  $B_k \equiv 0$  for all  $k \in \{1, \dots, l\}$ . Thus  $\sum_{i \in S_k} f_i = \sum_{i \in S_k} \varphi_{i,j_k} f_{j_k} = B_k f_{j_k} \equiv 0$  for all  $k \in \{1, \dots, l\}$ . Lemma 17 is proved.  $\square$

*Proof of Theorem 3.* By assumptions of the theorem, holomorphic functions

$$G_j = H_j(f) = \sum_{i=0}^n a_{ji} f_i,$$

satisfying

$$\{\tau_p(G_j^{-1}(0))\} \subset \{G_j^{-1}(0)\}, \quad j \in \{1, \dots, q\},$$

where  $H_j : \sum_{i=0}^n a_{ji} \omega_i = 0$ , and  $\{\cdot\}$  denotes a multiset with counting multiplicities of its elements. We say that  $i \sim j$  if  $G_i = \alpha G_j$  for some  $\alpha \in \phi_p^0 \setminus \{0\}$ . Therefore, the set of indexes  $\{1, \dots, q\}$  may be split into disjoint equivalence classes  $S_j$ ,

$$\{1, \dots, q\} = \cup_{j=1}^l S_j$$

for some  $l \leq q$ .

The first, we assume that  $S_j$  has as most  $q - N - 1$  elements for some  $j \in \{1, \dots, l\}$ . Put  $R = Q \setminus S_j$  then,  $|R| \geq N + 1$ . Let  $s_0 \in S_j$  and put  $U = R \cup \{s_0\}$ . Without loss of generality, we may assume that  $U = \{s_0, \dots, s_{N+1}\}$ . Then since the  $\{H_j\}_{j=1}^q$  are in  $N$ -subgeneral position, there exist  $\alpha_j \in \mathbb{C} \setminus \{0\}$  such that  $\sum_{j=0}^{N+1} \alpha_j H_{s_j} = 0$  and therefore, we have  $\sum_{j=0}^{N+1} \alpha_j H_{s_j}(f) = \sum_{j=0}^{N+1} \alpha_j G_{s_j} \equiv 0$ . By assumptions of the theorem, we can see that all zeros of  $\alpha_j G_{s_j}$  are forward invariant with respect to the rescaling  $\tau_p(z) = pz$ . We have

$$G := (\alpha_0 G_{s_0} : \dots : \alpha_{N+1} G_{s_{N+1}})$$

is a meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^{N+1}(\mathbb{C})$  with its order  $\sigma(G) \leq \sigma(f) = 0$ . By Lemma 17, we have  $\alpha_0 G_{s_0} \equiv 0$ . Hence,  $H_{s_0}(f) \equiv 0$ . This implies that the image  $f(\mathbb{C}^m)$  is included in the hyperplane  $H_{s_0}$  of  $\mathbb{P}^n(\mathbb{C})$ . We may consider  $f$  be a meromorphic mapping of  $\mathbb{C}^m$  into  $\mathbb{P}^{n-1}(\mathbb{C})$ .

The second, we assume that  $S_j$  has at least  $q - N$  elements for all  $j \in \{1, \dots, l\}$ . Then

$$l \leq \frac{q}{q - N}.$$

Since  $\{H_j\}_{j=1}^q$  is in  $N$ -subgeneral position, we can choose a subset  $V \subset \{1, \dots, q\}$  with  $|V| = n + 1$  such that  $\{H_j\}_{j \in V}$  is linearly independent. Put  $V_j = V \cap S_j$  for each  $1 \leq j \leq l$ . Then we have  $V = \cup_{j=1}^l V_j$ . Since each  $V_j$  gives rise to  $|V_j| - 1$  equations over the field  $\phi_p^0$ , it is easy to see that there are at least

$$\sum_{j=1}^l (|V_j| - 1) = n + 1 - l \geq n + 1 - \frac{q}{q - N} = n - \frac{N}{q - N}$$

linearly independent relations over the field  $\phi_p^0$ . It follows that the image of  $f$  is contained in a projective linear subspace over  $\phi_p^0$  of dimension  $\leq \left\lfloor \frac{N}{q - N} \right\rfloor$ .

Obviously, if  $q \geq 2N + 1$  then  $\left\lfloor \frac{N}{q - N} \right\rfloor = 0$ , and therefore  $f(z) = f(pz)$ . The Theorem 3 is proved.  $\square$

### References

- [1] D. C. Barnett, R. G. Halburd, R. J. Korhonen, and W. Morgan, *Nevanlinna theory for the  $q$ -difference operator and meromorphic solutions of  $q$ -difference equations*, Proc. Roy. Soc. Edinburgh Sect. A. **137** (2007), no. 3, 457–474.
- [2] T. B. Cao, *Difference analogues of the second main theorem for meromorphic functions in several complex variables*, Math. Nachr. **287** (2014), no. 5-6, 530–545.
- [3] T. B. Cao and R. Korhonen, *A new version of the second main theorem for meromorphic mappings intersecting hyperplanes in several complex variables*, J. Math. Anal. Appl. **444** (2016), no. 2, 1114–1132.
- [4] H. Cartan, *Sur les zeros des combinaisons linéaires de pfonctions holomorphes données*, Mathematica Cluj **7** (1933), 531.
- [5] H. Fujimoto, *The uniqueness problem of meromorphic maps into the complex projective space*, Nagoya Math. J. **58** (1975), 1–23.
- [6] ———, *Nonintegrated defect relation for meromorphic maps of complete Kähler manifolds into  $\mathbb{P}^{N_1}(\mathbb{C}) \times \dots \times \mathbb{P}^{N_k}(\mathbb{C})$* , Japan. J. Math. (N.S.) **11** (1985), no. 2, 233–264.
- [7] R. G. Halburd and R. J. Korhonen, *Nevanlinna theory for the difference operator*, Ann. Acad. Sci. Fenn. Math. **31** (2006), no. 2, 463–478.
- [8] R. Halburd, R. Korhonen, and K. Tohge, *Holomorphic curves with shift-invariant hyperplane preimages*, Trans. Amer. Math. Soc. **366** (2014), no. 8, 4267–4298.
- [9] W. K. Hayman, *On the characteristic of functions meromorphic in the plane and of their integrals*, Proc. Lond. Math. Soc. (3) **14** (1965), 93–128.
- [10] P.-C. Hu, P. Li, and C.-C. Yang, *Unicity of Meromorphic Mappings*, Vol. 1 of Advances in Complex Analysis and Its Applications, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [11] Z.-B. Huang, *Value distribution and uniqueness on  $q$ -differences of meromorphic functions*, Bull. Korean Math. Soc. **50** (2013), no. 4, 1157–1171.
- [12] R. Korhonen, *A difference Picard theorem for meromorphic functions of several variables*, Comput. Methods Funct. Theory **12** (2012), no. 1, 343–361.
- [13] E. I. Nochka, *On the theory of meromorphic functions*, Sov. Math. Dokl. **27** (1983), 377–381.

- [14] J. Noguchi, *A note on entire pseudo-holomorphic curves and the proof of Cartan-Nochka's theorem*, Kodai Math. J. **28** (2005), no. 2, 336–346.
- [15] X. Qi, K. Liu, and L. Yang, *Value results of a meromorphic function  $f(z)$  and  $f(qz)$* , Bull. Korean Math. Soc. **48** (2011), no. 6, 1235–1243.
- [16] Z. T. Wen, *The  $q$ -difference theorems for meromorphic functions of several variables*, Abstr. Appl. Anal. **2014** (2014), ID 736021, 6 pp.
- [17] Z.-T. Wen and Z. Ye, *Wiman-Valiron theorem for  $q$ -difference*, Annales Academiæ Scientiarum Fennicæ Mathematica **41** (2016), 305–312.
- [18] P. M. Wong, H. F. Law, and P. P. W. Wong, *A second main theorem on  $\mathbb{P}^n$  for difference operator*, Sci. China Ser. A **52** (2009), no. 12, 2751–758.
- [19] J. Zhang and R. Korhonen, *On the Nevanlinna characteristic of  $f(qz)$  and its applications*, J. Math. Anal. Appl. **369** (2010), no. 2, 537–544.

THI TUYET LUONG  
DEPARTMENT OF MATHEMATICS  
NATIONAL UNIVERSITY OF CIVIL ENGINEERING  
55 GIAI PHONG STR., HANOI, VIETNAM  
*E-mail address:* luongtuyetdtxd@gmail.com

DANG TUYEN NGUYEN  
DEPARTMENT OF MATHEMATICS  
NATIONAL UNIVERSITY OF CIVIL ENGINEERING  
55 GIAI PHONG STR., HANOI, VIETNAM  
*E-mail address:* tuyenn@nuce.edu.vn

DUC THOAN PHAM  
DEPARTMENT OF MATHEMATICS  
NATIONAL UNIVERSITY OF CIVIL ENGINEERING  
55 GIAI PHONG STR., HANOI, VIETNAM  
*E-mail address:* thoanpd@nuce.edu.vn