

LOGHARMONIC MAPPINGS WITH TYPICALLY REAL ANALYTIC COMPONENTS

ZAYID ABDULHADI, NAJLA M. ALARIFI, AND ROSIHAN M. ALI

ABSTRACT. This paper treats the class of normalized logharmonic mappings $f(z) = zh(z)\overline{g(z)}$ in the unit disk satisfying $\varphi(z) = zh(z)g(z)$ is analytically typically real. Every such mapping f admits an integral representation in terms of its second dilatation function and a function of positive real part with real coefficients. The radius of starlikeness and an upper estimate for arclength are obtained. Additionally, it is shown that f maps the unit disk into a domain symmetric with respect to the real axis when its second dilatation has real coefficients.

1. Introduction

Let $\mathcal{H}(U)$ be the linear space of analytic functions defined in the unit disk $U = \{z : |z| < 1\}$ of the complex plane \mathbb{C} . Let B denote the set of self-maps $a \in \mathcal{H}(U)$, and B_0 its subclass consisting of $a \in B(U)$ with $a(0) = 0$. A logharmonic mapping in U is a solution of the nonlinear elliptic partial differential equation

$$(1.1) \quad \overline{\left(\frac{f_z(z)}{f(z)}\right)} = a(z)\frac{f_z(z)}{f(z)},$$

where the second dilatation function a lies in B . Thus the Jacobian

$$J_f = |f_z|^2(1 - |a|^2)$$

is positive, and all non-constant logharmonic mappings are sense-preserving and open in U .

If f is a non-constant logharmonic mapping which vanishes only at $z = 0$, then [4] shows that f admits the representation

$$(1.2) \quad f(z) = z^m |z|^{2\beta m} h(z)\overline{g(z)},$$

Received December 11, 2017; Revised April 1, 2018; Accepted May 16, 2018.

2010 *Mathematics Subject Classification.* Primary 30C35, 30C45.

Key words and phrases. logharmonic mappings, typically real functions, radius of starlikeness, arclength.

where m is a positive integer, $\operatorname{Re} \beta > -1/2$, and $h, g \in \mathcal{H}(U)$ satisfy $g(0) = 1$ and $h(0) \neq 0$. The exponent β in (1.2) depends only on $a(0)$ and is given by

$$\beta = \overline{a(0)} \frac{1 + a(0)}{1 - |a(0)|^2}.$$

Note that $f(0) \neq 0$ if and only if $m = 0$, and that a univalent logharmonic mapping vanishes at the origin if and only if $m = 1$, that is, f has the form

$$f(z) = z|z|^{2\beta} h(z) \overline{g(z)},$$

where $0 \notin (hg)(U)$. This class has been studied extensively over recent years in [1–8].

As further evidence of its importance, note that $F(\zeta) = \log f(e^\zeta)$ are univalent harmonic mappings of the half-plane $\{\zeta : \operatorname{Re} \zeta < 0\}$. Studies on univalent harmonic mappings can be found in [9–13, 15–17], which are closely related to the theory of minimal surfaces (see [19, 20]).

An analytic function φ in U is typically real if $\varphi(z)$ is real whenever z is real and nonreal elsewhere. Similarly, a logharmonic mapping f in U is typically real if $f(z)$ is real whenever z is real and nonreal elsewhere. Investigations into typically real logharmonic mappings was initiated by Abdulhadi in [2].

Denote by HG the class of analytic functions $\varphi(z) = zh(z)g(z)$, where h and g in $\mathcal{H}(U)$ are normalized by $h(0) = 1 = g(0)$, and $0 \notin (hg)(U)$. This paper treats the class T_{Ra} of logharmonic mappings $f(z) = zh(z)\overline{g(z)}$ satisfying $\varphi(z) = zh(z)g(z) \in HG$ is analytically typically real in U .

In Section 2, every mapping $f \in T_{Ra}$ is shown to admit an integral representation in terms of its second dilatation function and a function of positive real part with real coefficients. The radius of starlikeness is also obtained for the class T_{Ra} , as well as an upper estimate for its arclength.

For an analytic univalent function $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$, it is known [2] that f is typically real if and only if the image $f(U)$ is a domain symmetric with respect to the real axis. However, this characterization no longer holds for logharmonic maps, that is, it is not true that a univalent logharmonic mapping $F(z) = zh(z)\overline{g(z)} \in T_{Ra}$ if and only if the image $F(U)$ is a symmetric domain with respect to the real axis.



As an illustration, Figure 1 shows the mapping $F(z) = z(1+iz/3)(1+i\bar{z}/3) \in T_{Ra}$ but $F(U)$ is not a symmetric domain with respect to the real axis.

On the other hand, Figure 2 shows the mapping

$$F(z) = z \exp \{ \operatorname{Re} (4z/(1-z)) \} (1-\bar{z})/(1-z)$$

which does not belong to the class T_{Ra} , but yet maps U onto a symmetric domain with respect to the real axis $F(U)$.

In Section 3 we explore conditions on the dilatation a that would ensure a logharmonic mapping $f(z) = zh(z)\overline{g(z)} \in T_{Ra}$ necessarily satisfies $f(U)$ is symmetric with respect to the real axis. Sufficient conditions for univalent

FIGURE 1. Graph of $F(z) = z\left(1 + \frac{iz}{3}\right)\left(1 + \frac{i\bar{z}}{3}\right)$.FIGURE 2. Graph of $F(z) = z \frac{1-\bar{z}}{1-z} \exp \left\{ \operatorname{Re} \left(\frac{4z}{1-z} \right) \right\}$.

logharmonic mappings to belong to the class T_{Ra} in some subdisk of U are also determined.

2. An integral representation and radius of starlikeness

Let us denote by $\mathcal{P}_{\mathbb{R}}$ the class of normalized analytic functions with positive real part and with real coefficients in U . The following result gives a representation of $f \in T_{Ra}$ in terms of the dilatation a and $p \in \mathcal{P}_{\mathbb{R}}$.

Theorem 1. *Let $f = zh(z)\overline{g(z)}$ belongs to T_{Ra} with respect to $a \in B_0$. Then*

$$f(z) = \frac{zp(z)}{1-z^2} \exp \left(-2i \operatorname{Im} \int_0^z \frac{a(s)}{1+a(s)} \left(\frac{1+s^2}{s(1-s^2)} + \frac{p'(s)}{p(s)} \right) ds \right)$$

for some $p \in \mathcal{P}_{\mathbb{R}}$.

Proof. Let $\varphi(z) = zh(z)g(z)$, and

$$(2.1) \quad f(z) = \varphi(z) \frac{\overline{g(z)}}{g(z)}.$$

It follows from (1.1) that

$$\frac{g'(z)}{g(z)} = \frac{a(z)}{1+a(z)} \frac{\varphi'(z)}{\varphi(z)},$$

which readily yields

$$(2.2) \quad g(z) = \exp \int_0^z \frac{a(s)}{1+a(s)} \frac{\varphi'(s)}{\varphi(s)} ds.$$

Substituting (2.2) into (2.1) yields

$$f(z) = \varphi(z) \exp \left(-2i \operatorname{Im} \int_0^z \frac{a(s)}{1+a(s)} \frac{\varphi'(s)}{\varphi(s)} ds \right).$$

It is known [21] that every typically real analytic function φ has the form $(1-z^2)\varphi(z) = zp(z)$ for some $p \in \mathcal{P}_{\mathbb{R}}$, which yields the desired result. \square

In the next result, we obtain an estimate on the radius of starlikeness for the class T_{Ra} .

Theorem 2. *Let $f(z) = zh(z)\overline{g(z)} \in T_{Ra}$ with respect to $a \in B_0$. Then f maps the disk $|z| < 3 - 2\sqrt{2}$ onto a starlike domain.*

Proof. The function f maps the circle $|z| = r$ onto a starlike curve provided

$$\frac{\partial}{\partial \theta} \arg f(re^{i\theta}) = \operatorname{Im} \left(\frac{\partial}{\partial \theta} \log f(re^{i\theta}) \right) = \operatorname{Re} \frac{zf_z - \bar{z}f_{\bar{z}}}{f} > 0.$$

With $\varphi(z) = zh(z)g(z)$, a short computation gives

$$\operatorname{Re} \frac{zf_z - \bar{z}f_{\bar{z}}}{f} = \operatorname{Re} \left(\frac{1-a(z)}{1+a(z)} \frac{z\varphi'(z)}{\varphi(z)} \right)$$

for some $a \in B_0$.

Next let

$$q(z) = \frac{1-a(z)}{1+a(z)} \frac{z\varphi'(z)}{\varphi(z)},$$

and $\sigma(z) = \rho_0 z$. Kirwan [18] has shown that the radius of starlikeness for typically real analytic functions φ is $\rho_0 = \sqrt{2} - 1$. Thus $\operatorname{Re} \{ \zeta \varphi'(\zeta) / \varphi(\zeta) \}_{(\sigma(z))} > 0$, and so $q(\sigma(z))$ is subordinated to $((1+z)/(1-z))^2$ in U .

Writing $p(z) = (1+z)/(1-z)$, it follows from [14, p. 84] that

$$\left| p(z) - \frac{1+r^2}{1-r^2} \right| \leq \frac{2r}{1-r^2}.$$

Thus $|\arg(p(z))| < \pi/4$ provided $|z| < \rho_0$, where ρ_0 is a smallest positive root of the equation $r^2 - 2\sqrt{2}r + 1 = 0$. The function $f(z) = zh(z)\overline{g(z)}$ is thus starlike in the disk $|z| < \rho_0^2 = 3 - 2\sqrt{2}$. \square

The next result gives an upper estimate for arclength of all mappings f in the class T_{Ra} .

Theorem 3. *Let $f(z) = zh(z)\overline{g(z)} \in T_{Ra}$ with respect to $a \in B_0$. Then for $|z| = r$, an upper bound for arclength $L(r)$ is given by*

$$L(r) \leq \frac{2\pi r(1+r)^2}{(1-r)^4}.$$

Proof. Let C_r denote the image of the circle $|z| = r < 1$ under the mapping $w = f(z)$. Then

$$\begin{aligned} L(r) &= \int_{C_r} |df| = \int_0^{2\pi} |zf_z - \bar{z}f_{\bar{z}}| d\theta \\ &\leq M(r) \int_0^{2\pi} \left| \frac{zf_z - \bar{z}f_{\bar{z}}}{f} \right| d\theta, \end{aligned}$$

where $|f(z)| \leq M(r)$, $0 < r < 1$. Let $\varphi(z) = zh(z)g(z)$. Since φ is a typically real analytic function, and $|f| = |\varphi|$, then $|z\varphi'(z)/\varphi(z)| \leq (1+r)/(1-r)$ and $M(r) \leq r/(1-r)^2$.

Further

$$\begin{aligned} \left| \frac{zf_z - \bar{z}f_{\bar{z}}}{f} \right| &= \left| \frac{z\varphi'(z)}{\varphi(z)} - 2\operatorname{Re} \left(\frac{a(z)}{1+a(z)} \frac{z\varphi'(z)}{\varphi(z)} \right) \right| \\ &\leq \frac{1+r}{1-r} + 2\frac{r}{1-r} \frac{1+r}{1-r}, \end{aligned}$$

and thus,

$$L(r) \leq \frac{2\pi r(1+r)^2}{(1-r)^4}. \quad \square$$

3. Logharmonic mappings in the class T_{Ra}

The following result is readily established, and thus the proof is omitted. It describes the geometry of a logharmonic function in the class T_{Ra} when its second dilatation has real coefficients.

Theorem 4. *Let $f(z) = zh(z)\overline{g(z)} \in T_{Ra}$ be a sense-preserving logharmonic mapping in U . If the second dilatation function a has real coefficients, that is, $a(\bar{z}) = \overline{a(z)}$, then $f(U)$ is symmetric with respect to the real axis.*

The final result derives sufficient conditions for $f \in T_{Ra}$ in some subdisk of U .

Theorem 5. *Let $f(z) = zh(z)\overline{g(z)}$ be a univalent sense-preserving logharmonic mapping in U normalized by $h(0) = 1 = g(0)$, where its second dilatation function a has real coefficients. Further, suppose $f(U) = \Omega$, $\Omega \neq \mathbb{C}$, is a strictly starlike Jordan domain. If $f(U)$ is a symmetric domain with respect to the real axis, and $|a(z)| \leq k < 1$ in U , then $\varphi(z) = zh(z)g(z)$ is typically real in the disk $|z| < \sqrt{2} - 1$.*

Proof. The domain Ω is a strictly starlike Jordan domain provided each radial ray from 0 intersects the boundary $\partial\Omega$ of $\Omega = f(U)$ at exactly one point of \mathbb{C} . Further, since $|a(z)| \leq k < 1$ in U , it follows from [7, Lemma 2.4] that there is only one univalent logharmonic mapping from U onto Ω which is a solution of (1.1) normalized by $f(0) = 0$ and $h(0) = 1 = g(0)$.

Since a has real coefficients, then $\overline{a(\bar{z})} = a(z)$. On the other hand, the mapping $F(z) = \overline{f(\bar{z})}$ is also univalent and logharmonic in U , where $F(U) = f(U)$.

If $F(z) = zH(z)\overline{G(z)} = z\overline{h(\bar{z})}g(\bar{z})$ with $H(z) = \overline{h(\bar{z})}$ and $G(z) = \overline{g(\bar{z})}$, then F satisfies the normalization $F(0) = 0, H(0) = 1 = G(0)$, and F is a solution of

$$\frac{\overline{F_z(z)}}{F(z)} = \overline{a(\bar{z})} \frac{F_z(z)}{F(z)} = a(z) \frac{F_z(z)}{F(z)}.$$

Thus, F is a logharmonic mapping with respect to the same a , and consequently, $f(z) \equiv F(z)$ in U . This implies f has real coefficients, and so $\psi(z) = zh(z)/g(z) = f(z)/|g(z)|^2$ has real coefficients.

Direct calculations yield

$$\frac{g'(z)}{g(z)} = \frac{a(z)}{1 - a(z)} \frac{\psi'(z)}{\psi(z)},$$

which upon integrating leads to

$$g(z) = \exp \int_0^z \frac{a(t) - \psi'(t)}{1 - a(t) \psi(t)} dt.$$

Then g , and so does h , have real coefficients, and thus $\varphi(z) = zh(z)g(z)$ has real coefficients. Furthermore, [5, Theorem 3.1] shows that φ is starlike univalent in the disk $|z| < \rho$, where $\rho = \sqrt{2} - 1$. Thus φ is typically real in the disk $|z| < \sqrt{2} - 1$. \square

Acknowledgment. The authors are indebted to the referee for the insightful suggestions that helped improve the clarity and conciseness of this manuscript. The third author gratefully acknowledged support from Universiti Sains Malaysia research grant 1001/PMATHS/811280.

References

- [1] Z. Abdulhadi, *Close-to-starlike logharmonic mappings*, Internat. J. Math. Math. Sci. **19** (1996), no. 3, 563–574.
- [2] ———, *Typically real logharmonic mappings*, Int. J. Math. Math. Sci. **31** (2002), no. 1, 1–9.
- [3] Z. AbdulHadi, N. M. Alareefi, and R. M. Ali, *On the convex-exponent product of logharmonic mappings*, J. Inequal. Appl. **2014** (2014), 485, 10 pp.
- [4] Z. Abdulhadi and R. M. Ali, *Univalent logharmonic mappings in the plane*, Abstr. Appl. Anal. **2012** (2012), Art. ID 721943, 32 pp.
- [5] ———, *On rotationally starlike logharmonic mappings*, Math. Nachr. **288** (2015), no. 7, 723–729.

- [6] Z. Abdulhadi and D. Bshouty, *Univalent functions in $H \cdot \overline{H}(D)$* , Trans. Amer. Math. Soc. **305** (1988), no. 2, 841–849.
- [7] Z. Abdulhadi and W. Hengartner, *Univalent harmonic mappings on the left half-plane with periodic dilatations*, in Univalent functions, fractional calculus, and their applications (Kōriyama, 1988), 13–28, Ellis Horwood Ser. Math. Appl, Horwood, Chichester, 1989.
- [8] Z. Abdulhadi and Y. Abu Muhanna, *Starlike log-harmonic mappings of order α* , JIPAM. J. Inequal. Pure Appl. Math. **7** (2006), no. 4, Article 123, 6 pp.
- [9] Y. Abu-Muhanna and A. Lyzzaik, *The boundary behaviour of harmonic univalent maps*, Pacific J. Math. **141** (1990), no. 1, 1–20.
- [10] J. Clunie and T. Sheil-Small, *Harmonic univalent functions*, Ann. Acad. Sci. Fenn. Ser. A I Math. **9** (1984), 3–25.
- [11] P. L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften, **259**, Springer-Verlag, New York, 1983.
- [12] P. Duren and G. Schober, *A variational method for harmonic mappings onto convex regions*, Complex Variables Theory Appl. **9** (1987), no. 2-3, 153–168.
- [13] ———, *Linear extremal problems for harmonic mappings of the disk*, Proc. Amer. Math. Soc. **106** (1989), no. 4, 967–973.
- [14] A. W. Goodman, *Univalent Functions. Vol. I*, Mariner Publishing Co., Inc., Tampa, FL, 1983.
- [15] W. Hengartner and G. Schober, *On the boundary behavior of orientation-preserving harmonic mappings*, Complex Variables Theory Appl. **5** (1986), no. 2-4, 197–208.
- [16] ———, *Harmonic mappings with given dilatation*, J. London Math. Soc. (2) **33** (1986), no. 3, 473–483.
- [17] S. H. Jun, *Univalent harmonic mappings on $\Delta = \{z : |z| > 1\}$* , Proc. Amer. Math. Soc. **119** (1993), no. 1, 109–114.
- [18] W. E. Kirwan, *Extremal problems for the typically real functions*, Amer. J. Math. **88** (1966), 942–954.
- [19] J. C. C. Nitsche, *Lectures on Minimal Surfaces. Vol. 1*, translated from the German by Jerry M. Feinberg, Cambridge University Press, Cambridge, 1989.
- [20] R. Osserman, *A Survey of Minimal Surfaces*, second edition, Dover Publications, Inc., New York, 1986.
- [21] W. Rogosinski, *Über positive harmonische Entwicklungen und typisch-reelle Potenzreihen*, Math. Z. **35** (1932), no. 1, 93–121.

ZAYID ABDULHADI
 DEPARTMENT OF MATHEMATICS
 AMERICAN UNIVERSITY OF SHARJAH
 SHARJAH, BOX 26666, UAE
 Email address: zahadi@aus.edu

NAJLA M. ALARIFI
 DEPARTMENT OF MATHEMATICS
 IMAM ABDULRAHMAN BIN FAISAL UNIVERSITY
 DAMMAM 31113, KINGDOM OF SAUDI ARABIA
 Email address: najarifi@hotmail.com

ROSIHAN M. ALI
 SCHOOL OF MATHEMATICAL SCIENCES
 UNIVERSITI SAINS MALAYSIA
 11800 USM, PENANG, MALAYSIA
 Email address: rosihan@usm.my