

## SR-ADDITIVE CODES

SAADOUN MAHMOUDI AND KARIM SAMEI

ABSTRACT. In this paper, we introduce  $SR$ -additive codes as a generalization of the classes of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$  and  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes, where  $S$  is an  $R$ -algebra and an  $SR$ -additive code is an  $R$ -submodule of  $S^\alpha \times R^\beta$ . In particular, the definitions of bilinear forms, weight functions and Gray maps on the classes of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$  and  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes are generalized to  $SR$ -additive codes. Also the singleton bound for  $SR$ -additive codes and some results on one weight  $SR$ -additive codes are given. Among other important results, we obtain the structure of  $SR$ -additive cyclic codes. As some results of the theory, the structure of cyclic  $\mathbb{Z}_2\mathbb{Z}_4$ ,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ ,  $\mathbb{Z}_2\mathbb{Z}_2[u]$ ,  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$  and  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$ -additive codes are presented.

### 1. Introduction

An important class of additive codes is  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes. A subgroup of  $\mathbb{Z}_2^\alpha \times \mathbb{Z}_4^\beta$ , where  $\alpha$  and  $\beta$  are positive integers, is called a  $\mathbb{Z}_2\mathbb{Z}_4$ -additive code. A comprehensive study on  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes has been introduced in [9] by Borges et al. The studies on  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes and their algebraic structures have attracted many researchers; see [2, 6–9, 13, 15–17].

$\mathbb{Z}_2\mathbb{Z}_4$ -additive codes were generalized to  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes [4, 21]. Also  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes is another generalization of  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes which has been introduced by Aydogdu et al. [3].

Recently, Aydogdu and Siap generalized  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes and  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes to  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes [5]. Also,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes have been studied in [10]. Also additive codes were studied over direct product of chain rings in [11].

Note that in  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes and  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes,  $\mathbb{Z}_2$  is a  $\mathbb{Z}_4$ -algebra and  $\mathbb{Z}_{2^s}$ -algebra; respectively. Also in  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes,  $\mathbb{Z}_2$  is considered as a  $\mathbb{Z}_2[u]$ -algebra and  $\mathbb{Z}_{p^r}$  is a  $\mathbb{Z}_{p^s}$ -algebra in  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes. Also in additive codes over product of two chain rings, one of the rings is an algebra over another ring.

---

Received October 18, 2018; Accepted December 18, 2018.

2010 *Mathematics Subject Classification.* 94B15.

*Key words and phrases.* additive code, chain ring, Galois ring.

In this paper, we generalize above codes to  $SR$ -additive codes, where  $S$  is an  $R$ -algebra. In this generalization, a subset  $C$  of  $S^\alpha \times R^\beta$  is called an  $SR$ -additive code if  $C$  is an  $R$ -submodule of  $S^\alpha \times R^\beta$ . We present the structure of  $SR$ -additive cyclic codes. Also we give the structure of  $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes,  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2)$ -additive cyclic codes and cyclic codes over direct product of chain rings as results of this theory, which the structure of these codes are the main parts of [2], [10], [22] and [11]; respectively.

Also, we obtain the structure of  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$  and  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$ -additive cyclic codes as other results of this theory.

In Section 4, we define an inner product over  $SR$ -additive codes which is a generalization of the inner products over  $\mathbb{Z}_2\mathbb{Z}_4$ ,  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ ,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ ,  $\mathbb{Z}_2\mathbb{Z}_2[u]$  additive codes. We show that the dual code of any  $SR$ -additive cyclic code is also an  $SR$ -additive cyclic code.

In Section 5, we find the Singleton bound for  $SR$ -additive codes. As examples, the Singleton bound for  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes and  $\mathbb{Z}_2[u]\mathbb{Z}_2$ -additive codes are given. In Section 6, we investigate one weight  $SR$ -additive codes. In particular, one weight  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes are determined.

Throughout this paper  $R$  and  $S$  are finite commutative rings such that  $S$  is an  $R$ -algebra.

## 2. Preliminaries

In this section, we remind some facts of  $R$ -additive codes which are applied throughout this paper. Also the structure of cyclic codes over some rings are given.

**Definition 2.1.** Let  $S$  be an  $R$ -algebra with a ring homomorphism  $f : R \rightarrow S$ . A nonempty subset  $C$  of  $S^n$  is called  $R$ -additive code if  $C$  is an  $R$ -submodule of  $S^n$ , where the scalar multiplication is defined as follows: for  $r \in R$  and  $(a_0, a_1, \dots, a_{n-1}) \in C$ , we have

$$r \cdot (a_0, a_1, \dots, a_{n-1}) = (f(r)a_0, f(r)a_1, \dots, f(r)a_{n-1}).$$

**Example 2.2** (Linear codes). Let  $R$  be a commutative ring with identity. A subset  $C$  of  $R^n$  is called a linear code if  $C$  is an  $R$ -submodule of  $R^n$ . Now consider  $R$  as  $R$ -algebra with identity homomorphism. Clearly, the subset  $C$  of  $R^n$  is a linear code if and only if  $C$  is an  $R$ -additive code.

Above example shows that  $R$ -additive codes is a generalization of linear codes. The following example give some special cases which  $R$ -additive codes and linear codes are the same.

**Example 2.3.** (1) Let  $f : R \rightarrow S$  be a ring isomorphism. In this case,  $R$ -additive codes over  $S$  are exactly linear codes over  $S$ .

(2) Let  $S = R/I$ , where  $I$  is an ideal of  $R$  and  $f : R \rightarrow R/I$  is the natural homomorphism. For any nonempty subset  $C$  of  $S^n$ , we have  $I.C = 0$ . Hence

$R$ -additive codes over  $S$  are exactly linear codes over  $S$ . Moreover, if  $f : R \rightarrow S$  is a surjective ring homomorphism then  $R$ -additive codes over  $S$  are exactly linear codes.

**Example 2.4** (Additive codes). Let  $S$  be a local ring of characteristic  $p^r$ . A subset  $C$  of  $S^n$  is called an additive code if  $C$  is a subgroup of  $S^n$  under addition. But we have the injective ring homomorphism  $f : \mathbb{Z}_{p^r} \rightarrow S, x \mapsto x.1_S$ . It is easy to see that additive codes are exactly  $\mathbb{Z}_{p^r}$ -submodules of  $S^n$ . In other words, additive codes over  $S$  are exactly  $\mathbb{Z}_{p^r}$ -additive codes over  $S$ .

**Example 2.5** ( $\mathbb{F}_q$ -linear codes over  $\mathbb{F}_{q^t}$ ). A subset  $C$  of  $(\mathbb{F}_{q^t})^n$  is called an  $\mathbb{F}_q$ -linear code over  $\mathbb{F}_{q^t}$  of length  $n$ , if  $C$  is an  $\mathbb{F}_q$ -submodule of  $(\mathbb{F}_{q^t})^n$ . Clearly these codes are  $R$ -additive codes, where  $R = \mathbb{F}_q$  and  $S = \mathbb{F}_{q^t}$ .

For a positive integer  $n$ , let  $R_n = R[x]/\langle x^n - 1 \rangle$  and  $S_n = S[x]/\langle x^n - 1 \rangle$ . Consider the following correspondence map.

$$(1) \quad \begin{aligned} \pi : S^n &\longrightarrow S_n, \\ (a_0, a_1, \dots, a_{n-1}) &\longmapsto a_0 + a_1x + \dots + a_{n-1}x^{n-1} + \langle x^n - 1 \rangle. \end{aligned}$$

Clearly  $\pi$  is an  $R$ -module isomorphism. We will identify  $S^n$  with  $S_n$  under  $\pi$  and for simplicity, we write the polynomial  $a_0 + a_1x + \dots + a_{n-1}x^{n-1}$  for the residue class  $a_0 + a_1x + \dots + a_{n-1}x^{n-1} + \langle x^n - 1 \rangle$ . The following proposition gives the structure of cyclic  $R$ -additive codes.

**Proposition 2.6** ([19, Proposition 3.1]). *Let  $\pi$  be the correspondence map defined in (1). Then a nonempty subset  $C$  of  $S^n$  is a cyclic  $R$ -additive code if and only if  $\pi(C)$  is an  $R_n$ -submodule of  $S_n$ .*

Let  $\omega$  be a weight function over  $S$ . If  $A_S = \text{Max}\{\omega(x) : x \in S\}$ , then we have the following bound for minimum weight of  $R$ -additive codes.

**Theorem 2.7** ([20, Theorem 3.5]). *Let  $R$  be a finite chain ring and  $S$  be a free  $R$ -algebra of  $\dim_R(S) = m$ . If there exists a nondegenerate bilinear form  $\beta : S \times S \rightarrow R$ , then  $\lfloor \frac{d_\omega(C)-1}{A_S} \rfloor \leq n - \lceil \frac{\text{rank}(C)}{m} \rceil$ .*

Now we remind the structure of cyclic codes over a chain ring  $R$  of length  $n$  coprime to  $\text{Char}(R)$ . Also the structure of cyclic codes over  $\mathbb{Z}_2 + u\mathbb{Z}_2$ ,  $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$  and  $\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2$  for an arbitrary length are given.

**Theorem 2.8.** *Let  $R$  be a chain ring with the maximal ideal  $\mathfrak{m} = \langle \gamma \rangle$  of nilpotency index  $s$  and  $C$  be a cyclic code of length  $n$  over  $R$ , where  $(n, \text{Char}(R)) = 1$ . Then*

- (1) *There is a unique set of pairwise co-prime monic polynomials  $g_0, \dots, g_s$  over  $R$  (possibly, some of them are equal to 1) such that  $g_0g_1 \cdots g_s = x^n - 1$  in  $R[x]$  and  $C = \langle \hat{g}_1, \gamma\hat{g}_2, \dots, \gamma^{s-1}\hat{g}_s \rangle$ , where  $\hat{g}_i = \prod_{j \neq i} g_j$ . Moreover,  $|C| = |R/\mathfrak{m}|^{\sum_{i=0}^{s-1} (s-i) \deg g_{i+1}}$ .*
- (2) *If  $h_i = g_0g_{i+2} \cdots g_s$  for  $i = 0, 1, \dots, s-2$  and  $h_{s-1} = g_0$ . Then  $h_{s-1}|h_{s-2}| \cdots |h_0|(x^n - 1)$ , and  $C = \langle h_0 + \gamma h_1 + \dots + \gamma^{s-1}h_{s-1} \rangle$ .*

*Proof.* Part (1) follows from Theorem 3.4 in [12]. We have part (2) by Theorem 3.5 in [12] and Theorem 2.4 in [11].  $\square$

The following corollary is a result of Proposition 2.8.

**Corollary 2.9.** *Let  $C$  be a cyclic code of length  $n$  over  $R = \mathbb{Z}_{p^s}$ , where  $(n, p) = 1$ . Then there exists a set of polynomials  $h_0, h_1, \dots, h_{s-1}$  in  $R[x]$  such that  $h_0 | (x^n - 1)$ ,  $h_i | h_{i-1}$  for  $i = 1, \dots, s-1$  and  $C = \langle h_0 + ph_1 + \dots + p^{s-1}h_{s-1} \rangle$ . Moreover if  $\widehat{h}_i = \frac{h_{i-1}}{h_i}$  for  $i \geq 1$  and  $\widehat{h}_0 = \frac{x^n - 1}{h_0}$ , then  $|C| = p^d$ , where  $d = \sum_{i=0}^{s-1} (s-i) \deg \widehat{h}_i$ . In special case, if  $n$  is odd and  $C$  is a cyclic code of length  $n$  over  $R = \mathbb{Z}_4$ , then  $C = \langle g(x) + 2a(x) \rangle$ , where  $a(x) | g(x) | (x^n - 1)$  in  $\mathbb{Z}_4[x]$ . In this case,  $|C| = 2^{2t_1 + t_2}$ , where  $t_1 = \deg \frac{x^n - 1}{g(x)}$  and  $t_2 = \deg \frac{g(x)}{a(x)}$ .*

**Theorem 2.10** ([1, Theorem 1]). *Let  $C$  be a cyclic code over  $\mathbb{Z}_2 + u\mathbb{Z}_2$  of length  $n$ . Then*

- (1) *If  $n$  is odd, then  $(\mathbb{Z}_2 + u\mathbb{Z}_2)_n$  is principal ideal ring and  $C = \langle g(x) + ua(x) \rangle$ , where  $g(x)$  and  $a(x)$  are polynomials in  $\mathbb{Z}_2[x]$  such that  $a(x) | g(x) | (x^n - 1) \pmod{2}$ .*
- (2) *If  $n$  is not odd, then*
  - (a)  *$C = \langle g(x) + up(x) \rangle$  such that  $g(x) | (x^n - 1) \pmod{2}$ ,  $(g(x) + up(x)) | (x^n - 1)$  in  $\mathbb{Z}_2 + u\mathbb{Z}_2$  and  $g(x) | p(x) \left( \frac{x^n - 1}{g(x)} \right)$ . Or*
  - (b)  *$C = \langle g(x) + up(x), ua(x) \rangle$  such that  $g(x)$ ,  $a(x)$  and  $p(x)$  are polynomials in  $\mathbb{Z}_2[x]$ . And  $a(x) | g(x) | (x^n - 1) \pmod{2}$ ,  $a(x) | p(x) \left( \frac{x^n - 1}{g(x)} \right)$  and  $\deg a(x) > \deg p(x)$ .*

**Theorem 2.11** ([1, Theorem 2]). *Let  $C$  be a cyclic code over  $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$  of length  $n$ , then*

- (1) *If  $n$  is odd, then  $(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)_n$  is principal ideal ring.  $C = \langle g(x) + ua_1(x) + u^2a_2(x) \rangle$ , where  $a_1(x)$ ,  $a_2(x)$  and  $g(x)$  are polynomials in  $\mathbb{Z}_2[x]$  such that  $a_2(x) | a_1(x) | g(x) | (x^n - 1) \pmod{2}$ .*
- (2) *If  $n$  is not odd, then*
  - (a)  *$C = \langle g + up_1 + u^2p_2 \rangle$ , where  $p_2 | p_1 | g | (x^n - 1) \pmod{2}$ ,  $(g + up_1) | (x^n - 1)$  in  $\mathbb{Z}_2 + u\mathbb{Z}_2$  and  $(g + up_1 + u^2p_2) | (x^n - 1)$  in  $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$  and  $\deg p_2 < \deg p_1$ .*
  - (b)  *$C = \langle g + up_1 + u^2p_2, u^2a_2 \rangle$ , where  $a_2 | g | (x^n - 1) \pmod{2}$ ,  $(g + up_1) | (x^n - 1)$  in  $\mathbb{Z}_2 + u\mathbb{Z}_2$ ,  $g(x) | p_1 \left( \frac{x^n - 1}{g(x)} \right)$  and  $a_2$  divides  $p_1 \left( \frac{x^n - 1}{g(x)} \right)$  and  $p_2 \left( \frac{x^n - 1}{g(x)} \right) \left( \frac{x^n - 1}{g(x)} \right)$  and  $\deg p_2 < \deg a_2$ . Or*
  - (c)  *$C = \langle g + up_1 + u^2p_2, ua_1 + u^2q_1, u^2a_2 \rangle$ , where  $a_2 | a_1 | g | (x^n - 1) \pmod{2}$ ,  $a_1 | p_1 \left( \frac{x^n - 1}{g(x)} \right)$  and  $a_2$  divides  $q_1 \left( \frac{x^n - 1}{a_1(x)} \right)$  and  $p_2 \left( \frac{x^n - 1}{g(x)} \right) \left( \frac{x^n - 1}{a_1(x)} \right)$ . Moreover,  $\deg p_2 < \deg a_2$ ,  $\deg q_1 < \deg a_2$  and  $\deg p_1 < \deg a_1$ .*

The following theorem gives the structure of cyclic codes over the non Frobenius ring  $\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2 = \{0, 1, u, v, 1 + u, 1 + v, u + v, 1 + u + v\}$ .

**Theorem 2.12.** *Let  $C$  be a cyclic code over  $R = \mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2$  of length  $n$ . Then  $C$  has a unique representation as follows:*

$$C = \langle g + up_1 + vp_2, ua_1 + vq_1, va_2 \rangle,$$

where

- (1)  $a_2|a_1|g|(x^n - 1)$  and  $a_1|p_1(\frac{x^n-1}{g})$ ,
- (2)  $a_2|q_1(\frac{x^n-1}{a_1})$  and  $a_2|p_2(\frac{x^n-1}{g})(\frac{x^n-1}{a_1})$ ,
- (3)  $\deg p_2, \deg q_1 < \deg a_2$ .

Moreover if  $n$  is odd, then  $C = \langle g + ua_1, va_2 \rangle$ , where  $a_2|a_1|g|(x^n - 1)$ .

*Proof.* See Theorems 1 and 2, Lemmas 3 and 4 and Corollary 1 in [18].  $\square$

### 3. SR-additive cyclic codes

The structure of  $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes investigated in [2]. As generalizations of these codes, recently  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$  and  $\mathbb{Z}_2\mathbb{Z}_2[u]$  additive codes have been introduced in [3] and [5]. Also the generator polynomials of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes were given in [10]. Moreover, additive codes studied over direct product of chain rings with the same residue fields in [11]. In this section, we define and extend these codes to SR-additive codes, where  $R$  is a finite commutative ring and  $S$  is a finite commutative  $R$ -algebra. A theory to find the generators of SR-additive cyclic codes is given. As results, we obtain the generators of  $\mathbb{Z}_2\mathbb{Z}_4$ ,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ ,  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive cyclic codes. Also the results in [11] on the structure of cyclic codes over direct product of chain rings with the same residue fields are given as a result of the theory. Moreover the structure of  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2)$ ,  $(\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$  and  $(\mathbb{Z}_2 + u\mathbb{Z}_2)(\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2)$ -additive cyclic codes as new examples of SR-additive cyclic codes are given, which we can not obtain their structures by previous works.

**Definition 3.1.** Let  $\alpha$  and  $\beta$  be two positive integers. A nonempty subset  $C$  of  $S^\alpha \times R^\beta$  is called an SR-additive code if  $C$  is an  $R$ -submodule with the following scalar multiplication: for  $r \in R$  and  $(s_\alpha, r_\beta) = (s_0, s_1, \dots, s_{\alpha-1}, r_0, r_1, \dots, r_{\beta-1}) \in C$ ,

$$r \cdot (s_\alpha, r_\beta) = (f(r)s_\alpha, rr_\beta) = (f(r)s_0, f(r)s_1, \dots, f(r)s_{\alpha-1}, rr_0, rr_1, \dots, rr_{\beta-1}).$$

We say that an SR-additive code  $C$  is cyclic if  $(s_{\alpha-1}, s_0, \dots, s_{\alpha-2}, r_{\beta-1}, r_0, \dots, r_{\beta-2}) \in C$  whenever  $(s_0, s_1, \dots, s_{\alpha-1}, r_0, r_1, \dots, r_{\beta-1}) \in C$ .

Consider the map  $\pi' : S^\alpha \times R^\beta \rightarrow S_\alpha \times R_\beta, (s_0, s_1, \dots, s_{\alpha-1}, r_0, r_1, \dots, r_{\beta-1}) \mapsto (s_0 + s_1x + \dots + s_{\alpha-1}x^{\alpha-1} + \langle x^\alpha - 1 \rangle, r_0 + r_1x + \dots + r_{\beta-1}x^{\beta-1} + \langle x^\beta - 1 \rangle)$ . Clearly  $\pi'$  is an  $R$ -module isomorphism. We will identify  $S^\alpha \times R^\beta$  with  $S_\alpha \times R_\beta$  under  $\pi'$  and for simplicity we write  $(s_0 + s_1x + \dots + s_{\alpha-1}x^{\alpha-1}, r_0 + r_1x + \dots + r_{\beta-1}x^{\beta-1})$  for above residue class.

**Lemma 3.2.** *A subset  $C$  of  $S^\alpha \times R^\beta$  is an SR-additive cyclic code if and only if  $\pi'(C)$  is an  $R[x]$ -submodule of  $S_\alpha \times R_\beta$ .*

*Proof.* Clearly  $S_\alpha \times R_\beta$  is an  $R[x]$ -module. Since  $\pi'$  is an  $R$ -module isomorphism,  $C$  is an  $R$ -submodule if and only if  $\pi'(C)$  is an  $R$ -submodule. Now for an element  $(s_\alpha, r_\beta) = (s_0, s_1, \dots, s_{\alpha-1}, r_0, r_1, \dots, r_{\beta-1}) \in C$ , the cyclic shift  $\sigma(s_\alpha, r_\beta) = (s_{\alpha-1}, s_0, \dots, s_{\alpha-2}, r_{\beta-1}, r_0, \dots, r_{\beta-2}) \in C$  if and only if  $x\pi'(s_\alpha, r_\beta) = \pi'(\sigma(s_\alpha, r_\beta)) \in \pi'(C)$ . This completes the proof.  $\square$

We identify  $C$  with  $\pi'(C)$ . Now we find the generator polynomials of  $C$ .

**Theorem 3.3.** *A subset  $C$  of  $S_\alpha \times R_\beta$  is an  $SR$ -additive cyclic code if and only if  $C = \langle (g_1, 0), \dots, (g_s, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle_{R[x]}$  such that*

- (1)  $C_1 = \langle f_1, \dots, f_r \rangle_{R[x]}$  is a cyclic linear code over  $R$  of length  $\beta$ ,
- (2)  $C_2 = \langle g_1, \dots, g_s \rangle_{R[x]}$  is a cyclic  $R$ -additive code over  $S$  of length  $\alpha$ ,
- (3)  $h_1, \dots, h_r$  are elements of  $S_\alpha$ ,
- (4)  $|C| = |C_1||C_2|$ .

*Proof.* Let  $C \subseteq S_\alpha \times R_\beta$  be an  $SR$ -additive cyclic code. Clearly the projection map  $\phi : C \rightarrow R_\beta$  is an  $R[x]$ -homomorphism. Hence  $Im(\phi)$  is an  $R[x]$ -submodule of  $R_\beta$ . As  $\langle x^\beta - 1 \rangle \cdot Im(\phi) \subseteq \langle x^\beta - 1 \rangle \cdot R_\beta = 0$ ,  $Im(\phi)$  is an ideal of  $R_\beta$ . In other words  $Im(\phi)$  is a linear cyclic code over  $R$  of length  $\beta$ , say  $C_1$ . Let  $C_1 = \langle f_1, \dots, f_r \rangle_{R[x]} = \langle \phi(h_1, f_1), \dots, \phi(h_r, f_r) \rangle_{R[x]}$ . Now,  $\ker \phi$  is an  $R[x]$ -submodule of  $C$ . Let  $C_2 = \{g \in S_\alpha : (g, 0) \in \ker \phi\}$ , then clearly  $C_2$  is an  $R[x]$ -submodule of  $S_\alpha$ . Since  $\langle x^\alpha - 1 \rangle \cdot C_2 \subseteq \langle x^\alpha - 1 \rangle \cdot S_\alpha = 0$ ,  $C_2$  is an  $R_\alpha$ -module. In other words  $C_2$  is a cyclic  $R$ -additive code of length  $\alpha$  over  $S$ . If  $C_2 = \langle g_1, \dots, g_s \rangle_{R_\alpha}$ , then  $\ker \phi = \langle (g_1, 0), \dots, (g_s, 0) \rangle_{R[x]}$ . Therefore  $C = \langle (g_1, 0), \dots, (g_s, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle_{R[x]}$ . Since  $\phi$  is an  $R[x]$ -homomorphism,  $\frac{C}{\ker \phi} \cong C_1$ , hence  $|C| = |\ker \phi||C_1| = |C_2||C_1|$ .  $\square$

**Proposition 3.4.** *With the above assumptions, let  $f : R \rightarrow S$  be a surjective ring homomorphism and  $C = \langle (g_1, 0), \dots, (g_s, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle_{R[x]}$  be an  $SR$ -additive cyclic code. Also let  $\{g_{i_1}, \dots, g_{i_t}\}$  be a subset of  $\{g_1, \dots, g_s\}$  such that  $g_{i_j}$  is monic for all  $j; j = 1, \dots, t$ . Then we can assume that  $\deg h_i < \min\{\deg g_{i_j} : 1 \leq j \leq t\}$  for all  $i; 1 \leq i \leq r$ .*

*Proof.* Since  $f$  is surjective, every  $R$ -additive code over  $S$  is linear. In particular,  $C_2$  is a cyclic linear code over  $S$ . Let  $g_j$  be monic and  $\deg h_i \geq \deg g_j$  for some  $i$ . Let  $\deg h_i - \deg g_j = \ell$  and  $a \in S$  be the leading coefficient of  $h_i$ . Then  $(h_i, f_i) = (h_i - ax^\ell g_j, f_i) + ax^\ell (g_j, 0)$ . Thus  $\langle (h_i, f_i), (g_j, 0) \rangle = \langle (h_i - ax^\ell g_j, f_i), (g_j, 0) \rangle$ . Hence we can use  $h_i - ax^\ell g_j$  instead of  $h_i$ . By this method we can reduce  $\deg h_i$ .  $\square$

**Proposition 3.5.** *Let  $C = \langle (g_1, 0), \dots, (g_s, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle_{R[x]}$  be an  $SR$ -additive cyclic code as in Theorem 3.3. Then*

$$(x^\beta - 1)h_i \in C_2 = \langle g_1, \dots, g_s \rangle_{R[x]}.$$

*Proof.* Clearly  $(x^\beta - 1)(h_i, f_i) = ((x^\beta - 1)h_i, 0) \in \ker \phi$ . Hence  $(x^\beta - 1)h_i \in C_2 = \langle g_1, \dots, g_s \rangle_{R[x]}$ .  $\square$

**Corollary 3.6** ( $(R/\mathfrak{m})R$ -additive cyclic codes). *Let  $R$  be a finite local ring with the unique maximal ideal  $\mathfrak{m}$  and  $C \subseteq (R/\mathfrak{m})^\alpha \times R^\beta$  be an  $(R/\mathfrak{m})R$ -additive cyclic code. Then  $C = \langle (g, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle$  with the following conditions:*

- (a)  $g|x^\alpha - 1$  over  $(R/\mathfrak{m})$ ,
- (b)  $h_i \in (R/\mathfrak{m})_\alpha$ ,
- (c)  $C_1 = \langle f_1, \dots, f_r \rangle$  is a linear cyclic code over  $R$ .

*Proof.*  $R/\mathfrak{m}$  is an  $R$ -algebra with the natural ring homomorphism  $f : R \rightarrow R/\mathfrak{m}$ . Since  $f$  is surjective,  $R$ -additive codes over  $R/\mathfrak{m}$  are linear over  $R/\mathfrak{m}$ . Now, we have the results by Theorem 3.3.  $\square$

$\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes is an example of  $(R/\mathfrak{m})R$ -additive cyclic codes. This class of codes is discussed in [2]. We obtain the structure of these codes as a result of above discussion.

**Corollary 3.7** ( $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes). *Let  $C \subseteq (\mathbb{Z}_2)_\alpha \times (\mathbb{Z}_4)_\beta$  be a  $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic code. If  $\beta$  is an odd integer, then*

- (1)  $C = \langle (h(x), 0), (\ell(x), g(x) + 2a(x)) \rangle$ , where
  - (a)  $h(x)$  is a monic polynomial over  $\mathbb{Z}_2$  such that  $h(x)|(x^\alpha - 1)$ ,
  - (b)  $a(x)|g(x)|(x^\beta - 1)$  in  $\mathbb{Z}_4[x]$ ,
  - (c)  $\ell(x) \in (\mathbb{Z}_2)_\alpha$  and  $\deg \ell(x) < \deg h(x)$ .
- (2) If  $t_1 = \deg \frac{x^\beta - 1}{g(x)}$ ,  $t_2 = \deg \frac{g(x)}{a(x)}$  and  $t = \deg h(x)$ , then  $|C| = 2^{2t_1 + t_2 + \alpha - t}$ .

*Proof.* By above corollary,  $C = \langle (h(x), 0), (\ell_1, f_1), \dots, (\ell_r, f_r) \rangle$ , where  $h(x)$  is a monic polynomial over  $\mathbb{Z}_2$  such that  $h(x)|(x^\alpha - 1)$ . Also  $C_1 = \langle f_1, \dots, f_r \rangle$  is a linear cyclic code over  $\mathbb{Z}_4$ . By Corollary 2.9, there exist polynomials  $g(x)$  and  $a(x)$  over  $\mathbb{Z}_4$  such that  $C_1 = \langle g(x) + 2a(x) \rangle$ , where  $a(x)|g(x)|(x^\beta - 1)$  in  $\mathbb{Z}_4[x]$ . Hence  $C = \langle (h(x), 0), (\ell(x), g(x) + 2a(x)) \rangle$ , where  $\ell(x) \in (\mathbb{Z}_2)_\alpha$  and  $\deg \ell(x) < \deg h(x)$ . By Corollary 2.9,  $|C_1| = 2^{2t_1 + t_2}$ , where  $t_1 = \deg \frac{x^\beta - 1}{g(x)}$  and  $t_2 = \deg \frac{g(x)}{a(x)}$ . Also  $|C_2| = |\langle h(x) \rangle| = 2^{\alpha - t}$ , where  $t = \deg h(x)$ . Therefore by Theorem 3.3,  $|C| = |C_1||C_2| = 2^{2t_1 + t_2 + \alpha - t}$ .  $\square$

Another example of  $SR$ -additive codes is the class of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes (see [10]). We give the structure of these codes as another result of above discussion.

**Corollary 3.8** ( $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes). *Let  $1 \leq r < s$  and  $C \subseteq (\mathbb{Z}_{p^r})_\alpha \times (\mathbb{Z}_{p^s})_\beta$  be a  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic code. If  $(p, \beta) = 1$  and  $(p, \alpha) = 1$ , then*

- (1)  $C = \langle (h'_0 + ph'_1 + \dots + p^{r-1}h'_{r-1}, 0), (\ell(x), h_0 + ph_1 + \dots + p^{s-1}h_{s-1}) \rangle$ , where
  - (a)  $h_0, h_1, \dots, h_{s-1}$  are polynomials in  $\mathbb{Z}_{p^s}[x]$  such that  $h_0|(x^\beta - 1)$  and  $h_i|h_{i-1}$  for  $i = 1, \dots, s-1$ ,
  - (b)  $h'_0, h'_1, \dots, h'_{r-1}$  are polynomials in  $\mathbb{Z}_{p^r}[x]$  such that  $h'_0|(x^\alpha - 1)$  and  $h'_i|h'_{i-1}$  for  $i = 1, \dots, r-1$ .

$$(2) |C| = p^{d_1+d_2}, \text{ where } d_1 = \sum_{i=0}^{s-1} (s-i) \deg \widehat{h}_i \text{ and } d_2 = \sum_{i=0}^{r-1} (r-i) \deg \widehat{h}'_i.$$

*Proof.* Since  $f : \mathbb{Z}_{p^s} \rightarrow \mathbb{Z}_{p^r}$  is surjective, by the same argument of Corollary 3.7,  $C = \langle (h(x), 0), (\ell(x), g(x)) \rangle$ , where  $g(x) \in (\mathbb{Z}_{p^s})_\beta$  is a generator of a cyclic code over  $\mathbb{Z}_{p^s}$  of length  $\beta$ ,  $h(x) \in (\mathbb{Z}_{p^r})_\alpha$  is a generator of a cyclic code over  $\mathbb{Z}_{p^r}$  of length  $\alpha$  and  $\ell(x) \in (\mathbb{Z}_{p^r})_\alpha$  is a polynomial. By Corollary 2.9, there exists a set of polynomials  $h_0, h_1, \dots, h_{s-1}$  in  $\mathbb{Z}_{p^s}[x]$  such that  $h_0 | (x^\beta - 1)$  and  $h_i | h_{i-1}$  for  $i = 1, \dots, s-1$  and  $g(x) = h_0 + ph_1 + \dots + p^{s-1}h_{s-1}$ . Similarly, there exists a set of polynomials  $h'_0, h'_1, \dots, h'_{r-1}$  in  $\mathbb{Z}_{p^r}[x]$  such that  $h'_0 | (x^\alpha - 1)$  and  $h'_i | h'_{i-1}$  for  $i = 1, \dots, r-1$  and  $h(x) = h'_0 + ph'_1 + \dots + p^{r-1}h'_{r-1}$ . In this case,  $|C| = p^{d_1+d_2}$ , where  $d_1 = \sum_{i=0}^{s-1} (s-i) \deg \widehat{h}_i$  and  $d_2 = \sum_{i=0}^{r-1} (r-i) \deg \widehat{h}'_i$ .  $\square$

Recently,  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes generalized to codes over direct product of two finite chain rings in some special case [11]. More precisely, let  $R_1$  and  $R_2$  be two chain rings with the maximal ideals  $\mathfrak{m}_1 = \langle \gamma_1 \rangle$  and  $\mathfrak{m}_2 = \langle \gamma_2 \rangle$  of the nilpotency indexes  $e_1$  and  $e_2$ ; respectively. Let  $e_1 \leq e_2$ , and  $R_1$  and  $R_2$  have the same residue field  $R_1/\mathfrak{m}_1 = R_2/\mathfrak{m}_2 = \mathbb{F}$ . If  $a_1 \in R_1$  and  $a_2 \in R_2$ , then  $a_1$  and  $a_2$  can be uniquely written as follows:

$$a_1 = a_{1,0} + a_{1,1}\gamma_1 + \dots + a_{1,e_1-1}\gamma_1^{e_1-1}, \quad a_2 = a_{2,0} + a_{2,1}\gamma_2 + \dots + a_{2,e_2-1}\gamma_2^{e_2-1},$$

where the  $a_{1,i}$ s and  $a_{2,i}$ s can be viewed as elements in  $\mathbb{F}$  (see [14, Lemma 2]). Now define  $\psi : R_2 \rightarrow R_1$  by  $\psi(\sum_{i=0}^{e_2-1} a_i \gamma_2^i) = \sum_{i=0}^{e_1-1} a_i \gamma_1^i$ . It is easy to see that  $\psi$  is a ring homomorphism. Hence  $R_1$  is an  $R_2$ -algebra. For positive integers  $\alpha$  and  $\beta$ , an  $R_2$ -submodule  $C \subseteq R_1^\alpha \times R_2^\beta$  is called an  $R_1R_2$ -additive code. When  $\alpha$  and  $\beta$  are coprime integers with  $\text{Char}(R_i/\mathfrak{m})$ , the structure of these codes have been given (see [11, Theorem 4.3]). Now we obtain the structure of these codes as a result of the structure of  $SR$ -additive codes.

**Corollary 3.9** (Additive cyclic codes over direct product of finite chain rings). *With above assumptions, let  $C \subseteq (R_1)_\alpha \times (R_2)_\beta$  be an  $R_1R_2$ -additive cyclic code. If  $\alpha$  and  $\beta$  are coprime integers with  $\text{Char}(R_i/\mathfrak{m})$ , Then*

- (1)  $C = \langle (h'_0 + \gamma_1 h'_1 + \dots + \gamma_1^{e_1-1} h'_{e_1-1}, 0), (\ell(x), h_0 + \gamma_2 h_1 + \dots + \gamma_2^{e_2-1} h_{e_2-1}) \rangle$ , where
  - (a)  $h_0, h_1, \dots, h_{e_2-1}$  are polynomials in  $R_2[x]$  such that  $h_0 | (x^\beta - 1)$  and  $h_i | h_{i-1}$  for  $i = 1, \dots, e_2 - 1$ ,
  - (b)  $h'_0, h'_1, \dots, h'_{e_1-1}$  are polynomials in  $R_1[x]$  such that  $h'_0 | (x^\alpha - 1)$  and  $h'_i | h'_{i-1}$  for  $i = 1, \dots, e_1 - 1$ .
- (2)  $|C| = p^{d_1+d_2}$ , where  $d_1 = \sum_{i=0}^{e_2-1} (e_2 - i) \deg \widehat{h}_i$  and  $d_2 = \sum_{i=0}^{e_1-1} (e_1 - i) \deg \widehat{h}'_i$ .

*Proof.* By the same argument as Corollary 3.8, it follows from Theorem 3.3 and Theorem 2.8.  $\square$

Now we give new examples of  $SR$ -additive codes. First we give some examples of additive codes over direct products of chain rings that we can not



obtain their structures by [11]; see Corollaries 3.10, 3.11 and 3.12. Note that in [11], they considered an additive code  $C \subseteq R_1^\alpha \times R_2^\beta$  over the chain rings  $R_1$  and  $R_2$  in a case that  $\alpha$  and  $\beta$  are coprime integers with  $\text{Char}(R_i/\mathfrak{m})$ . But in the structure of SR-additive codes we haven't any restriction on  $\alpha$  and  $\beta$ .

Let  $R_1 = \mathbb{Z}_2$ ,  $R_2 = \mathbb{Z}_2 + u\mathbb{Z}_2 = \{0, 1, u, 1 + u\}$  such that  $u^2 = 0$  and  $R_3 = \mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2 = \{0, 1, u, 1 + u, u^2, 1 + u^2, 1 + u + u^2, u + u^2\}$  such that  $u^3 = 0$ . By the following maps,  $R_i$  is an  $R_j$ -algebra for  $1 \leq i < j \leq 3$ .

$$\begin{aligned} f_{2,1} : R_2 &\longrightarrow R_1; & \lambda_0 + \lambda_1 u &\longmapsto \lambda_0, \\ f_{3,1} : R_3 &\longrightarrow R_1; & \lambda_0 + \lambda_1 u + \lambda_2 u^2 &\longmapsto \lambda_0, \\ f_{3,2} : R_3 &\longrightarrow R_2; & \lambda_0 + \lambda_1 u + \lambda_2 u^2 &\longmapsto \lambda_0 + \lambda_1 u. \end{aligned}$$

We want to describe  $R_i R_j$ -additive cyclic codes for  $1 \leq i < j \leq 3$ . First we find the generators of  $R_1 R_2$ -additive cyclic codes which are known as  $\mathbb{Z}_2 \mathbb{Z}_2[u]$ -additive codes and studied in [3, 22].

**Corollary 3.10** ( $R_1 R_2$ -additive cyclic codes). *Let  $C \subseteq (R_1)_\alpha \times (R_2)_\beta$  be an  $R_1 R_2$ -additive cyclic code.*

- (1) *If  $\beta$  is odd, then  $C = \langle (h(x), 0), (\ell(x), g(x) + ua(x)) \rangle$  such that  $h(x)|(x^\alpha - 1) \pmod{2}$ ,  $\ell(x) \in (\mathbb{Z}_2)_\alpha$  and  $g(x) + ua(x) \in (R_2)_\beta$  with the same condition as the part (1) of Theorem 2.10.*
- (2) *If  $\beta$  is not odd, then*
  - (a)  *$C = \langle (h(x), 0), (\ell(x), g(x) + up(x)) \rangle$ , where  $h(x)$  and  $\ell(x)$  are such as (1).  $g(x)$  and  $p(x)$  have the same conditions as Theorem 2.10 part 2(a). Or*
  - (b)  *$C = \langle (h(x), 0), (\ell_1(x), g(x) + up(x)), (\ell_2(x), ua(x)) \rangle$ , where  $h(x)$  and  $\ell_i(x)$  are such as (1).  $g(x)$ ,  $p(x)$  and  $a(x)$  have the same conditions as Theorem 2.10 part 2(b).*

*Proof.* By Corollary 3.6,  $C = \langle (h(x), 0), (\ell_1, f_1), \dots, (\ell_r, f_r) \rangle$ , where  $h(x)$  is a monic polynomial over  $R_1$  such that  $h(x)|(x^\alpha - 1)$ . Also  $C_1 = \langle f_1, \dots, f_r \rangle$  is a linear cyclic code over  $R_2$ . Now we have the result by Theorem 2.10.  $\square$

**Corollary 3.11** ( $R_1 R_3$ -additive cyclic codes). *Let  $C \subseteq (R_1)_\alpha \times (R_3)_\beta$  be an  $R_1 R_3$ -additive cyclic code.*

- (1) *If  $\beta$  is odd, then  $C = \langle (h(x), 0), (\ell(x), g(x) + ua_1(x) + u^2 a_2(x)) \rangle$ , where  $h(x)$ ,  $\ell(x)$  are elements of  $\mathbb{Z}_2[x]$ ,  $h(x)|(x^\alpha - 1)$  in  $\mathbb{Z}_2[x]$  and  $g, a_1, a_2$  have the same conditions as Theorem 2.11 part (1).*
- (2) *If  $\beta$  is not odd, then*
  - (a)  *$C = \langle (h(x), 0), (\ell(x), g(x) + up_1(x) + u^2 p_2(x)) \rangle$ , where  $\ell, h$  are such as (1) and  $g, p_1, p_2$  have the same conditions as Theorem 2.11 part 2(a).*
  - (b)  *$C = \langle (h(x), 0), (\ell_1(x), g(x) + up_1(x) + u^2 p_2(x)), (\ell_2(x), u^2 a_2(x)) \rangle$ , where  $\ell_i$  and  $h$  are such as (1) and  $g, p_1, p_2, a_2$  have the same conditions as Theorem 2.11 part 2(b).*

- (c)  $C = \langle\langle (h(x), 0), (\ell_1(x), g(x) + up_1(x) + u^2p_2(x)), (\ell_2(x), ua_1(x) + u^2q_1(x)), (\ell_3, u^2a_2(x)) \rangle\rangle$ , where  $\ell_i$  and  $h$  are such as (1) and  $g, p_1, p_2, a_1, q_1, a_2$  have the same conditions as Theorem 2.11 part 2(c).

*Proof.* By the same argument as Corollary 3.10, it follows from Corollary 3.6 and Theorem 2.11.  $\square$

**Corollary 3.12** ( $R_2R_3$ -additive cyclic codes). *Let  $C \subseteq (R_2)_\alpha \times (R_3)_\beta$  be an  $R_2R_3$ -additive cyclic code.*

- (1) *If  $\beta$  and  $\alpha$  are odd, then  $C = \langle\langle (h(x), 0), (\ell(x), g(x) + ua_1(x) + u^2a_2(x)) \rangle\rangle$ , where  $h(x), \ell(x)$  are elements of  $(R_2)_\alpha$ .  $h(x)$  is a generator of a code such as Theorem 2.10 part (1) and  $g, a_1, a_2$  have the same conditions as Theorem 2.11 part (1).*
- (2) *If  $\beta$  is odd and  $\alpha$  is not odd, then*
  - (a)  $C = \langle\langle (g + up, 0), (\ell, f) \rangle\rangle$ , where  $g, p$  have the same conditions as Theorem 2.10 part 2(a).  $\ell \in (R_2)_\alpha$  and  $f \in (R_3)_\beta$  is a generator of a code such as Theorem 2.11 part (1). Or
  - (b)  $\langle\langle (g + up, 0), (ua, 0), (\ell, f) \rangle\rangle$ , where  $g, p, a$  are polynomials with the same conditions as Theorem 2.10 part 2(b).  $\ell \in (R_2)_\alpha$  and  $f \in (R_3)_\beta$  is a generator of a code such as Theorem 2.11 part (1).
- (3) *If  $\alpha$  is odd and  $\beta$  is not odd, then*
  - (a)  $\langle\langle (f, 0), (\ell, g + ua_1 + u^2a_2) \rangle\rangle$ , where  $\ell \in (R_2)_\alpha$ ,  $f$  is a generator of a code such as Theorem 2.10 part (1) and  $g, a_1, a_2$  are such as Theorem 2.11 part 2(a). Or
  - (b)  $C = \langle\langle (f, 0), (\ell_1, g + up_1 + u^2p_2), (\ell_2, u^2a_2) \rangle\rangle$ , where  $f$  and  $\ell_i$  are such as (a) and  $g, p_1, p_2, a_2$  have the same conditions as Theorem 2.11 part 2(b). Or
  - (c)  $C = \langle\langle (f, 0), (\ell_1, g + up_1 + u^2p_2), (\ell_2, ua_1 + u^2q_1), (\ell_3, u^2a_2) \rangle\rangle$ , where  $f$  and  $\ell_i$  are such as (a) and  $g, p_1, p_2, a_1, a_2, q_1$  have the same conditions as Theorem 2.11 part 2(c).
- (4) *If  $\alpha$  and  $\beta$  are not odd, then we have one of the following states.*
  - (a)  $C = \langle\langle (g_1, 0), (\ell_1, f_1) \rangle\rangle$ , where  $g_1$  is a generator of a code in Theorem 2.10 part 2(a),  $f_1$  is a generator of a code in Theorem 2.11 part 2(a) and  $\ell_1$  is an elements of  $(R_2)_\alpha$ .
  - (b)  $C = \langle\langle (g_1, 0), (\ell_1, f_1), (\ell_2, f_2) \rangle\rangle$ , where  $g_1$  is a generator of a code in Theorem 2.10 part 2(a),  $f_i$  are generators of a code in Theorem 2.11 part 2(b) and  $\ell_i$  are elements of  $(R_2)_\alpha$ .
  - (c)  $C = \langle\langle (g_1, 0), (\ell_1, f_1), (\ell_2, f_2), (\ell_3, f_3) \rangle\rangle$ , where  $g_1$  is a generator of a code in Theorem 2.10 part 2(a),  $f_i$  are generators of a code in Theorem 2.11 part 2(c) and  $\ell_i$  are elements of  $(R_2)_\alpha$ .
  - (d)  $C = \langle\langle (g_1, 0), (g_2, 0), (\ell_1, f_1) \rangle\rangle$ , where  $g_i$  are generators of a code in Theorem 2.10 part 2(b),  $f_1$  is a generator of a code in Theorem 2.11 part 2(a) and  $\ell_1$  is an element of  $(R_2)_\alpha$ .

- (e)  $C = \langle (g_1, 0), (g_2, 0), (\ell_1, f_1), (\ell_2, f_2) \rangle$ , where  $g_i$  are generators of a code in Theorem 2.10 part 2(b),  $f_i$  are generators of a code in Theorem 2.11 part 2(b) and  $\ell_i$  is an element of  $(R_2)_\alpha$ .
- (f)  $C = \langle (g_1, 0), (g_2, 0), (\ell_1, f_1), (\ell_2, f_2), (\ell_3, f_3) \rangle$ , where  $g_i$  are generators of a code in Theorem 2.10 part 2(b),  $f_i$  are generators of a code in Theorem 2.11 part 2(c) and  $\ell_i$  are elements of  $(R_2)_\alpha$ .

*Proof.* By Theorem 3.3,  $C = \langle (g_1, 0), \dots, (g_s, 0), (h_1, f_1), \dots, (h_r, f_r) \rangle_{R[x]}$  such that  $C_1 = \langle f_1, \dots, f_r \rangle_{R_3[x]}$  is a cyclic linear code over  $R_3$  of length  $\beta$  and  $C_2 = \langle g_1, \dots, g_s \rangle_{R_3[x]}$  is a cyclic  $R_3$ -additive code over  $R_2$  of length  $\alpha$ . Since  $f_{3,2} : R_3 \rightarrow R_2$  is a surjective map,  $C_2$  is a linear code over  $R_2$ . Now the result follows from Theorems 2.10 and 2.11.  $\square$

Now we give some examples that the ring  $R$  in  $SR$ -additive codes is not a chain ring (moreover this ring is not a Frobenius ring). Let  $R_4 = \mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2 = \{0, 1, u, v, 1+u, 1+v, u+v, 1+u+v\}$  such that  $u^2 = v^2 = uv = 0$ . This ring is not a chain ring. Moreover  $R_4$  is a non Frobenius ring. Consider the rings  $R_1 = \mathbb{Z}_2$  and  $R_2 = \mathbb{Z}_2 + u\mathbb{Z}_2$  in above corollaries. It is easy to see that the following maps are ring homomorphisms:

$$\begin{aligned} f_{4,1} : R_4 &\longrightarrow R_1; & \lambda_1 + \lambda_2 u + \lambda_3 v &\longmapsto \lambda_1, \\ f_{4,2} : R_4 &\longrightarrow R_2; & \lambda_1 + \lambda_2 u + \lambda_3 v &\longmapsto \lambda_1 + \lambda_2 u. \end{aligned}$$

Hence  $R_4$  is an  $R_i$ -algebra for  $i = 1, 2$ . Now we want to describe  $R_1 R_4$  and  $R_2 R_4$ -additive cyclic codes.

**Corollary 3.13** ( $R_1 R_4$ -additive cyclic codes). *Let  $C \subseteq (R_1)_\alpha \times (R_4)_\beta$  be an  $R_1 R_4$ -additive cyclic code. Then  $C = \langle (f, 0), (h_1, g + up_1 + vp_2), (h_2, ua_1 + vq_1), (h_3, va_2) \rangle$ , where  $f|(x^\alpha - 1)$ ,  $h_i \in (R_1)_\alpha$  and  $p_1, p_2, q_1, a_1, a_2$  have the same conditions as Theorem 2.12. Moreover if  $\beta$  is odd, then  $C = \langle (f, 0), (h_1, g + ua_1), (h_2, va_2) \rangle$ , where  $a_2 | a_1 | g | (x^n - 1)$ .*

*Proof.* It follows from Corollary 3.6 and Theorem 2.12.  $\square$

**Corollary 3.14** ( $R_2 R_4$ -additive cyclic codes). *Let  $C \subseteq (R_2)_\alpha \times (R_4)_\beta$  be an  $R_2 R_4$ -additive cyclic code. Then*

- (1) *If  $\alpha$  is odd, then  $C = \langle (g + ua, 0), (h_1, g_1 + up_1 + vp_2), (h_2, ua_1 + vq_1), (h_3, va_2) \rangle$ , where  $g$  and  $a$  are polynomials in  $\mathbb{Z}_2[x]$  such that  $a|g|(x^\alpha - 1) \pmod{2}$ ,  $h_i \in (R_2)_\alpha$  and  $p_1, p_2, q_1, g_1, a_1, a_2$  have the same conditions as Theorem 2.12.*
- (2) *If  $\alpha$  is not odd, then*
  - (a)  $C = \langle (g + up, 0), (h_1, g_1 + up_1 + vp_2), (h_2, ua_1 + vq_1), (h_3, va_2) \rangle$  such that  $g|(x^\alpha - 1) \pmod{2}$ ,  $(g + up)|(x^\alpha - 1)$  in  $\mathbb{Z}_2 + u\mathbb{Z}_2$  and  $g|p(\frac{x^\alpha - 1}{g})$ .  
Or
  - (b)  $C = \langle (ua, 0), (g + up, 0), (h_1, g_1 + up_1 + vp_2), (h_2, ua_1 + vq_1), (h_3, va_2) \rangle$  such that  $g, a$  and  $p$  are polynomials in  $\mathbb{Z}_2[x]$ .  $a|g|(x^\alpha - 1) \pmod{2}$ ,  $a|p(\frac{x^\alpha - 1}{g})$  and  $\deg a > \deg p$ .

Where  $h_i \in (R_2)_\alpha$  and  $p_1, p_2, q_1, g_1, a_1, a_2$  have the same conditions as Theorem 2.12.

*Proof.* It follows from Theorem 3.3 and Theorem 2.12.  $\square$

In the above examples the ring homomorphisms between  $R_i$  and  $R_j$  are surjective, hence cyclic  $R_i R_j$ -additive codes are constructed by linear cyclic codes over  $R_i$  and  $R_j$ . But when  $f$  is not surjective to construct cyclic  $SR$ -additive codes we need the structure of  $R$ -additive codes over  $S$ . See the following examples.

**Example 3.15.** Let  $R_1 = \mathbb{Z}_2$  and  $R_2 = \mathbb{Z}_2 + u\mathbb{Z}_2$  be the rings in above corollaries. Then  $R_2$  is an  $R_1$ -algebra with the including map. Let  $C \subseteq (R_2)_\alpha \times (R_1)_\beta$  be an  $R_2 R_1$ -additive cyclic code. Then  $C = \langle (g_1, 0), \dots, (g_s, 0), (h, f) \rangle$ , where  $f|(x^\beta - 1)$ ,  $h \in (R_2)_\alpha$ , and  $C_1 = \langle g_1, \dots, g_s \rangle$  is a cyclic  $R_1$ -additive code over  $R_2$  ( $C_1$  is an additive cyclic code over  $R_2$ ).

**Example 3.16.** Let  $R = GR(p^s, m)$  and  $S = R[\xi] = GR(p^s, m\ell)$  be the Galois extension of  $R$ . Then  $S$  is an  $R$ -algebra with the including map. Let  $C \subseteq S_\alpha \times R_\beta$  be an  $SR$ -additive cyclic code. If  $\gcd(\beta, p) = 1$  and  $\gcd(\alpha, p) = 1$ , then  $C = \langle (g_1, 0), \dots, (g_\ell, 0), (h, f) \rangle$ , where  $C_2 = \langle f \rangle$  is a cyclic code over  $R$ ,  $C_1 = \langle g_1, \dots, g_\ell \rangle$  is a cyclic  $R$ -additive code over  $S$  of length  $\alpha$  and  $h \in S_\alpha$  is a polynomial.

#### 4. Duality of $SR$ -additive codes

In this section, we define a bilinear form on  $SR$ -additive codes which is a generalization of the bilinear forms over  $\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes in [2],  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes in [3] and  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes in [10].

**Definition 4.1.** Let  $\tau : S \rightarrow R$  be an  $R$ -module homomorphism, then

$$\begin{aligned} \beta' : (S^\alpha \times R^\beta) \times (S^\alpha \times R^\beta) &\longrightarrow R \\ ((x_1, y_1), (x_2, y_2)) &\longmapsto \tau(x_1.x_2) + (y_1.y_2) \end{aligned}$$

is an  $R$ -bilinear form where  $x_1.x_2$  and  $y_1.y_2$  are standard inner products. For an  $SR$ -additive code  $C$ ,  $C^\perp$  is the dual of  $C$  with respect to  $\beta'$ .

**Proposition 4.2.** Let  $R$  be a chain ring with maximal ideal  $\mathfrak{m} = \langle \gamma \rangle$  of nilpotency index  $e$ . If  $\beta'$  is a bilinear form on  $(R/\mathfrak{m})R$ -additive codes defined by an  $R$ -module homomorphism  $\tau : R/\mathfrak{m} \rightarrow R$ , then there is a unit element  $a \in R$  such that

$$\begin{aligned} \beta' : ((R/\mathfrak{m})^\alpha \times R^\beta) \times ((R/\mathfrak{m})^\alpha \times R^\beta) &\longrightarrow R \\ ((\bar{x}_1, y_1), (\bar{x}_2, y_2)) &\longmapsto a\gamma^{e-1}(x_1.x_2) + (y_1.y_2). \end{aligned}$$

Where  $\bar{x}_1 = (x_{1,i} + \mathfrak{m})$ ,  $\bar{x}_2 = (x_{2,i} + \mathfrak{m})$ , and  $x_1 = (x_{1,i})$  and  $x_2 = (x_{2,i})$ .

*Proof.* By the definition of  $\beta'$ , it suffices to determine  $\text{Hom}_R(R/\mathfrak{m}, R)$ . But we have the following  $R$ -module isomorphism

$$\begin{aligned} \text{Hom}_R(R/\mathfrak{m}, R) &\longrightarrow \text{Ann}_R(\mathfrak{m}) \\ \tau &\longmapsto \tau(1 + \mathfrak{m}). \end{aligned}$$

Since  $R$  is a chain ring and  $\text{Ann}_R(\mathfrak{m})$  is an ideal of  $R$ ,  $\text{Ann}_R(\mathfrak{m}) = \langle \gamma^j \rangle$  for some  $j$ ;  $1 \leq j \leq e$ . Clearly  $\gamma^{e-1}\mathfrak{m} = 0$ . On other hand  $\gamma^{e-2}\gamma \neq 0$ . Hence  $\text{Ann}_R(\mathfrak{m}) = \langle \gamma^{e-1} \rangle$ . Thus there is a unit element  $a \in R \setminus \mathfrak{m}$  such that  $\tau(1 + \mathfrak{m}) = a\gamma^{e-1}$ . Hence for  $r + \mathfrak{m} \in R/\mathfrak{m}$ ,  $\tau(r + \mathfrak{m}) = r\tau(1 + \mathfrak{m}) = ra\gamma^{e-1}$ . This completes the proof.  $\square$

Now we give some examples of this bilinear form over  $SR$ -additive codes, which we see some of them in [2] and [3].

**Corollary 4.3** (The bilinear form of  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes). *The following bilinear form is the only form on  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes defined by Definition 4.1.*

$$\beta' : (\mathbb{Z}_2^\alpha \times \mathbb{Z}_4^\beta) \times (\mathbb{Z}_2^\alpha \times \mathbb{Z}_4^\beta) \longrightarrow \mathbb{Z}_4, ((x_1, y_1), (x_2, y_2)) \longmapsto 2(x_1 \cdot x_2) + (y_1 \cdot y_2).$$

Where the elements  $x_1$  and  $x_2$  in the inner product  $2(x_1 \cdot x_2)$  are considered as elements of  $\mathbb{Z}_4^\beta$ ; naturally.

*Proof.*  $\mathbb{Z}_4$  is a chain ring with maximal ideal  $2\mathbb{Z}_4$  of nilpotency index 2. Also  $\frac{\mathbb{Z}_4}{2\mathbb{Z}_4} \cong \mathbb{Z}_2$ . Now we have the result by Proposition 4.2.  $\square$

**Proposition 4.4** (The bilinear forms of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes,  $r < s$ ). *Let  $\beta'$  be a bilinear form on  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes defined by Definition 4.1. Then  $\beta'$  is defined as follows:*

$$\begin{aligned} \beta' : (\mathbb{Z}_{p^r}^\alpha \times \mathbb{Z}_{p^s}^\beta) \times (\mathbb{Z}_{p^r}^\alpha \times \mathbb{Z}_{p^s}^\beta) &\longrightarrow \mathbb{Z}_{p^s}, \\ ((x_1, y_1), (x_2, y_2)) &\longmapsto ap^{s-r}(x_1 \cdot x_2) + (y_1 \cdot y_2), \end{aligned}$$

where  $a \in \mathbb{Z}_{p^s}$  and the elements  $x_1$  and  $x_2$  in the inner product  $ap^{s-r}(x_1 \cdot x_2)$  are considered as elements of  $\mathbb{Z}_{p^s}^\beta$ ; naturally.

*Proof.*  $\text{Hom}_{\mathbb{Z}_{p^s}}(\mathbb{Z}_{p^r}, \mathbb{Z}_{p^s}) = \text{Hom}_{\mathbb{Z}_{p^s}}(\frac{\mathbb{Z}_{p^s}}{p^r\mathbb{Z}_{p^s}}, \mathbb{Z}_{p^s}) \cong \text{Ann}_{\mathbb{Z}_{p^s}}(p^r\mathbb{Z}_{p^s}) = \langle p^{s-r} \rangle$ . Now by the same argument of Proposition 4.2 we have the result.  $\square$

Let  $R_1$  and  $R_2$  be the finite chain rings with the assumptions of Corollary 3.9. We have the isomorphism  $\psi : \frac{R_2}{\gamma_2^{\epsilon_1}R_2} \rightarrow R_1$ . Let  $p : R_2 \rightarrow \frac{R_2}{\gamma_2^{\epsilon_1}R_2}$  be defined naturally. Hence  $\iota = p^{-1}\psi^{-1} : R_1 \rightarrow R_2$  is well defined, where  $p^{-1}$  is a right inverse of  $p$ . The following proposition gives the bilinear forms over direct product of chain rings.

**Proposition 4.5** (The bilinear forms of additive codes over product of chain rings). *Let  $R_1$  and  $R_2$  be the finite chain rings with the assumptions Corollary*

3.9. If  $\beta'$  is a bilinear form on  $R_1R_2$ -additive codes defined by Definition 4.1, then  $\beta'$  is defined as follows:

$$\begin{aligned} \beta' : (R_1^\alpha \times R_2^\beta) \times (R_1^\alpha \times R_2^\beta) &\longrightarrow R_2, \\ ((x_1, y_1), (x_2, y_2)) &\longmapsto a\gamma^{e_2-e_1}l(x_1.x_2) + (y_1.y_2), \end{aligned}$$

where  $a \in R_2$ .

*Proof.*  $\text{Hom}_{R_2}(R_1, R_2) = \text{Hom}_{R_2}(\frac{R_2}{\gamma_2^{e_1}R_2}, R_2) \cong \text{Ann}_{R_2}(\gamma_2^{e_1}R_2) = \gamma_2^{e_2-e_1}R_2$ . Now by the same argument of Proposition 4.2 we have the result.  $\square$

**Proposition 4.6** (The bilinear forms of  $R_iR_j$ -additive codes,  $i < j$ ). *Let  $R_i$  and  $R_j$  be such as Corollaries 3.10, 3.11, 3.12, 3.13, 3.14. Then, we have the following bilinear forms on  $R_iR_j$ -additive codes.*

$$\begin{aligned} \beta_{1,2} : (R_1^\alpha \times R_2^\beta) \times (R_1^\alpha \times R_2^\beta) &\rightarrow R_2, ((x_1, y_1), (x_2, y_2)) \mapsto u(x_1.x_2) + y_1.y_2, \\ \beta_{1,3} : (R_1^\alpha \times R_3^\beta) \times (R_1^\alpha \times R_3^\beta) &\rightarrow R_3, ((x_1, y_1), (x_2, y_2)) \mapsto u^2(x_1.x_2) + y_1.y_2, \\ \beta_{2,3} : (R_2^\alpha \times R_3^\beta) \times (R_2^\alpha \times R_3^\beta) &\rightarrow R_3, ((x_1, y_1), (x_2, y_2)) \mapsto au(x_1.x_2) + y_1.y_2, \\ \beta_{1,4} : (R_1^\alpha \times R_4^\beta) \times (R_1^\alpha \times R_4^\beta) &\rightarrow R_4, ((x_1, y_1), (x_2, y_2)) \mapsto h(x_1.x_2) + y_1.y_2, \\ \beta_{2,4} : (R_2^\alpha \times R_4^\beta) \times (R_2^\alpha \times R_4^\beta) &\rightarrow R_4, ((x_1, y_1), (x_2, y_2)) \mapsto h(x_1.x_2) + y_1.y_2, \end{aligned}$$

where  $a \in R_3$ ,  $h \in R_4u + R_4v$ .

*Proof.*  $R_2$  and  $R_3$  are chain rings with maximal ideals  $R_2\langle u \rangle$  and  $R_3\langle u \rangle$  of nilpotency indices 2 and 3; respectively. Also  $\frac{R_2}{\langle u \rangle} \cong \frac{R_3}{\langle u \rangle} \cong R_1$ . Hence we have the bilinear forms  $\beta_{1,2}$  and  $\beta_{1,3}$  by Proposition 4.2. To obtain  $\beta_{2,3}$ ,  $\beta_{1,4}$  and  $\beta_{2,4}$  note that

$$\begin{aligned} \text{Hom}_{R_3}(R_2, R_3) &\cong \text{Hom}_{R_3}(\frac{R_3}{\langle u^2 \rangle}, R_3) \cong \text{Ann}_{R_3}(\langle u^2 \rangle) = \langle u \rangle, \\ \text{Hom}_{R_4}(R_1, R_4) &\cong \text{Hom}_{R_4}(\frac{R_4}{\langle u, v \rangle}, R_4) \cong \text{Ann}_{R_4}(\langle u, v \rangle) = R_4u + R_4v, \\ \text{Hom}_{R_4}(R_2, R_4) &\cong \text{Hom}_{R_4}(\frac{R_4}{\langle v \rangle}, R_4) \cong \text{Ann}_{R_4}(\langle v \rangle) = R_4u + R_4v. \end{aligned}$$

Now by the same argument of the proof of Proposition 4.2 we have the result.  $\square$

**Proposition 4.7.** *Let  $\tau : S \rightarrow R$  be an  $R$ -module homomorphism and  $C \subseteq S^\alpha \times R^\beta$  be an  $SR$ -additive cyclic code. If  $C^\perp$  is the dual of  $C$  with respect to the bilinear form defined by  $\tau$  in Definition 4.1, then  $C^\perp$  is an  $SR$ -additive cyclic code.*

*Proof.* Clearly  $C^\perp$  is an  $R$ -submodule of  $S^\alpha \times R^\beta$ , hence  $C^\perp$  is an  $SR$ -additive code. Now let

$$(x, y) = (x_0 \cdots x_{\alpha-1}, y_0 \cdots y_{\beta-1}) \in C^\perp \text{ and}$$

$$\phi(x, y) = (x_{\alpha-1} \cdots x_{\alpha-2}, y_{\beta-1} \cdots y_{\beta-2}).$$

Let  $j = \text{lcm}(\alpha, \beta)$  and  $(v, w) \in C$ . Since  $C$  is cyclic,  $\phi^{j-1}(v, w) \in C$ . Now

$$\begin{aligned} (v, w) \cdot \phi(x, y) &= \tau(v \cdot \phi(x)) + w \cdot \phi(y) \\ &= \tau(\phi^{j-1}(v) \cdot x) + \phi^{j-1}(w) \cdot y \\ &= \phi^{j-1}(v, w) \cdot (x, y) = 0. \end{aligned}$$

Therefore  $\phi(x, y) \in C^\perp$  and hence  $C^\perp$  is cyclic.  $\square$

### 5. Singleton Bounds for SR-additive codes

Aydogdu and Siap obtained some bounds on the minimum distance of  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes [4]. In this section, we generalize the definitions of weight functions and Gray maps on the classes of  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$  and  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes to SR-additive codes. We obtain singleton bounds for SR-additive codes. As results, singleton bounds for  $\mathbb{Z}_2\mathbb{Z}_2[u]$  and  $\mathbb{Z}_2[u]\mathbb{Z}_2$ -additive codes are given.

**Definition 5.1.** Let  $T$  be a commutative finite ring. For every  $x = (x_1, \dots, x_n) \in T^n$  and  $t \in T$ , the complete weight of  $x$  is defined by

$$n_t(x) := |\{i : x_i = t\}|.$$

For  $t \in T \setminus \{0\}$ , let  $a_t$  be a positive integer, and set  $a_0 = 0$ . The general weight function over  $T$  is defined as follows:

$$\omega_T(x) := \sum_{t \in T} a_t n_t(x).$$

Now let  $\omega_R$  and  $\omega_S$  be two weight functions over  $R$  and  $S$ . A weight function  $\omega$  over  $S^\alpha \times R^\beta$  is defined as follows: for  $(x, y) \in S^\alpha \times R^\beta$ ,  $\omega(x, y) = \omega_S(x) + \omega_R(y)$ .

**Definition 5.2.** Let  $n_s \in \mathbb{N}$  be a positive integer. A map  $\phi : R \rightarrow S^{n_s}$  with the following conditions is called a gray map:

- (a)  $\phi$  is injective.
- (b) for  $x, y \in R$ ,  $\omega_R(x - y) = \omega_S(\phi(x) - \phi(y))$ .

A gray map  $\phi$  is called  $R$ -linear if  $\phi$  is an  $R$ -module homomorphism.  $\phi$  generalize on  $R^\beta$  naturally; for  $x = (x_1, \dots, x_\beta) \in R^\beta$ ,  $\phi(x) = (\phi(x_1), \dots, \phi(x_\beta)) \in S^{n_s\beta}$ . We generalize  $\phi$  to a map  $\Phi$  over  $S^\alpha \times R^\beta$  as follows:

$$\begin{aligned} \Phi : S^\alpha \times R^\beta &\longrightarrow S^{\alpha+n_s\beta} \\ (x, y) &\longmapsto (x, \phi(y)). \end{aligned}$$

Clearly for any  $(x, y) \in S^\alpha \times R^\beta$ ,  $\omega(x, y) = \omega_S(\Phi(x, y))$ . Moreover  $\Phi$  is an injective map. Now let  $C \subseteq S^\alpha \times R^\beta$  be an SR-additive code, the minimum general weight of  $C$  is

$$d_\omega(C) := \min\{\omega(x, y) : (x, y) \in C \setminus \{0\}\}.$$

Let  $A_s = \max\{a_s : s \in S\}$ . The following theorem gives Singleton Bounds for  $SR$ -additive codes.

**Theorem 5.3.** *With above notations, let  $R$  be a chain ring and  $S$  be an  $R$ -algebra with a ring homomorphism  $f : R \rightarrow S$ . If  $C \subseteq S^\alpha \times R^\beta$  is an  $SR$ -additive code such that  $\Phi(C) \subseteq S^{\alpha+n_s\beta}$  is an  $R$ -additive code, then*

- (1) *If  $S$  is a principal ideal ring and  $f$  is surjective, then  $\lfloor \frac{d_\omega(C)-1}{A_s} \rfloor \leq \alpha + n_s\beta - \text{rank}(C)$ .*
- (2) *If  $S$  is a free  $R$ -algebra of dimension  $m$ , then  $\lfloor \frac{d_\omega(C)-1}{A_s} \rfloor \leq \alpha + n_s\beta - \lceil \frac{\text{rank}(C)}{m} \rceil$ .*

*Proof.* (1)  $\Phi(C)$  is an  $R$ -additive code. Since  $f$  is surjective, hence  $\Phi(C)$  is a linear code over  $S$ . If  $d_{\omega_s}(\Phi(C))$  is the minimum weight of  $\Phi(C)$  with respect to the weight function  $\omega_s$ , then by Theorem 3.7 of [20], we have that  $\lfloor \frac{d_{\omega_s}(\Phi(C))-1}{A_s} \rfloor \leq \alpha + n_s\beta - \text{rank}(\Phi(C))$ . But  $d_{\omega_s}(\Phi(C)) = d_\omega(C)$  and  $\text{rank}(\Phi(C)) = \text{rank}(C)$ . This completes the proof of part (1).

(2)  $\Phi(C)$  is an  $R$ -additive code and  $S$  is a free  $R$ -algebra. Hence by Theorem 2.7,  $\lfloor \frac{d_{\omega_s}(\Phi(C))-1}{A_s} \rfloor \leq \alpha + n_s\beta - \lceil \frac{\text{rank}(\Phi(C))}{m} \rceil$ . Since  $d_{\omega_s}(\Phi(C)) = d_\omega(C)$  and  $\text{rank}(\Phi(C)) = \text{rank}(C)$ , we have the result.  $\square$

**Corollary 5.4.** *With above assumptions, let  $\omega_S = \omega_H$  be the Hamming weight. Then*

- (1) *If  $S$  is a free  $R$ -algebra of dimension  $m$ , then  $d_\omega(C) \leq \alpha + n_s\beta - \lceil \frac{\text{rank}(C)}{m} \rceil + 1$ .*
- (2) *If  $S$  is a principal ideal ring and  $f$  is surjective, then  $d_\omega(C) \leq \alpha + n_s\beta - \text{rank}(C) + 1$ .*

*Remark 5.5.* Let  $R$  be a finite commutative ring and  $S$  be a finite  $R$ -algebra with a surjective ring homomorphism  $f : R \rightarrow S$ . With above assumptions, if  $\omega_S = \omega_H$  is the Hamming weight, then  $d_\omega(C) \leq \alpha + n_s\beta - \log_{|S|}|C| + 1$ .

*Proof.* Since  $f$  is surjective,  $\Phi(C)$  is a linear code over  $S$ . By the singleton bound for linear codes we have the result.  $\square$

**Example 5.6.** Consider  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes. The Lee weight over  $\mathbb{Z}_2[u] = \{0, 1, u, 1+u\}$  is defined as follows:

$$\omega_L(0) = 0, \omega_L(1) = 1, \omega_L(u) = 2, \omega_L(1+u) = 1.$$

For any element  $(x, y) = (x_0, \dots, x_{\alpha-1}; y_0, \dots, y_{\beta-1}) \in \mathbb{Z}_2^\alpha \times \mathbb{Z}_2[u]^\beta$ , the weight function  $\omega$  is defined in the following way:

$$\omega(x, y) = \sum_{i=0}^{\alpha-1} \omega_H(x_i) + \sum_{i=0}^{\beta-1} \omega_L(y_i),$$



where  $\omega_H$  is the hamming weight over  $\mathbb{Z}_2$  and  $\omega_L$  is the Lee weight over  $\mathbb{Z}_2[u]$ . Now we have the following Gray map:

$$\begin{aligned}\phi : \mathbb{Z}_2[u] &\longrightarrow \mathbb{Z}_2^2 \\ a + bu &\longmapsto (b, a + b).\end{aligned}$$

It is easy to see that  $\omega_L(a + bu) = \omega_H(b, a + b)$  for any element  $a + bu \in \mathbb{Z}_2[u]$ . This map generalizes to the Gray map  $\Phi$ :

$$\begin{aligned}\phi : \mathbb{Z}_2^\alpha \times \mathbb{Z}_2[u]^\beta &\longrightarrow \mathbb{Z}_2^{\alpha+2\beta} \\ (x, y) &\longmapsto (x, \phi(y)).\end{aligned}$$

Clearly  $\omega(x, y) = \omega_H(\phi(x, y))$ . Now if  $C \subseteq \mathbb{Z}_2^\alpha \times \mathbb{Z}_2[u]^\beta$  is a  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive code, then we have the following bounds for minimum weight  $d_\omega(C)$ :

$$\begin{aligned}d_\omega(C) &\leq \alpha + 2\beta - \text{rank}(C) + 1, \\ d_\omega(C) &\leq \alpha + 2\beta - \log_2 |C| + 1.\end{aligned}$$

**Example 5.7.** Consider  $\mathbb{Z}_2[u]\mathbb{Z}_2$ -additive codes in Example 3.15. The subset  $C \subseteq \mathbb{Z}_2[u]^\alpha \times \mathbb{Z}_2^\beta$  is a  $\mathbb{Z}_2[u]\mathbb{Z}_2$ -additive code if and only if  $C$  is a subgroup under addition. For any element  $(x, y) \in \mathbb{Z}_2[u]^\alpha \times \mathbb{Z}_2^\beta$ , the weight function  $\omega$  is defined as follows:

$$\omega(x, y) = \omega_L(x) + \omega_H(y),$$

where  $\omega_L$  is the Lee weight over  $\mathbb{Z}_2[u]$  in above example and  $\omega_H$  is the Hamming weight over  $\mathbb{Z}_2$ . Let  $j : \mathbb{Z}_2 \rightarrow \mathbb{Z}_2[u]$  be the including map. We define a Gray map as follows:

$$\begin{aligned}\Phi : \mathbb{Z}_2[u]^\alpha \times \mathbb{Z}_2^\beta &\longrightarrow \mathbb{Z}_2[u]^{\alpha+\beta} \\ (x, y) &\longmapsto (x, j(y)).\end{aligned}$$

It is easy to see that  $\omega(x, y) = \omega_L(\Phi(x, y))$ . Since  $\mathbb{Z}_2[u]$  is a free  $\mathbb{Z}_2$ -algebra of dimension 2, by Theorem 5.3, we have the following bound for minimum weight:

$$\lfloor \frac{d_\omega(C) - 1}{2} \rfloor \leq \alpha + \beta - \lceil \frac{\text{rank}(C)}{2} \rceil.$$

## 6. One weight SR-additive codes

Recently, Dougherty et al. described one weight  $\mathbb{Z}_2\mathbb{Z}_4$ -additive codes [15]. In this section, we generalize this theory over SR-additive codes where  $S$  and  $R$  are chain rings. As applications of the theory, we obtain some results on one weight  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes (with respect to homogeneous weight) and one weight  $\mathbb{Z}_{2^r}\mathbb{Z}_{2^s}$ -additive codes (with respect to Lee weight). In particular, we obtain the structure of one weight  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes. First we remind the following definition of a pre-homogeneous weight in [23].

**Definition 6.1.** Let  $T$  be a commutative finite ring. A weight function  $\omega_T : T \rightarrow \mathbb{R}$  is pre-homogeneous if  $a_0 = 0$  and there exists a constant  $c_T > 0$  such that for  $t \neq 0$ ,

$$\sum_{t' \in \langle t \rangle} a_{t'} = c_T |\langle t \rangle|,$$

where  $\langle t \rangle$  is the principal ideal generated by an element  $t$  of  $T$ . In this case  $c_T$  is called the average weight.

**Example 6.2** ([23, Example 3.7]). Let  $R = \mathbb{Z}_{2^s}$ . Then Lee weight is pre-homogeneous with average weight  $c_R = 2^{s-2}$ .

**Lemma 6.3.** Let  $R$  and  $S$  be two chain rings, where  $S$  is an  $R$ -algebra with a surjective ring homomorphism  $f : R \rightarrow S$ . Also let  $\omega_S$  and  $\omega_R$  be two pre-homogeneous weights with average weights  $c_R$  and  $c_S$ . If  $C \subseteq S^\alpha \times R^\beta$  is an  $SR$ -additive code with no all zero columns, then

$$\sum_{c \in C} \omega(c) = |C|(\alpha c_S + \beta c_R),$$

where  $\omega$  is the weight function defined by  $\omega_S$  and  $\omega_R$  over  $S^\alpha \times R^\beta$ .

*Proof.* Let  $S$  a chain ring with maximal ideal  $\mathfrak{m} = \langle \gamma \rangle$  of nilpotency index  $v$ . Write the codewords of  $C$  as rows of a matrix  $G$ . Consider the column  $j$  of  $G$ , where  $1 \leq j \leq \alpha$ . Let  $J$  be the ideal of  $S$  generated by all elements of the column  $j$ . Then there exists  $1 \leq t \leq v$  that  $J = \langle \gamma^t \rangle$ . Since  $f$  is surjective and  $C$  is an  $R$ -submodule, any element of  $J$  is an element of the column  $j$ . Now we show that any two elements of  $J$  have the same repetition number in the column  $j$ . Consider two elements  $\gamma^t$  and  $\gamma^{t+1}$  of  $J$  with the repetition numbers  $n_t$  and  $n_{t+1}$ , respectively. Since  $\gamma^{t+1} = \gamma \gamma^t$ , hence  $n_t \leq n_{t+1}$ . On the other hand  $\gamma^t(\gamma - 1) = \gamma^{t+1} - \gamma^t$ . Since  $\gamma - 1$  is invertible,  $\gamma^t = (\gamma - 1)^{-1}(\gamma^{t+1} - \gamma^t)$ . Hence  $n_{t+1} \leq n_t$  and hence  $n_t = n_{t+1}$ . Thus all elements of  $J$  have the same repetition number  $\frac{|C|}{|J|}$  in the column  $j$ . Therefore the sum of the weights of all elements of the column  $j$  is equal to

$$\frac{|C|}{|J|} \left( \sum_{s \in J} a_s \right) = \frac{|C|}{|J|} (c_S |J|) = |C| c_S.$$

By the same argument, the sum of the weights of all elements of the columns of  $\beta$  coordinates is equal to  $|C| c_R$ . Therefore

$$\sum_{c \in C} \omega(c) = |C|(\alpha c_S + \beta c_R). \quad \square$$

**Theorem 6.4.** With the assumptions of above lemma, let  $C \subseteq S^\alpha \times R^\beta$  be a one weight  $SR$ -additive code with weight  $m$  such that there exists no zero columns in the generator matrix of  $C$ . Then there exists a unique positive integer  $\lambda$  such that  $m = \lambda |C|$  and  $\alpha c_S + \beta c_R = \lambda(|C| - 1)$ .

*Proof.* By above lemma, we have that

$$\sum_{c \in C} \omega(c) = |C|(\alpha c_S + \beta c_R).$$

On the other hand, the sum of the weights of all codewords is  $(|C|-1)m$ . Hence  $|C|(\alpha c_S + \beta c_R) = (|C|-1)m$ . But  $\gcd(|C|, (|C|-1)) = 1$ . Therefore there exists a positive integer  $\lambda$  such that  $m = \lambda|C|$  and hence  $\alpha c_S + \beta c_R = \lambda(|C|-1)$ .  $\square$

Let  $T$  be a finite chain ring with maximal ideal  $\langle \gamma \rangle$ , nilpotency index  $e$ , and residue field  $T/\langle \gamma \rangle = \mathbb{F}_{p^k}$ . A homogenous weight is defined as follows

$$\omega_{hom}(t) = \begin{cases} (p^k - 1)p^{k(e-2)}, & t \in T \setminus \langle \gamma^{e-1} \rangle; \\ p^{k(e-1)}, & t \in \langle \gamma^{e-1} \rangle \setminus \langle 0 \rangle; \\ 0, & t = 0. \end{cases}$$

**Lemma 6.5.** *With above assumptions, let  $T$  be a chain ring. Then  $\omega_{hom}$  is pre-homogeneous with average weight  $c_T = (p^k - 1)p^{k(e-2)}$ .*

*Proof.* Let  $\langle t \rangle$  be an ideal of  $T$ . By the structure of chain rings,  $\langle t \rangle = \langle \gamma^j \rangle$  for some  $j$ ;  $1 \leq j \leq e$ . Hence  $|\langle \gamma^{e-1} \rangle| = |\langle \gamma^j \rangle| = p^{k(e-j)}$ . Therefore

$$\begin{aligned} \sum_{t' \in \langle t \rangle} a_{t'} &= \sum_{t' \in \langle \gamma^j \rangle \setminus \langle \gamma^{e-1} \rangle} a_{t'} + \sum_{t' \in \langle \gamma^{e-1} \rangle} a_{t'} \\ &= (p^k - 1)p^{k(e-2)}(|\langle \gamma^j \rangle| - |\langle \gamma^{e-1} \rangle|) + p^{k(e-1)}(|\langle \gamma^{e-1} \rangle| - 1) \\ &= (p^k - 1)p^{k(e-2)}(p^{k(e-j)} - p^k) + p^{k(e-1)}(p^k - 1) \\ &= (p^k - 1)p^{k(e-2)}p^{k(e-j)} \\ &= c_T |\langle t \rangle|. \end{aligned}$$

This completes the proof.  $\square$

**Theorem 6.6.** *Let  $\omega$  be the weight function defined by  $\omega_{hom}$  over  $\mathbb{Z}_{p^r}$  and  $\mathbb{Z}_{p^s}$  on  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes. If  $C \subseteq \mathbb{Z}_{p^r}^\alpha \times \mathbb{Z}_{p^s}^\beta$  is a one weight  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive code with weight  $m$  such that there exists no zero columns in the generator matrix of  $C$ , then there exists a unique positive integer  $\lambda$  such that  $m = \lambda|C|$  and  $(p-1)p^{r-2}(\alpha + p^{s-r}\beta) = \lambda(|C| - 1)$ .*

*Proof.* By Lemma 6.5,  $c_{\mathbb{Z}_{p^r}} = (p-1)p^{r-2}$  and  $c_{\mathbb{Z}_{p^s}} = (p-1)p^{s-2}$ . Now we have the result by Theorem 6.4.  $\square$

By Example 6.2, the Lee weight over  $\mathbb{Z}_{2^r}$  and  $\mathbb{Z}_{2^s}$  is pre-homogeneous. Hence we have the following result on one weight  $\mathbb{Z}_{2^r}\mathbb{Z}_{2^s}$ -additive codes.

**Theorem 6.7.** *Let  $C \subseteq \mathbb{Z}_{2^r}^\alpha \times \mathbb{Z}_{2^s}^\beta$  be a  $\mathbb{Z}_{2^r}\mathbb{Z}_{2^s}$ -additive code. Consider the weight  $\omega$  defined by Lee weight over  $\mathbb{Z}_{2^r}$  and  $\mathbb{Z}_{2^s}$ . If  $C$  is a one weight  $\mathbb{Z}_{2^r}\mathbb{Z}_{2^s}$ -additive code with weight  $m$  such that there exists no zero columns in the generator matrix of  $C$ , then there exists a unique positive integer  $\lambda$  such that  $m = \lambda|C|$  and  $2^{r-2}(\alpha + 2^{s-r}\beta) = \lambda(|C| - 1)$ .*

*Proof.* It follows from Example 6.2 and Theorem 6.4.  $\square$

The structure of  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes is studied in [4]. If a  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive code  $C \subseteq \mathbb{Z}_2^\alpha \times \mathbb{Z}_{2^s}^\beta$  is isomorphic to an abelian structure  $\mathbb{Z}_2^{k_0+k_s} \times \mathbb{Z}_{2^s}^{k_1} \times \cdots \times \mathbb{Z}_4^{k_{s-1}}$ , then we say that  $C$  is of type  $(\alpha, \beta; k_0, k_1, k_2, \dots, k_s)$ . The following theorem gives the structure of one weight  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes which is a generalization of Theorem 3.10 in [15].

**Theorem 6.8.** *Let  $C \subseteq \mathbb{Z}_2^\alpha \times \mathbb{Z}_{2^s}^\beta$  be a one weight  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive code of type  $(\alpha, \beta; k_0, k_1, k_2, \dots, k_s)$  with weight  $m$ . Let  $k = k_0 + sk_1 + (s-1)k_2 + \cdots + k_s$ . Then there exists a positive integer  $\lambda$  such that  $m = \lambda 2^{k-1}$ , where  $\alpha$  and  $\beta$  satisfy  $\alpha + 2^{s-1}\beta = \lambda(2^k - 1)$ . Furthermore, if  $m$  is an odd integer, then  $\alpha$  is odd and  $C = \{(0_\alpha, 0_\beta), (1_\alpha, 2_\beta^{s-1})\}$ , where  $1_\alpha = (1, \dots, 1) \in \mathbb{Z}_2^\alpha$  and  $2_\beta^{s-1} = (2^{s-1}, \dots, 2^{s-1}) \in \mathbb{Z}_{2^s}^\beta$ .*

*Proof.* By Lemma 6.3,  $\sum_{c \in C} \omega(c) = |C|(\frac{\alpha}{2} + 2^{s-2}\beta) = \frac{|C|}{2}(\alpha + 2^{s-1}\beta)$ . On the other hand, the sum of the weights of all codewords is  $(|C| - 1)m$ . But  $\gcd(\frac{|C|}{2}, (|C| - 1)) = \gcd(2^{k-1}, 2^k - 1) = 1$ . Therefore there exists a positive integer  $\lambda$  such that  $m = \lambda \frac{|C|}{2} = \lambda 2^{k-1}$  and hence  $\alpha + 2^{s-1}\beta = \lambda(2^k - 1)$ .

If  $m$  is odd, then  $\lambda 2^{k-1}$  is odd. Hence  $\lambda$  is odd and  $k = 1$ . Moreover the equality  $m = \lambda = \alpha + 2^{s-1}\beta$  implies that  $\alpha$  is odd. Since  $|C| = 2$  and  $(1_\alpha, 2_\beta^{s-1})$  is the only word with weight  $\alpha + 2^{s-1}\beta$  and addition order 2, we have that  $C = \{(0_\alpha, 0_\beta), (1_\alpha, 2_\beta^{s-1})\}$ .  $\square$

## References

- [1] T. Abualrub and I. Siap, *Cyclic codes over the rings  $\mathbb{Z}_2 + u\mathbb{Z}_2$  and  $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$* , Des. Codes Cryptogr. **42** (2007), no. 3, 273–287.
- [2] T. Abualrub, I. Siap, and N. Aydin,  *$\mathbb{Z}_2\mathbb{Z}_4$ -additive cyclic codes*, IEEE Trans. Inform. Theory **60** (2014), no. 3, 1508–1514.
- [3] I. Aydogdu, T. Abualrub, and I. Siap, *On  $\mathbb{Z}_2\mathbb{Z}_2[u]$ -additive codes*, Int. J. Comput. Math. **92** (2015), no. 9, 1806–1814.
- [4] I. Aydogdu and I. Siap, *The structure of  $\mathbb{Z}_2\mathbb{Z}_{2^s}$ -additive codes: bounds on the minimum distance*, Appl. Math. Inf. Sci. **7** (2013), no. 6, 2271–2278.
- [5] ———, *On  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive codes*, Linear Multilinear Algebra **63** (2015), no. 10, 2089–2102.
- [6] J. J. Bernal, J. Borges, C. Fernández-Córdoba, and M. Villanueva, *Permutation decoding of  $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes*, Des. Codes Cryptogr. **76** (2015), no. 2, 269–277.
- [7] M. Bilal, J. Borges, S. T. Dougherty, and C. Fernández-Córdoba, *Maximum distance separable codes over  $\mathbb{Z}_4$  and  $\mathbb{Z}_2 \times \mathbb{Z}_4$* , Des. Codes Cryptogr. **61** (2011), no. 1, 31–40.
- [8] J. Borges and C. Fernández-Córdoba, *There is exactly one  $\mathbb{Z}_2\mathbb{Z}_4$ -cyclic 1-perfect code*, Des. Codes Cryptogr. **85** (2017), no. 3, 557–566.
- [9] J. Borges, C. Fernández-Córdoba, J. Pujol, J. Rifa, and M. Villanueva,  *$\mathbb{Z}_2\mathbb{Z}_4$ -linear codes: generator matrices and duality*, Des. Codes Cryptogr. **54** (2010), no. 2, 167–179.
- [10] J. Borges, C. Fernández-Córdoba, and R. Ten-Valls, *On  $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$ -additive cyclic codes*, (2016).
- [11] ———, *Linear and cyclic codes over direct product of chain rings*, Math. Meth. Appl. Sci. (2017).

- [12] H. Q. Dinh and S. R. López-Permouth, *Cyclic and negacyclic codes over finite chain rings*, IEEE Trans. Inform. Theory **50** (2004), no. 8, 1728–1744.
- [13] S. T. Dougherty and C. Fernández-Córdoba,  $\mathbb{Z}_2\mathbb{Z}_4$ -additive formally self-dual codes, Des. Codes Cryptogr. **72** (2014), no. 2, 435–453.
- [14] S. T. Dougherty, H. Liu, and Y. H. Park, *Lifted codes over finite chain rings*, Math. J. Okayama Univ. **53** (2011), 39–53.
- [15] S. T. Dougherty, H. Liu, and L. Yu, *One weight  $\mathbb{Z}_2\mathbb{Z}_4$  additive codes*, Appl. Algebra Engrg. Comm. Comput. **27** (2016), no. 2, 123–138.
- [16] C. Fernández-Córdoba, J. Pujol, and M. Villanueva,  $\mathbb{Z}_2\mathbb{Z}_4$ -linear codes: rank and kernel, Des. Codes Cryptogr. **56** (2010), no. 1, 43–59.
- [17] J. Rifà, F. I. Solov'eva, and M. Villanueva, *On the intersection of  $\mathbb{Z}_2\mathbb{Z}_4$ -additive perfect codes*, IEEE Trans. Inform. Theory **54** (2008), no. 3, 1346–1356.
- [18] K. Samei and M. R. Alimoradi, *Cyclic codes over the ring  $F_2 + uF_2 + vF_2$* , Comput. Appl. Math. **37** (2018), no. 3, 2489–2502.
- [19] K. Samei and S. Mahmoudi, *Cyclic R-additive codes*, Discrete Math. **340** (2017), no. 7, 1657–1668.
- [20] ———, *Singleton bounds for R-additive codes*, Adv. Math. Commun. **12** (2018), no. 1, 107–114.
- [21] K. Samei and S. Sadeghi, *Maximum distance separable codes over  $\mathbb{Z}_2 \times \mathbb{Z}_{2^s}$* , J. Algebra Appl. **17** (2018), no. 7, 1850136, 12 pp.
- [22] B. Srinivasulu and M. Bhaintwal,  $\mathbb{Z}_2(\mathbb{Z}_2 + u\mathbb{Z}_2)$ -additive cyclic codes and their duals, Discrete Math. Algorithms Appl. **8** (2016), no. 2, 1650027, 19 pp.
- [23] J. A. Wood, *The structure of linear codes of constant weight*, Trans. Amer. Math. Soc. **354** (2002), no. 3, 1007–1026.

SAADOUN MAHMOUDI  
DEPARTMENT OF MATHEMATICS  
BU ALI SINA UNIVERSITY  
HAMEDAN, IRAN  
Email address: mahmoudi.math89@yahoo.com

KARIM SAMEI  
DEPARTMENT OF MATHEMATICS  
BU ALI SINA UNIVERSITY  
HAMEDAN, IRAN  
Email address: samei@ipm.ir