

## SEIBERG-WITTEN-LIKE EQUATIONS ON THE STRICTLY PSEUDOCONVEX $CR$ -3 MANIFOLDS

SERHAN EKER

ABSTRACT. In this paper, Seiberg-Witten-like equations are written down on 3-manifolds. Then, it has been proved that the  $L^2$ -solutions of these equations are trivial on  $\mathbb{R}^3$ . Finally, a global solution is obtained on the strictly pseudoconvex  $CR$ -3 manifolds for a given constant negative scalar curvature.

### 1. Introduction

The Seiberg-Witten equations, which are consisted of the curvature and Dirac equation, carries subtle information that can be used to investigate the topology and geometry of the manifolds [11, 21, 27]. Although these equations are defined in 4-manifolds, they are also handled by many authors on higher dimensional manifolds [4–6, 18, 21]. This paper is mainly interested with the Seiberg-Witten-like equations on 3-manifold. Seiberg-Witten-like equations on 3-manifold are studied to obtain the equations of motion of  $U(1)$  Chern-Simons theory coupled to a massless spinor field and investigated their moduli space of the gauge equivalence classes of their solutions [3]. Also, these equations are studied on a compact 3-manifold with boundary to show that solution space of these equations is a Banach manifold [16]. As mentioned above, Seiberg-Witten-like equations have been investigated by many authors in three dimensions [11, 14, 17, 23, 25]. In this paper, these equations are investigated with a different perspective. Just as in the 4-manifolds, there is a need for a  $\text{Spin}^c$ -structure in order to be able to construct spinor bundle on 3-manifolds. With all these, the Dirac equation can be defined on the spinor bundle. However, the definition of the curvature equation differs from the curvature equation which are defined on 4-manifolds. On 4-manifolds the curvature equation is defined as being dependent on the self-duality concept and also in higher dimensions the curvature equation is defined with the help of the generalised self-duality

---

Received December 28, 2018; Revised May 11, 2019; Accepted August 5, 2019.

2010 *Mathematics Subject Classification*. Primary 53C27, 15A66, 34L40.

*Key words and phrases*. Spin and  $\text{Spin}^c$  geometry, Clifford algebras, spinors, Seiberg-Witten equations.

concept [4–6]. Since in 3-manifolds self-duality concept is meaningless, the curvature equation is defined independently from the self-duality concept.

This paper is organized as follows. At first, some basic facts concerning contact metric manifold,  $\text{Spin}^c$ -structure and Dirac operator are introduced. Then, in Section 3, Seiberg-Witten-like equations on 3-manifolds are defined and some useful identities are obtained to determine Seiberg-Witten-like functional. Therefore, a bound to the solutions of these equations with the approach given in [15] is obtained. Furthermore, it is proved that  $L^2$ -solutions of these equations are trivial on  $\mathbb{R}^3$ . Finally, a global solution to these equations on the strictly pseudoconvex  $CR$ -3 manifolds is obtained for a given constant negative scalar curvature.

## 2. Some basic materials

### 2.1. Contact metric manifolds

Let  $M$  be a smooth 3-manifold. A smooth 1-form  $\alpha$  on  $M$  is called a contact form if  $\alpha \wedge (d\alpha) \neq 0$  everywhere on  $M$ . A hyperplane subbundle  $H$  of the tangent bundle  $TM$ , which is given by  $H = \ker \alpha$ , is induced by contact form  $\alpha$ . The Reeb vector field  $\xi$  is the unique vector field satisfying  $\alpha(\xi) = 1$  and  $d\alpha(\xi, \cdot) = 0$ . Then  $(M, \alpha)$  is called a contact manifold. The tangent bundle  $TM$  splits into  $TM = H \oplus \mathbb{R}\xi$ . Let  $X$  be any vector field on  $M$ . Then, the decomposition of  $X$  can be written as

$$X = X_H + f\xi,$$

where  $f \in C^\infty(M, \mathbb{R})$  and  $X_H$  is the horizontal part of  $X$ .

If  $(M, \alpha)$  is a contact manifold, the pair  $(H, d\alpha|_H)$  is a symplectic vector bundle and an almost complex structure  $J_H$  can be fixed on  $H$ , which is compatible with  $d\alpha|_H$ . Since  $J_H^2 = -I_d$ , the following eigenspaces decomposition can be given by:

$$\Lambda_H^1(M) = H \otimes_{\mathbb{R}} \mathbb{C} = \Lambda_H^{1,0}(M) \oplus \Lambda_H^{0,1}(M),$$

where

$$\Lambda_H^{1,0}(M) = \{Z \in H \otimes_{\mathbb{R}} \mathbb{C} \mid J_H Z = iZ\},$$

$$\Lambda_H^{0,1}(M) = \{Z \in H \otimes_{\mathbb{R}} \mathbb{C} \mid J_H Z = -iZ\}.$$

The complexification of  $\Lambda_H^s(M)$  is decomposed as follows

$$\Lambda_H^s(M) = \sum_{q+r=s} \Lambda_H^{q,r}(M),$$

where  $\Lambda^{q,r}(M)_H = \text{span}\{u \wedge v \mid u \in \Lambda^q(\Lambda_H^{1,0}(M)), v \in \Lambda^r(\Lambda_H^{0,1}(M))\}$ . Also,  $J_H$  can be extended to an endomorphism  $J$  of the tangent bundle  $TM$  by setting  $J\xi = 0$ . At this point  $J^2 = -Id + \alpha \otimes \xi$  is satisfied. With this in mind,  $g_\alpha$  defines a Riemannian metric on  $TM$  given by

$$g_\alpha(X, Y) = d\alpha(X, JY) + \alpha(X)\alpha(Y).$$

The metric  $g_\alpha$  is called a Webster metric and is said to be associated to  $\alpha$ . Moreover,  $g_\alpha$  satisfies the following relations:

$$\begin{aligned} g_\alpha(X, Y) &= \alpha(X), \quad g_\alpha(JX, Y) = d\alpha(X, Y), \\ g_\alpha(JX, JY) &= g_\alpha(X, Y) - \alpha(X)\alpha(Y), \end{aligned}$$

for any  $X, Y \in \chi(M)$ . We call  $(M, g_\alpha, \alpha, \xi, J)$  as a contact metric manifold. For more information see [1, 2, 20].

On the contact metric manifold  $(M, g_\alpha, \alpha, \xi, J)$ , the generalized Tanaka-Webster connection  $\nabla^{TW}$  is given by:

$$\nabla_X^{TW} Y = \nabla_X Y - (\nabla_X \alpha)(Y)\xi - \alpha(X)\nabla_Y \xi - \alpha(X)\alpha(Y),$$

where  $\nabla$  is the Levi-Civita connection and  $X, Y \in \chi(M)$  [24]. Also, the generalized Tanaka-Webster connection  $\nabla^{TW}$  satisfies  $\nabla^{TW} \alpha = 0$  and  $\nabla^{TW} g_\alpha = 0$ . Moreover, if  $J$  is integrable, i.e.,  $\nabla^{TW} J = 0$ , then the contact metric manifold  $(M, g_\alpha, \alpha, \xi, J)$  is called a strictly pseudoconvex  $CR$  manifold [19, 20].

## 2.2. Spin<sup>c</sup>-structure and Dirac operator

A complex vector bundle  $\mathbb{S}$  can be constructed by a given Spin<sup>c</sup> representation  $\kappa_3 : \text{Spin}^c(3) \rightarrow \text{Aut}(\Delta_3)$  and denoted by  $\mathbb{S} = P_{\text{Spin}^c(3)} \times_{\kappa_3} \Delta_3$ . Also this complex vector bundle is called a spinor bundle for a given Spin<sup>c</sup>-structure on  $M$  [8].  $\kappa : \mathbb{R}^3 \rightarrow \text{End}(\mathbb{S})$  is a linear map satisfying the following conditions:

$$\kappa(v)^* + \kappa(v) = 0, \quad \kappa(v)^* \kappa(v) = |v|^2 \mathbb{I}$$

for every  $v \in \mathbb{R}^3$ . Then,

$$\rho : \Lambda^2(T^*M) \rightarrow \text{End}(\mathbb{S})$$

can be defined on the frames by extending map  $\kappa : TM \rightarrow \text{End}(\mathbb{S})$  of  $\kappa$ . Let  $\{e_1, e_2, e_3\}$  be orthonormal frame on open subset  $U \subset M$ . Then

$$\alpha = \sum_{i < j} \alpha_{ij} e^i \wedge e^j \rightarrow \rho(\alpha) = \sum_{i < j} \alpha_{ij} \kappa(e_i) \kappa(e_j).$$

$\rho$  can be extended to complex valued 2-forms such that

$$\rho : \Lambda^2(T^*M) \otimes \mathbb{C} \rightarrow \text{End}(\mathbb{S}).$$

A connection  $\nabla^A$  on  $\mathbb{S}$ , which is called a spinor covariant derivative operator, is obtained by using an  $i\mathbb{R}$ -valued connection 1-form  $A \in \Omega(M, i\mathbb{R})$  and the Levi-Civita connection  $\nabla$  on  $M$ . At this point the definition of the Dirac operator  $D_A : \Gamma(\mathbb{S}) \rightarrow \Gamma(\mathbb{S})$  can be given by

$$D_A(\Psi) = \sum_{i=1}^3 \kappa(e_i) \nabla_{e_i}^A \Psi,$$

where  $\{e_1, e_2, e_3\}$  is any positively oriented local orthonormal frame of  $TM$  [13]. A Spin<sup>c</sup>-structure is needed to describe the Dirac operator on contact metric manifold. It is known that any contact metric manifold admits a canonical Spin<sup>c</sup>-structure. By using this canonical Spin<sup>c</sup>-structure, an associated

canonical spinor bundle can be constructed and described by the following isomorphism:

$$\mathbb{S} \cong \Omega^{0,*}(M),$$

where  $\Omega^{0,*}(M)$  is the direct sum of  $\Omega(M)^{0,0} \oplus \Omega(M)^{0,1}$ . Furthermore, on this spinor bundle, the Clifford multiplication “ $\cdot$ ” is given by:

$$V \cdot \Psi = \sqrt{2} \left( (V_H^{0,1})^* \wedge \Psi - \iota(V_H^{0,1})\Psi \right) + i(-1)^{\deg \Psi + 1} \eta(V)\psi,$$

where  $V_H$  denotes the horizontal part of  $V$ . According to these multiplication one can easily obtain  $\xi\psi = i(-1)^{\deg \psi + 1}\psi$ .

The spinor bundle  $\mathbb{S}$  carries a natural Hermitian metric, denoted by  $(\cdot, \cdot)$  and for any vector field  $X$  and spinor field  $\Psi, \Phi$  satisfies [12]

$$(1) \quad (X \cdot \Psi, \Phi) = -(\Psi, X \cdot \Phi).$$

Also, the norm  $\|\cdot\|$  in the Hilbert space  $L^2$  is defined as [8, 21],

$$(2) \quad \|\Psi\|^2 = \sqrt{\int_M |\Psi|^2 dvol}.$$

On the  $2n+1$ -dimensional contact metric manifold  $(M, g_\alpha, \alpha, \xi, J)$  equipped with a  $\text{Spin}^c$ -structure, each unitary connection  $A$  on  $L$  induces a spinorial connection  $\nabla^A$  on  $\mathbb{S}$  with the generalized Tanaka-Webster connection  $\nabla^{TW}$ . Then according to  $\nabla^A$  the Kohn-Dirac operator  $D_H^A$  is defined as follows [20]:

$$(3) \quad D_H^A = \sum_{i=1}^{2n} \kappa(e_i)(\nabla_{e_i}^A),$$

where  $\{e_i\}$  is a local orthonormal frame of  $H$ . The Dirac operator  $D_A$  is defined by [20]

$$(4) \quad D_A = D_H^A + \xi \cdot \nabla_\xi^A.$$

Also, by considering strictly pseudoconvex CR manifolds with  $\Omega_H^{0,*}(M)$  associated spinor bundle the Dirac type operator is defined as follows:

Let

$$(5) \quad \bar{\partial}_H : \Omega_H^{0,r}(M) \longrightarrow \Omega_H^{0,r+1}(M), \quad \bar{\partial}_H^* : \Omega_H^{0,r}(M) \longrightarrow \Omega_H^{0,r-1}$$

respectively given by:

$$\bar{\partial}_H = \sum_{i=1}^n \bar{Z}_i^* \wedge \nabla_{\bar{Z}_i}^{TW}, \quad \bar{\partial}_H^* = - \sum_{i=1}^n \iota(\bar{Z}_i)^* \wedge \nabla_{\bar{Z}_i}^{TW},$$

where  $\nabla^{TW}$  is the extension of the generalized Webster-Tanaka connection to  $\Omega_H^{0,*}(M)$  and  $\iota$  is the contraction operator.

It follows from (3) that we have on  $\Omega_H^{0,*}(M)$

$$(6) \quad \mathcal{H} = \sqrt{2} \sum_{r=0}^n (\bar{\partial}_H + \bar{\partial}_H^*) + \sum_{r=0}^n (-1)^{r+1} \sqrt{-1} \cdot \nabla_\xi^{TW}.$$

Since  $\mathbb{S} \cong \Omega_H^{0,*}(M)$ , (4) coincides with (6).

### 3. Seiberg-Witten-like equations on 3-manifolds

In this section, we write down Seiberg-Witten-like equation on 3-manifold  $M$ . Then, we get explicit forms of these equations on  $\mathbb{R}^3$ .

**Definition.** Let  $M$  be a 3-manifold endowed with a  $\text{Spin}^c(3)$ -structure and  $A$  be the fixed connection on  $U(1)$ -principal bundle. Then, for any  $\Psi \in \Gamma(\mathbb{S})$  Seiberg-Witten-like equations are defined by

$$\begin{aligned} D_A(\Psi) &= 0, \\ F_A &= \frac{1}{4}\sigma(\Psi), \end{aligned}$$

where  $F_A = dA$  is the imaginary-valued curvature 2-form of the connection  $A$  in the  $P_{\mathbb{S}^1}$ -bundle associated with the  $\text{Spin}^c$ -structure.

Moreover, the well known formula called the Schrödinger-Lichnerowicz formula is given by [8, 21]

$$(7) \quad D_A^* D_A \Psi = (\nabla^A)^* \nabla^A \Psi + \frac{s}{4} \Psi + \frac{1}{2} d_A \cdot \Psi,$$

where  $s$  is the scalar curvature of  $M$ ,  $(\nabla^A)^*$  is the adjoint of the covariant derivative operator  $\nabla^A$  and  $D_A^*$  is the adjoint of  $D_A$ .

#### 3.1. Seiberg-Witten-like equations on $\mathbb{R}^3$

Let  $\kappa : \mathbb{R}^3 \rightarrow \text{End}(\mathbb{C}^2)$  be the  $\text{Spin}^c(3)$ -structure which is defined on generators  $\{e_1, e_2, e_3\}$  by the followings:

$$\begin{aligned} \kappa(e_1) &= \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad \kappa(e_2) = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \\ \kappa(e_3) &= \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \quad \kappa(d\alpha) = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, \end{aligned}$$

where  $d\alpha = e^1 \wedge e^2$ . Note that there is no decomposition of spinor space over  $\mathbb{R}^3$  contrary to the case  $\mathbb{R}^4$  [8]. The  $\text{Spin}^c$ -connection  $\nabla^A$  on  $\mathbb{R}^3$  is given by

$$\nabla_j^A \Psi = \frac{\partial \Psi}{\partial x_j} + \frac{1}{2} A_j \Psi,$$

where  $A_j : \mathbb{R}^3 \rightarrow i\mathbb{R}$  for  $j = 1, 2, 3$  and  $\Psi : \mathbb{R}^3 \rightarrow \mathbb{C}^2$  are smooth maps. Then the associated connection on the line bundle  $P_{\mathbb{S}^1}$  is the connection 1-form and represented by

$$A = \sum_{i=1}^3 A_i dx^i \in \Omega(\mathbb{R}^3, i\mathbb{R})$$

and its curvature 2-form is given by

$$F_A = \sum_{i < j}^3 F_{ij} dx^i \wedge dx^j \in \Omega^2(\mathbb{R}^3, i\mathbb{R})$$

where  $F_{ij} = \left( \frac{\partial A_j}{\partial x_i} - \frac{\partial A_i}{\partial x_j} \right)$  for  $i, j = 1, 2, 3$ . Then the Dirac operator  $D_A : \Gamma(\mathbb{S}) \rightarrow \Gamma(\mathbb{S})$  on  $\mathbb{R}^3$  can be written with respect to given  $\text{Spin}^c$ -connection  $\nabla^A$  as follows:

$$D_A \Psi = \sum_{i=1}^3 \kappa(e_i) \nabla_{e_i}^A \Psi.$$

Therefore, the Dirac equation in the flat case is given by

$$\begin{aligned} D_A(\Psi) &= \kappa(e_1) \nabla_{e_1}^A \Psi + \kappa(e_2) \nabla_{e_2}^A \Psi + \kappa(e_3) \nabla_{e_3}^A \Psi \\ &= \sum_{i=1}^3 \kappa(e_i) \left( \nabla_{e_i}^A \Psi \right) \\ &= \sum_{i=1}^3 \kappa(e_i) \begin{bmatrix} \frac{\partial \psi_1}{\partial x_i} + \frac{1}{2} A_i \psi_1 \\ \frac{\partial \psi_2}{\partial x_i} + \frac{1}{2} A_i \psi_2 \end{bmatrix} \\ &= \begin{bmatrix} i \left( \frac{\partial \psi_2}{\partial x_2} + \frac{\partial \psi_1}{\partial x_3} \right) + \frac{\partial \psi_2}{\partial x_1} + \frac{1}{2} \left( A_1 \psi_2 + i(A_3 \psi_1 + A_2 \psi_2) \right) \\ i \left( -\frac{\partial \psi_2}{\partial x_3} + \frac{\partial \psi_1}{\partial x_2} \right) - \frac{\partial \psi_1}{\partial x_1} + \frac{1}{2} \left( -A_1 \psi_1 + i(A_2 \psi_1 - A_3 \psi_2) \right) \end{bmatrix}. \end{aligned}$$

Let us consider the complexified space  $\Lambda^2(\mathbb{R}^3) \otimes \mathbb{C}$  and  $F_A$  be the curvature form of the imaginary valued connection 1-form  $A$ . Then,

$$F_A = \sum_{i < j}^3 F_{ij} dx^i \wedge dx^j.$$

The curvature equation is defined by

$$F_A = \frac{1}{4} \sigma(\Psi),$$

where  $\sigma(\Psi)$  is an imaginary valued 2-form defined by the formula

$$\sigma(\Psi)(X, Y) = \langle X \cdot Y \cdot \Psi, \Psi \rangle + \langle X, Y \rangle |\Psi|^2$$

for any  $\Psi \in \Gamma(\mathbb{S})$ . The map  $\sigma : \Gamma(\mathbb{S}) \rightarrow \Omega^1(M, i\mathbb{R})$  is called a quadratic map.

The explicit form of the second equation can be expressed as follows:

$$\begin{aligned} F_{12} &= -\frac{i}{4} \left( |\psi_1|^2 - |\psi_2|^2 \right), \\ F_{13} &= \frac{i}{4} \left( \overline{\psi_2} \psi_1 + \overline{\psi_1} \psi_2 \right), \\ F_{23} &= \frac{1}{4} \left( \overline{\psi_2} \psi_1 - \overline{\psi_1} \psi_2 \right). \end{aligned}$$

#### 4. Seiberg-Witten-like functional

The Energy functional consistent with the 3-dimensional Seiberg-Witten-like equations is defined by

$$E(A, \Psi) = \int_M \left( |D_A \Psi|^2 + |F_A - \frac{1}{4} \sigma(\Psi)|^2 \right) dvol.$$

Note that solutions of 3-dimensional Seiberg-Witten-like equations are zeros of Energy functional. In this section, we obtain some useful identities related with spinors and their image under the quadratic map  $\sigma$ . With the aid of the following lemma we obtain another form of Seiberg-Witten-like functional and we get a bound for the solutions of Seiberg-Witten-like equations.

**Lemma 4.1.** *Let  $\kappa : TM \rightarrow \text{End}(\mathbb{S})$  be a  $\text{Spin}^c$ -structure on a compact oriented smooth 3-dimensional Riemannian manifold  $M$ . Then, the following equalities hold*

$$(8) \quad (\sigma(\Psi)\Psi, \Psi) = (\sigma(\Psi), \sigma(\Psi)) = |\Psi|^4,$$

where  $\Psi \in \Gamma(\mathbb{S})$  and  $\sigma(\Psi) \in \Omega^2(M, i\mathbb{R})$ .

Lemma 4.1 can be proved with an easy computation.

In the following, we give a bound to the negative constant scalar curvature of  $(M, g)$  by using the usual Laplacian on defined as follows [8, 21],

$$(9) \quad \Delta|\Psi|^2 = 2((\nabla^A)^* \nabla^A \Psi, \Psi) - 2(\nabla^A \Psi, \nabla^A \Psi),$$

where  $\Psi \in \Gamma(\mathbb{S})$  and  $(\nabla^A)^*$  is the adjoint of the covariant derivative operator  $\nabla^A$ .

**Lemma 4.2.** *Let  $(A, \Psi)$  be a solution of  $D_A \Psi = 0$ ,  $F_A = \frac{1}{4} \sigma(\Psi)$  over a compact smooth 3-dimensional Riemannian manifold  $(M, g)$  with a negative constant scalar curvature  $s$ . Then, at each point*

$$\frac{|\Psi(x)|^2}{2} \leq -s_{\min}, \quad \text{where } s_{\min} = \min\{s(m) : m \in M\}.$$

*Proof.* At a point  $x$  where  $|\Psi(x)|^2$  attains its maximum we have  $0 \leq \Delta|\Psi|^2$ . Then

$$\begin{aligned} 0 \leq \Delta|\Psi|^2 &= 2((\nabla^A)^* \nabla^A \Psi, \Psi) - 2(\nabla^A \Psi, \nabla^A \Psi) \\ &\leq 2((\nabla^A)^* \nabla^A \Psi, \Psi) \\ &= 2(D_A^* D_A \Psi - \frac{s}{4} \Psi - \frac{1}{2} dA \cdot \Psi, \Psi) \\ &= \left( -\frac{s}{2} \Psi - dA \cdot \Psi, \Psi \right), \\ &= -\frac{s}{2} |\Psi|^2 - (dA \cdot \Psi, \Psi) \\ &= -\frac{s}{2} |\Psi|^2 - \frac{1}{4} (\sigma(\Psi)\Psi, \Psi) \end{aligned}$$

$$= -\frac{s}{2}|\Psi|^2 - \frac{1}{4}|\Psi|^4.$$

Now, if  $|\Psi(x)|^2 > 0$ , then  $0 \leq -\frac{s}{2}|\Psi|^2 - \frac{1}{4}|\Psi|^4_{\max}$  and  $\frac{1}{2}|\Psi(x_{\max})|^2 \leq -s_{\min}$ .  $\square$

**Lemma 4.3.** *Under the same conditions as in Lemma 4.2, the following inequality is satisfied*

$$|F_A| \leq \frac{1}{2}|s|.$$

*Proof.*

$$\begin{aligned} |F_A|^2 &= \left| \frac{1}{4}\sigma(\Psi) \right|^2 = \left( \frac{1}{4}\sigma(\Psi), \frac{1}{4}\sigma(\Psi) \right) \\ &= \frac{1}{16}(\sigma(\Psi), \sigma(\Psi)) \\ &= \frac{1}{16}|\Psi|^4. \end{aligned}$$

As a result,  $|F_A| = \frac{1}{4}|\Psi|^2 \leq \frac{-s}{2} \leq \frac{1}{2}|s|$ .  $\square$

Since 3-dimensional Hyperbolic space is a Riemannian manifold with negative constant curvature and it satisfies Lemma 4.2 and Lemma 4.3 [26]. In addition, manifolds of negative constant curvature are given in [10].

**Lemma 4.4.** *On the compact oriented smooth 3-dimensional Riemannian manifold  $(M, g)$ , by considering Seiberg-Witten-like equation:*

$$(10) \quad D_A \Psi = 0, \quad F_A = \frac{1}{4}\sigma(\Psi),$$

*the Seiberg-Witten-like functional is obtained as follows*

$$E(A, \Psi) = \int_M \left( |F_A|^2 + |\nabla^A \Psi|^2 + \frac{s}{4}|\Psi|^2 + \frac{1}{16}|\Psi|^4 \right) dvol.$$

*Proof.* Using the Schrodinger-Lichnerowicz formula given in (7), we have

$$(11) \quad \int_M |D_A \Psi|^2 dvol = \int_M \left[ |\nabla^A \Psi|^2 + \frac{s}{4}|\Psi|^2 + \left( \frac{1}{2}dA \cdot \Psi, \Psi \right) \right] dvol.$$

Since  $F_A$  and  $\sigma(\Psi)$  are 2-forms with purely imaginary values, calculating their length amounts to

$$\left| F_A - \frac{1}{4}\sigma(\Psi) \right|^2 = |F_A|^2 - \frac{1}{2}(dA \cdot \Psi, \Psi) + \frac{1}{16}|\sigma(\Psi)|^2.$$

This implies

$$\begin{aligned} E(A, \Psi) &= \int_M \left[ \left| F_A - \frac{1}{4}\sigma(\Psi) \right|^2 + |D_A \Psi|^2 \right] dvol \\ (12) \quad &= \int_M \left[ |F_A|^2 + |\nabla^A \Psi|^2 + \frac{s}{4}|\Psi|^2 + \frac{1}{16}|\Psi|^4 \right] dvol. \end{aligned} \quad \square$$

In 3-dimensional case i.e.,  $M = \mathbb{R}^3$ , the following lemma shows that  $L^2$ -solutions of these equations are trivial.



**Lemma 4.5.** *Let  $A \in \Omega^1(\mathbb{R}^3, i\mathbb{R})$  and  $\Psi \in C^\infty(\mathbb{R}^3, \mathbb{C}^2)$  the following equations are satisfied:*

$$(13) \quad \begin{aligned} \nabla_1^A \Psi + \nabla_2^A \Psi + \nabla_3^A \Psi &= 0, \\ F_{12} &= -\frac{i}{4}(|\psi_1|^2 - |\psi_2|^2) = -\frac{1}{4}\Psi^* K \Psi, \\ F_{13} &= \frac{i}{4}(\overline{\psi_2}\psi_1 + \overline{\psi_1}\psi_2) = \frac{1}{4}\Psi^* J \Psi, \\ F_{23} &= \frac{1}{4}(\overline{\psi_2}\psi_1 - \overline{\psi_1}\psi_2) = -\frac{1}{4}\Psi^* I \Psi, \end{aligned}$$

where

$$I = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad J = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, \quad K = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}.$$

Then

- (1) If  $\Psi \in L^2$ , then  $\Psi \equiv 0$ .
- (2) If  $E(A, \Psi) < \infty$ , then  $\Psi \equiv 0$  and  $F_A \equiv 0$ .

*Proof.* Let

$$(14) \quad \Delta = -\sum_{i=1}^3 \frac{\partial^2}{(\partial x^i)^2}$$

be the usual Laplacian on  $\mathbb{R}^3$ . At first we claim that

$$(15) \quad \Delta|\Psi|^2 = -2 \sum_{i=1}^3 \frac{\partial}{\partial x^i} \operatorname{Re}(\Psi, \nabla_i^A \Psi).$$

To proof our claim we compute

$$(16) \quad \begin{aligned} \frac{\partial}{\partial x^i} |\Psi|^2 &= \partial_i(\overline{\psi_1}\psi_1 + \overline{\psi_2}\psi_2) \\ &= \overline{\psi_1}\partial_i\psi_1 + \psi_1\partial_i\overline{\psi_1} + \overline{\psi_2}\partial_i\psi_2 + \psi_2\partial_i\overline{\psi_2}, \end{aligned}$$

and

$$(17) \quad \begin{aligned} (\Psi, \nabla_i^A \Psi) &= (\Psi, \partial_i \Psi + \frac{1}{2}A_i \Psi) \\ &= \overline{\psi_1}(\partial_i \psi_1 + \frac{1}{2}A_i \psi_1) + \overline{\psi_2}(\partial_i \psi_2 + \frac{1}{2}A_i \psi_2) \\ &= \overline{\psi_1}\partial_i\psi_1 + \overline{\psi_2}\partial_i\psi_2 + \frac{1}{2}A_i(|\psi_1|^2 + |\psi_2|^2). \end{aligned}$$

By inserting (17) into the following equality

$$(18) \quad \begin{aligned} 2\operatorname{Re}(\Psi, \nabla_i^A \Psi) &= (\Psi, \nabla_i^A \Psi) + \overline{(\Psi, \nabla_i^A \Psi)} \\ &= \overline{\psi_1}\partial_i\psi_1 + \psi_1\partial_i\overline{\psi_1} + \overline{\psi_2}\partial_i\psi_2 + \psi_2\partial_i\overline{\psi_2} \end{aligned}$$

is obtained. At the end, by comparing (16) with (18), one gets the following equality:

$$\frac{\partial}{\partial x^i} |\Psi|^2 = 2Re(\Psi, \nabla_i^A \Psi).$$

Also, this equality can be written as in the following

$$-\frac{\partial^2}{(\partial x^i)^2} |\Psi|^2 = -2 \frac{\partial}{\partial x^i} Re(\Psi, \nabla_i^A \Psi),$$

which means that (14) equals to (15).

Moreover, one can show that

$$(19) \quad \frac{\partial}{\partial x^i} \left( Re(\Psi, \nabla_i^A \Psi) \right) = |\nabla_i^A \Psi|^2 + Re(\Psi, \nabla_i^A \nabla_i^A \Psi).$$

To obtain this, firstly, the right side of (19) is computed:

$$(20) \quad \begin{aligned} 2 \frac{\partial}{\partial x^i} Re(\Psi, \nabla_i^A \Psi) &= \frac{\partial}{\partial x^i} (\overline{\psi_1} \partial_i \psi_1 + \psi_1 \partial_i \overline{\psi_1} + \overline{\psi_2} \partial_i \psi_2 + \psi_2 \partial_i \overline{\psi_2}) \\ &= \overline{\psi_1} \partial_i \partial_i \psi_1 + \partial_i \overline{\psi_1} \partial_i \psi_1 + \psi_1 \partial_i \partial_i \overline{\psi_1} + \partial_i \psi_1 \partial_i \overline{\psi_1} \\ &\quad + \overline{\psi_2} \partial_i \partial_i \psi_2 + \partial_i \overline{\psi_2} \partial_i \psi_2 + \psi_2 \partial_i \partial_i \overline{\psi_2} + \partial_i \psi_2 \partial_i \overline{\psi_2} \\ &= \overline{\psi_1} \partial_i \partial_i \psi_1 + \psi_1 \partial_i \partial_i \overline{\psi_1} + \overline{\psi_2} \partial_i \partial_i \psi_2 + \psi_2 \partial_i \partial_i \overline{\psi_2} \\ &\quad + \partial_i \psi_1 \partial_i \overline{\psi_1} + \partial_i \psi_2 \partial_i \overline{\psi_2}. \end{aligned}$$

Explicit form of  $|\nabla_i^A \Psi|^2$  is obtained as in the following:

$$(21) \quad \begin{aligned} 2|\nabla_i^A \Psi|^2 &= 2(\nabla_i^A \Psi, \nabla_i^A \Psi) \\ &= 2(\partial_i \Psi, \partial_i \Psi) + (\partial_i \Psi, A_i \Psi) + (A_i \Psi, \partial_i \Psi) + \frac{1}{2}(A_i \Psi, A_i \Psi) \\ &= 2\partial_i \overline{\psi_1} \partial_i \psi_1 + 2\partial_i \overline{\psi_2} \partial_i \psi_2 + 2Re(\partial_i \Psi, A_i \Psi) + \frac{1}{2}|A_i \Psi|^2. \end{aligned}$$

Also,

$$(22) \quad \begin{aligned} \nabla_i^A \nabla_i^A \Psi &= (\partial_i + \frac{1}{2} A_i)(\partial_i \Psi + \frac{1}{2} A_i \Psi) \\ &= \partial_i \partial_i \Psi + \frac{1}{2} \partial_i (A_i \Psi) + \frac{1}{2} A_i \partial_i \Psi + \frac{1}{4} A_i^2 \Psi \\ &= \partial_i \partial_i \Psi + \frac{1}{2} A_i \partial_i \Psi + \frac{1}{2} \Psi \partial_i A_i + \frac{1}{2} A_i \partial_i \Psi + \frac{1}{4} A_i^2 \Psi \\ &= \partial_i \partial_i \Psi + A_i \partial_i \Psi + \frac{1}{2} \Psi \partial_i A_i + \frac{1}{4} A_i^2 \Psi \\ &= \partial_i \partial_i \Psi + A_i \partial_i \Psi + \frac{1}{2} \Psi \partial_i A_i - \frac{1}{4} |A_i|^2 \Psi. \end{aligned}$$

Hermitian inner product  $\Psi$  with (22) is calculated by

$$(\Psi, \nabla_i^A \nabla_i^A \Psi) = (\Psi, \partial_i \partial_i \Psi) + A_i (\Psi, \partial_i \Psi) + \frac{1}{2} \partial_i A_i (\Psi, \Psi) - \frac{1}{4} |A_i|^2 (\Psi, \Psi)$$

$$(23) \quad = \overline{\psi_1} \partial_i \partial_i \Psi_1 + \overline{\psi_2} \partial_i \partial_i \psi_2 + A_i(\Psi, \partial_i \Psi) + \frac{1}{2} \partial_i A_i |\Psi|^2 - \frac{1}{4} |A_i \Psi|^2.$$

The real part of (23) is

$$(24) \quad \begin{aligned} 2Re(\Psi, \nabla_i^A \nabla_i^A \Psi) &= (\Psi, \nabla_i^A \nabla_i^A \Psi) + \overline{(\Psi, \nabla_i \nabla_i \Psi)} \\ &= \overline{\psi_1} \partial_i \partial_i \Psi_1 + \overline{\psi_2} \partial_i \partial_i \psi_2 + A_i(\Psi, \partial_i \Psi) + \frac{1}{2} \partial_i A_i |\Psi|^2 \\ &\quad - \frac{1}{4} |A_i \Psi|^2 + \psi_1 \partial_i \partial_i \overline{\Psi_1} + \psi_2 \partial_i \partial_i \overline{\psi_2} - A_i(\overline{\Psi}, \partial_i \overline{\Psi}) \\ &\quad - \frac{1}{2} \partial_i A_i |\Psi|^2 - \frac{1}{4} |A_i \Psi|^2. \end{aligned}$$

Since

$$(25) \quad \begin{aligned} A_i(\Psi, \partial_i \Psi) &= (\overline{A_i \Psi}, \partial_i \Psi) = (-A_i \Psi, \partial_i \Psi) \\ &= -(A_i \Psi, \partial_i \Psi) \\ &= -(\partial_i \Psi, A_i \Psi), \end{aligned}$$

$$(26) \quad \begin{aligned} Re(A_i(\Psi, \partial_i \Psi)) &= -Re(\partial_i \Psi, A_i \Psi) \\ &= -Re(\partial_i \Psi, A_i \Psi) \end{aligned}$$

are obtained. Inserting (26) in (24), one has

$$\begin{aligned} 2Re(\Psi, \nabla_i \nabla_i \Psi) &= \overline{\psi_1} \partial_i \partial_i \psi_1 + \psi_1 \partial_i \partial_i \overline{\psi_1} + \overline{\psi_2} \partial_i \partial_i \psi_2 + \psi_2 \partial_i \partial_i \overline{\psi_2} \\ &\quad - 2Re(\partial_i \Psi, A_i \Psi) - \frac{1}{2} |A_i \Psi|^2. \end{aligned}$$

Since (20) is the sum of (21) and (24), (19) is proved.

Considering the scalar curvature  $s = 0$  and Dirac equation  $D_A \Psi = 0$  in (7), we get

$$(\nabla_i^A)^* \nabla_i^A \Psi + \frac{1}{2} dA \cdot \Psi = 0.$$

Since  $(\nabla^A)^* = -\nabla^A$  [8, 21], we obtain

$$(27) \quad \nabla_i^A \nabla_i^A \Psi = \frac{1}{2} F_A \cdot \Psi.$$

Inserting (27) in (19), we get the following equation:

$$(28) \quad \begin{aligned} \Delta |\Psi|^2 &= -2 \sum_{i=1}^3 |\nabla_i^A \Psi|^2 + Re(\Psi, \rho(F_A) \Psi) \\ &= -2 \sum_{i=1}^3 |\nabla_i^A \Psi|^2 - 2Re(\Psi, F_{12} K \Psi) - Re(\Psi, F_{13} J \Psi) - Re(\Psi, F_{23} I \Psi). \end{aligned}$$

By using (13) in the following Hermitian inner product

$$\begin{aligned}
(1) \quad (\Psi, F_{12}K\Psi) &= \left( \Psi, \left( -\frac{1}{4}\Psi^*K\Psi \right) K\Psi \right) \\
&= \left( -\frac{1}{4}\Psi^*K\Psi \right) (\Psi, K\Psi) \\
&= -\frac{i}{4}(|\psi_1|^2 - |\psi_2|^2)i(|\psi_1|^2 - |\psi_2|^2) \\
&= \frac{1}{4}|\Psi^*K\Psi|^2
\end{aligned}$$

is obtained. Also the following holds

$$(29) \quad -Re(\Psi, F_{12}K\Psi) = -\frac{1}{4}|\Psi^*K\Psi|^2.$$

Similarly, by using (13), one can obtain

$$\begin{aligned}
(2) \quad (\Psi, F_{13}J\Psi) &= \left( \Psi, \left( \frac{1}{4}\Psi^*J\Psi \right) J\Psi \right) \\
&= \left( \frac{1}{4}\Psi^*J\Psi \right) (\Psi, J\Psi) \\
&= \frac{i}{4}(\overline{\psi_2}\psi_1 + \overline{\psi_1}\psi_2)i(\overline{\psi_1}\psi_2 + \overline{\psi_2}\psi_1) \\
&= -\frac{1}{4}|\Psi^*J\Psi|^2
\end{aligned}$$

and then

$$(30) \quad -Re(\Psi, F_{13}J\Psi) = \frac{1}{4}|\Psi^*J\Psi|^2.$$

At the end, with the aid of (13) the following identity holds

$$\begin{aligned}
(3) \quad (\Psi, F_{23}I\Psi) &= \left( \Psi, \left( -\frac{1}{4}\Psi^*I\Psi \right) I\Psi \right) \\
&= -\left( \frac{1}{4}\Psi^*I\Psi \right) (\Psi, I\Psi) \\
&= \frac{1}{4}(\overline{\psi_2}\psi_1 - \overline{\psi_1}\psi_2)(\overline{\psi_1}\psi_2 - \overline{\psi_2}\psi_1) \\
&= \frac{1}{4}|\Psi^*I\Psi|^2.
\end{aligned}$$

Then,

$$(31) \quad -Re(\Psi, F_{23}I\Psi) = -\frac{1}{4}|\Psi^*I\Psi|^2$$

is obtained. By inserting (29), (30), (31) into (28), one has

$$(32) \quad \Delta|\Psi|^2 = -2 \sum_{i=1}^3 |\nabla_i^A \Psi|^2 - \frac{1}{4}|\Psi^*K\Psi|^2 + \frac{1}{4}|\Psi^*J\Psi|^2 - \frac{1}{4}|\Psi^*I\Psi|^2.$$

Accordingly, the last three terms are obtained as:

$$\begin{aligned}
&|\Psi^*K\Psi|^2 - |\Psi^*J\Psi|^2 + |\Psi^*I\Psi|^2 \\
&= (|\psi_1|^2 - |\psi_2|^2)^2 + (\overline{\psi_2}\psi_1 + \overline{\psi_1}\psi_2)^2 + (\overline{\psi_2}\psi_1 - \overline{\psi_1}\psi_2)^2
\end{aligned}$$

$$\begin{aligned}
&= |\psi_1|^4 - 2|\psi_1|^2|\psi_2|^2 + |\psi_2|^4 - \overline{\psi_2}^2\psi_1^2 + 2|\psi_2|^2|\psi_1|^2 - \overline{\psi_1}^2\psi_2^2 \\
&\quad + \overline{\psi_1}^2\psi_2^2 + 2|\psi_1|^2|\psi_2|^2 + \overline{\psi_2}^2\psi_1^2 \\
&= (|\psi_1|^2 + |\psi_2|^2)^2 \\
(33) \quad &= |\Psi|^4.
\end{aligned}$$

After inserting (33) in (32), we get  $\Delta|\Psi|^2 \leq 0$  which means that the function

$$x \longrightarrow |\Psi(x)|^2 : \mathbb{R}^3 \longrightarrow \mathbb{R}$$

is subharmonic on  $\mathbb{R}^3$  [21]. As a result,  $|\Psi(x)|^2$  satisfies the mean value inequality for subharmonic functions. According to the Mean value theorem for subharmonic function, the following inequality is satisfied for any  $r > 0$  and any  $x \in \mathbb{R}^3$ ,

$$|\Psi(x)|^2 \leq \frac{3}{4\pi r^3} \int_{B_r(x)} |\Psi(x)|^2 dvol,$$

where  $B_r(x)$  is the closed ball of radius  $r$  about  $x$  [7, 9, 22]. If  $\Psi \in L^2$ ,  $\int_{\mathbb{R}^3} |\Psi(x)|^2 dvol < \infty$ . Hence,  $L^2$ -norm of  $\Psi$  is finite. Denoting the value of this integral by  $\kappa$ , we obtain

$$|\Psi(x)|^2 \leq \frac{3\kappa}{4\pi r^3}.$$

Since the  $L^2$ -norm of  $\Psi$  is finite it follows, by taking the limit  $r \rightarrow \infty$ , that  $\Psi(x) = 0$  for all  $x \in \mathbb{R}^3$ .

To prove second part, similar way is used. By inserting (33) into (32), one can provide

$$(34) \quad \Delta|\Psi|^2 = -2 \sum_{i=1}^3 |\nabla_i^A \Psi|^2 - \frac{1}{4} |\Psi|^4.$$

By means of standard identity from vector calculus, one has

$$\Delta(f \cdot g) = f \cdot \Delta g - 2\nabla f \cdot \nabla g + g \cdot \Delta f.$$

Taking  $g = f$ , one can provide

$$\Delta(f^2) = -2\nabla f \cdot \nabla f + 2f \cdot \Delta f,$$

so

$$\begin{aligned}
\Delta|\Psi|^4 &= \Delta\left(|\Psi|^2\right)^2 \\
&= -2\left(\nabla|\Psi|^2\right) \cdot \left(\nabla|\Psi|^2\right) + 2|\Psi|^2 \Delta|\Psi|^2
\end{aligned}$$

then

$$\Delta|\Psi|^4 = -2\left(\nabla|\Psi|^2\right) \cdot \left(\nabla|\Psi|^2\right) - 4|\Psi|^2 \sum_{i=1}^3 |\nabla_i^A \Psi|^2 - \frac{1}{2} |\Psi|^6.$$

Consequently,  $\Delta|\Psi|^4 \leq 0$  on  $\mathbb{R}^3$  so  $x \rightarrow |\Psi(x)|^4$  is subharmonic on  $\mathbb{R}^3$ . Thus, for every  $r > 0$  and every  $x \in \mathbb{R}^3$ ,

$$(35) \quad |\Psi(x)|^4 \leq \frac{3}{4\pi r^3} \int_{B_r(x)} |\Psi(x)|^4 dvol.$$

The assumptions  $E(A, \Psi) < \infty$  can be written as in the following

$$(36) \quad E(A, \Psi) = \int_M \left( |F_A|^2 + |\nabla^A \Psi|^2 + \frac{R}{4} |\Psi|^2 + \frac{1}{16} |\Psi|^4 \right) dvol < \infty.$$

From (36), one has

$$(37) \quad \int_{\mathbb{R}^3} |\Psi|^4 dvol < \infty.$$

This means that  $\Psi \equiv 0$  on  $\mathbb{R}^3$ . Consequently under the assumption  $E(A, \Psi) < \infty$ ,  $F_A \equiv 0$  since  $\Psi \equiv 0$ .  $\square$

### 5. A non-trivial solution to Seiberg-Witten-like equations on 3-dimensional contact metric manifolds

In this section, Seiberg-Witten-like equations on the 3-dimensional strictly pseudoconvex  $CR$ -3 manifolds are written and a global solution to these equations is given.

On the 3-dimensional strictly pseudoconvex  $CR$ -3 manifolds, the spinor bundle can be decomposed as follows:

$$\mathbb{S} \cong \Lambda_H^{0,1}(M) \oplus \Lambda_H^{0,0}(M),$$

where  $\Lambda_H^{0,1}(M)$  is the eigenspace corresponding to the eigenvalue  $i$  of the mapping  $\kappa(d\alpha) : \mathbb{S} \rightarrow \mathbb{S}$  and has dimension 1,  $\Lambda_H^{0,0}(M)$  is the eigenspace corresponding to the eigenvalue  $-i$  of the mapping  $\kappa(d\alpha) : \mathbb{S} \rightarrow \mathbb{S}$  and has dimension 1. If  $\Psi_0 \in \mathbb{S}$ , isomorphic to the constant function  $1 \in \Lambda_H^{0,0}(M)$ , then  $\Psi_0$  denotes the spinor corresponding to the constant function 1 in the chosen coordinates

$$\Psi_0 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

By using the expression of  $\sigma_H(\Psi)$  in the local coordinates and  $d\alpha \cdot \Psi_0 = -i\Psi_0$ , the following identity is obtained:

$$\sigma_H(\Psi_0) = id\alpha.$$

On the subbundle  $H$ , the Ricci form  $\rho_H$  is defined by

$$(38) \quad \rho_H(X, Y) = Ric(X, J_H Y) = g_\alpha(X, J_H Ric Y)$$

for any  $X, Y \in \Gamma(H)$ . Since on the strictly pseudoconvex  $CR$  manifold, the almost complex structure  $J_H$  is complex,

$$(39) \quad Ric(X, Y) = i\rho_H(X, Y)$$

for any  $X, Y \in \Gamma(H)$ .

**Proposition 5.1.** *Suppose that  $\rho_H$  be a Ricci form on the subbundle  $H$  and  $s_H$  be a scalar curvature of  $H$ . Then, one can satisfy the following identity:*

$$(40) \quad \rho = -\frac{s_H}{2}d\alpha.$$

*Proof.* According to the local coordinates, the almost complex structure  $J$  is given as follows:

$$J = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Since  $J \circ Ric = Ric \circ J$  commutative, the reduced form of the  $Ric$  is

$$Ric = \begin{bmatrix} R_{11} & 0 & 0 \\ 0 & R_{11} & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

By using (38) the explicit form of  $\rho_H$  is:

$$(41) \quad \rho_H = -R_{11}e^1 \wedge e^2 = -\frac{s_H}{2}d\alpha. \quad \square$$

In the following theorem, a special solution of Seiberg-Witten-like equations is given on the 3-dimensional strictly pseudoconvex contact metric manifold.

**Theorem 5.2.** *Let  $(M, g_\alpha, \alpha, \xi, J)$  be a strictly pseudoconvex CR-3 manifold. Then, for a given negative and constant scalar curvature  $s_H$ ,  $(A, \Psi = \sqrt{-2s_H}\Psi_0)$  is the solution of Seiberg-Witten-like equations.*

*Proof.* By using  $\Psi$  we get  $\sigma_H(\Psi) = id\alpha$ . Also it can be written as

$$(42) \quad \sigma_H(\Psi) = -2is_H d\alpha.$$

By using (39) and (42), one can satisfy,

$$(43) \quad F_A = Ric = i\rho_H = -i\frac{s_H}{2}d\alpha = \frac{1}{4}\sigma_H(\Psi).$$

Since  $\sigma_H(\Psi) = \sigma(\Psi)$ ,

$$(44) \quad F_A = \frac{1}{4}\sigma(\Psi)$$

is satisfied.

The following is easily hold. By using the spinor field  $\Psi_0$  corresponding to the constant function 1, one can obtain

$$(45) \quad \mathcal{H}(1) = \sqrt{2} \sum_{r=0}^n (\bar{\partial}_H + \bar{\partial}_H^*)(1) + \sum_{r=0}^n (-1)^{r+1} \sqrt{-1} \cdot \nabla_\xi^{TW}(1) = 0.$$

This means that  $D_{A_0}\Psi = D_H^{A_0}\Psi + \xi \cdot \nabla_\xi^{A_0}\Psi = 0$ .

As a result,  $(A, \Psi = \sqrt{-2s_H}\Psi_0)$  is the solution of Seiberg-Witten-like equations on the strictly pseudoconvex CR-3 manifold.  $\square$

A 3-dimensional Hiperbolic space with a negative and constant scalar curvature can be given for the Theorem (5.2) (see [26]).

### References

- [1] H. Baum, *Lorentzian twistor spinors and CR-geometry*, Differential Geom. Appl. **11** (1999), no. 1, 69–96. [https://doi.org/10.1016/S0926-2245\(99\)00020-0](https://doi.org/10.1016/S0926-2245(99)00020-0)
- [2] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds*, Progress in Mathematics, **203**, Birkhäuser Boston, Inc., Boston, MA, 2002. <https://doi.org/10.1007/978-1-4757-3604-5>
- [3] A. L. Carey, B. L. Wang, R. Zhang, and J. McCarthy, *Seiberg-Witten monopoles in three dimensions*, Lett. Math. Phys. **39** (1997), no. 3, 213–228. <https://doi.org/10.1023/A:1007319915035>
- [4] N. Değirmenci and Bulut, *Seiberg-Witten-like equations on 6-dimensional  $SU(3)$ -manifolds*, Balkan J. Geom. Appl. **20** (2015), no. 2, 23–31.
- [5] N. Değirmenci and N. Özdemir, *Seiberg-Witten-like equations on 7-manifolds with  $G_2$ -structure*, J. Nonlinear Math. Phys. **12** (2005), no. 4, 457–461.
- [6] S. Eker, *Seiberg-Witten equations on 8-dimensional manifolds with different self-duality*, Balkan J. Geom. Appl. **22** (2017), no. 2, 37–43.
- [7] C. Fefferman and E. M. Stein,  *$H^p$  spaces of several variables*, Acta Math. **129** (1972), no. 3-4, 137–193. <https://doi.org/10.1007/BF02392215>
- [8] T. Friedrich, *Dirac operators in Riemannian geometry*, translated from the 1997 German original by Andreas Nestke, Graduate Studies in Mathematics, **25**, American Mathematical Society, Providence, RI, 2000. <https://doi.org/10.1090/gsm/025>
- [9] J. E. Littlewood and G. H. Hardy, *Some properties of conjugate functions*, J. Reine Angew. Math. **167** (1932), 405–423. <https://doi.org/10.1515/crll.1932.167.405>
- [10] J. Lohkamp, *Metrics of negative Ricci curvature*, Ann. of Math. (2) **140** (1994), no. 3, 655–683. <https://doi.org/10.2307/2118620>
- [11] P. Kronheimer and T. Mrowka, *Monopoles and three-manifolds*, New Mathematical Monographs, **10**, Cambridge University Press, Cambridge, 2007. <https://doi.org/10.1017/CB09780511543111>
- [12] H. B. Lawson, Jr. and M.-L. Michelsohn, *Spin geometry*, Princeton Mathematical Series, **38**, Princeton University Press, Princeton, NJ, 1989.
- [13] J. W. Morgan, *The Seiberg-Witten equations and applications to the topology of smooth four-manifolds*, Mathematical Notes, **44**, Princeton University Press, Princeton, NJ, 1996.
- [14] T. Mrowka, P. Ozsváth, and B. Yu, *Seiberg-Witten monopoles on Seifert fibered spaces*, Comm. Anal. Geom. **5** (1997), no. 4, 685–791. <https://doi.org/10.4310/CAG.1997.v5.n4.a3>
- [15] G. L. Naber, *Topology, geometry, and gauge fields*, Texts in Applied Mathematics, **25**, Springer-Verlag, New York, 1997. <https://doi.org/10.1007/978-1-4757-2742-5>
- [16] T. Nguyen, *The Seiberg-Witten equations on manifolds with boundary I: the space of monopoles and their boundary values*, Comm. Anal. Geom. **20** (2012), no. 3, 565–676. <https://doi.org/10.4310/CAG.2012.v20.n3.a5>
- [17] L. I. Nicolaescu, *Adiabatic limits of the Seiberg-Witten equations on Seifert manifolds*, Comm. Anal. Geom. **6** (1998), no. 2, 331–392. <https://doi.org/10.4310/CAG.1998.v6.n2.a5>
- [18] ———, *Notes on Seiberg-Witten theory*, Graduate Studies in Mathematics, **28**, American Mathematical Society, Providence, RI, 2000. <https://doi.org/10.1090/gsm/028>
- [19] R. Petit, *Harmonic maps and strictly pseudoconvex CR manifolds*, Comm. Anal. Geom. **10** (2002), no. 3, 575–610. <https://doi.org/10.4310/CAG.2002.v10.n3.a5>



- [20] R. Petit, *Spin<sup>c</sup>-structures and Dirac operators on contact manifolds*, Differential Geom. Appl. **22** (2005), no. 2, 229–252. <https://doi.org/10.1016/j.difgeo.2005.01.003>
- [21] D. Salamon, *Spin geometry and Seiberg-Witten invariants*, Citeseer, 1996.
- [22] A. Shkheam, *Spaces of harmonic functions and harmonic quasiconformal mappings*, Doctoral Thesis, Belgrade, 2013.
- [23] A. I. Stipsicz, *Gauge theory and Stein fillings of certain 3-manifolds*, Turkish J. Math. **26** (2002), no. 1, 115–130.
- [24] S. Tanno, *Variational problems on contact Riemannian manifolds*, Trans. Amer. Math. Soc. **314** (1989), no. 1, 349–379. <https://doi.org/10.2307/2001446>
- [25] C. H. Taubes, *The Seiberg-Witten equations and the Weinstein conjecture*, Geom. Topol. **11** (2007), 2117–2202. <https://doi.org/10.2140/gt.2007.11.2117>
- [26] T. Jørgensen, *Compact 3-manifolds of constant negative curvature fibering over the circle*, Ann. of Math. (2) **106** (1977), no. 1, 61–72. <https://doi.org/10.2307/1971158>
- [27] E. Witten, *Monopoles and four-manifolds*, Math. Res. Lett. **1** (1994), no. 6, 769–796. <https://doi.org/10.4310/MRL.1994.v1.n6.a13>

SERHAN EKER  
DEPARTMENT MATHEMATICS  
AĞRI İBRAHİM ÇEÇEN UNIVERSITY  
AĞRI 04000, TURKEY  
Email address: [srhaneker@gmail.com](mailto:srhaneker@gmail.com)