

A NOTE ON COHOMOLOGICAL DIMENSION OVER COHEN-MACAULAY RINGS

IRAJ BAGHERIYEH, KAMAL BAHMANPOUR, AND GHADER GHASEMI

ABSTRACT. Let (R, \mathfrak{m}) be a Noetherian local Cohen-Macaulay ring and I be a proper ideal of R . Assume that $\beta_R(I, R)$ denotes the constant value of $\text{depth}_R(R/I^n)$ for $n \gg 0$. In this paper we introduce the new notion $\gamma_R(I, R)$ and then we prove the following inequalities:

$$\beta_R(I, R) \leq \gamma_R(I, R) \leq \dim R - \text{cd}(I, R) \leq \dim R/I.$$

Also, some applications of these inequalities will be included.

1. Introduction

Let R denote a commutative Noetherian ring (with identity), I, J be two ideals of R , and M be a finitely generated R -module. Ratliff in [15] conjectured that the set $\text{Ass}_R(M/J^n M)$ stabilizes for $n \gg 0$, when R is a Noetherian domain. Subsequently, Brodmann in [4] showed that if R is a Noetherian ring, then the sets $\text{Ass}_R(M/J^n M)$ and $\text{Ass}_R(J^n M/J^{n+1} M)$ are ultimately constant for large n . Also, based on this result, in [5] he showed that if R is a Noetherian ring, then the integers $\text{grade}(I, M/J^n M)$ and $\text{grade}(I, J^n M/J^{n+1} M)$ take constant values for large n . In particular, the integers $\text{depth}_R(M/J^n M)$ and $\text{depth}_R(J^n M/J^{n+1} M)$ take constant values for large n , when (R, \mathfrak{m}) is Noetherian local ring. In the sequel let $\beta_R(I, M)$ denote the constant value of $\text{depth}_R(M/I^n M)$ for $n \gg 0$, when R is local. In [1] it has been proved that if M is a non-zero finitely generated R -module and I is an ideal of R with $\beta_R(I, M) = \dim M$, then M annihilates by some power of I and so M is a Cohen-Macaulay R -module.

For an R -module M , the i th local cohomology module of M with support in $V(I)$ is defined as:

$$H_I^i(M) = \varinjlim_{n \geq 1} \text{Ext}_R^i(R/I^n, M).$$

We refer the reader to [6] or [10] for more details about local cohomology.

Received February 10, 2019; Revised July 4, 2019; Accepted July 26, 2019.

2010 *Mathematics Subject Classification.* Primary 13D45, 13E05.

Key words and phrases. canonical module, Cohen-Macaulay ring, cohomological dimension, local cohomology, Noetherian ring.

For an R -module M , the notion $\text{cd}(I, M)$, *the cohomological dimension of M with respect to I* , is defined as:

$$\text{cd}(I, M) = \sup\{i \in \mathbb{N}_0 : H_I^i(M) \neq 0\}$$

with the usual convention that the supremum of the empty set of integers is interpreted as $-\infty$. This notion have been studied by several authors (see [3, 7–9, 11, 12]).

In this paper first we shall introduce the new notion $\gamma_R(I, R)$ and then we will establish some new inequalities between $\text{cd}(I, R)$, $\beta_R(I, R)$ and $\gamma_R(I, R)$. Also, some applications of these inequalities will be included.

Throughout this paper, (R, \mathfrak{m}) will always be a commutative Noetherian local ring with non-zero identity. For each R -module M , we denote by $E_R(M)$ the injective envelope (or injective hull) of M . Also, for any ideal \mathfrak{b} of R , *the radical of \mathfrak{b}* , denoted by $\text{Rad}(\mathfrak{b})$, is defined to be the set $\{x \in R : x^n \in \mathfrak{b} \text{ for some } n \in \mathbb{N}\}$. For any other unexplained notation and terminology we refer the reader to [6] and [13].

2. The results

The main purpose of this section is to prove Theorem 2.3. But, first we need to introduce the new notion $\gamma_R(I, M)$.

Definition. Let (R, \mathfrak{m}) be a local ring and I be an ideal of R . For every finitely generated R -module M set $G(I, M) := \bigoplus_{n=1}^{\infty} M/I^n M$. Then we define $\gamma_R(I, M)$ as:

$$\gamma_R(I, M) := \inf\{i \in \mathbb{N}_0 : I \not\subseteq \text{Rad}(\text{Ann}_R H_{\mathfrak{m}}^i(G(I, M)))\},$$

with the convention that the infimum of the empty set of integers is interpreted as ∞ .

From the definition it follows that there exists a positive integer k such that

$$I^k \subseteq \text{Ann}_R \bigoplus_{j=0}^{\gamma_R(I, M)-1} H_{\mathfrak{m}}^j(G(I, M)).$$

Furthermore, since the local cohomology functor $H_{\mathfrak{m}}^i(-)$ commutes with the direct sums it follows that $I^k H_{\mathfrak{m}}^j(M/I^n M) = 0$ for each $0 \leq j \leq \gamma_R(I, M) - 1$ and each $n \geq 1$.

Now we consider the following question.

Question 1: Let (R, \mathfrak{m}) be a local Cohen-Macaulay ring and I be an ideal of R . Whether $\gamma_R(I, R) = \dim R - \text{cd}(I, R)$?

In this paper we shall prove that the inequality $\gamma_R(I, R) \leq \dim R - \text{cd}(I, R)$ always holds. So, Question 1 reduces to the following question:

Question 2: Let (R, \mathfrak{m}) be a local Cohen-Macaulay ring and I be an ideal of R . Whether $\gamma_R(I, R) \geq \dim R - \text{cd}(I, R)$?

The following easy lemma is needed in the proof of Theorem 2.2.

Lemma 2.1. *Let (R, \mathfrak{m}) be a local ring, I be an ideal of R and M be a finitely generated R -module. If $\text{cd}(I, M) = t \geq 1$, then $H_I^t(M) = IH_I^t(M)$.*

In particular,

$$I \not\subseteq \text{Rad}(\text{Ann}_R H_I^t(M)).$$

Proof. In view of [2, Lemma 2.8], $H_I^t(M) = IH_I^t(M)$. Since, $H_I^t(M) \neq 0$ it follows that $I^k H_I^t(M) = H_I^t(M) \neq 0$ for each $k \geq 1$ and hence

$$I \not\subseteq \text{Rad}(\text{Ann}_R H_I^t(M)). \quad \square$$

Now we are ready to state and prove the main result of this paper.

Theorem 2.2. *Let (R, \mathfrak{m}) be a local Cohen-Macaulay ring and I be an ideal of R . Then,*

$$\beta_R(I, R) \leq \gamma_R(I, R) \leq \dim R - \text{cd}(I, R) \leq \dim R/I.$$

In particular, if $\gamma_R(I, R) = \dim R/I$, then

$$\gamma_R(I, R) = \dim R - \text{cd}(I, R) \text{ and } \text{cd}(I, R) = \text{height } I.$$

Proof. Assume that $\beta_R(I, R) = t$. If $t = 0$, then it is clear that

$$\beta_R(I, R) \leq \gamma_R(I, R).$$

Now assume that $t \geq 1$. Then, by the definition there is a positive integer k such that for each $n \geq k$, $\text{depth}_R R/I^n = t$. Thus, for each $0 \leq j \leq t-1$ and each $n \geq k$, $H_{\mathfrak{m}}^j(R/I^n) = 0$. On the other hand, for each $0 \leq j \leq t-1$ and each $n < k$ it is clear that $I^k H_{\mathfrak{m}}^j(R/I^n) = 0$. Therefore, for each $0 \leq j \leq t-1$, $I^k H_{\mathfrak{m}}^j(G(I, R)) = 0$ and hence $I \subseteq \text{Rad}(\text{Ann}_R H_{\mathfrak{m}}^j(G(I, R)))$. Now, it follows from the definition that $\gamma_R(I, R) \geq t = \beta_R(I, R)$.

Let \widehat{R} denote the \mathfrak{m} -adic completion of R . Then $(\widehat{R}, \mathfrak{m}\widehat{R})$ is a Noetherian local Cohen-Macaulay ring such that $\gamma_{\widehat{R}}(I\widehat{R}, \widehat{R}) = \gamma_R(I, R)$, $\dim \widehat{R} = \dim R$, $\dim \widehat{R}/I\widehat{R} = \dim R/I$ and $\text{cd}(I\widehat{R}, \widehat{R}) = \text{cd}(I, R)$. Therefore, in order to prove the inequality $\gamma_R(I, R) \leq \dim R - \text{cd}(I, R)$, without loss of generality we may assume that R is a Noetherian complete local Cohen-Macaulay ring.

Let $\gamma_R(I, R) = \ell$ and assume that ω_R denote the canonical module of R . Then by the definition there exists a positive integer k such that for each $0 \leq j \leq \ell-1$ and each $n \geq 1$, $I^k H_{\mathfrak{m}}^j(R/I^n) = 0$. Therefore, in view of *Local Duality Theorem* for each $0 \leq j \leq \ell-1$ and each $n \geq 1$, $I^k \text{Ext}_R^{\dim R-j}(R/I^n, \omega_R) \simeq I^k D(H_{\mathfrak{m}}^j(R/I^n)) = 0$, where $D(-)$ denotes the Matlis dual functor $\text{Hom}_R(-, E_R(R/\mathfrak{m}))$. Thus, for each $0 \leq j \leq \ell-1$,

$$I^k H_I^{\dim R-j}(\omega_R) = I^k \left(\varinjlim_{n \geq 1} \text{Ext}_R^{\dim R-j}(R/I^n, \omega_R) \right) = 0.$$

Also, by the *Grothendieck's Vanishing Theorem*, $H_I^{\dim R+i}(\omega_R) = 0$ for each $i \geq 1$. Hence, by Lemma 2.1, $\text{cd}(I, \omega_R) \leq \dim R - \ell$.

Since,

$$\text{Supp } \omega_R = \text{Spec}(R) = \text{Supp } R,$$

it follows from [7, Theorem 2.2] that $\text{cd}(I, R) = \text{cd}(I, \omega_R)$ and so

$$\text{cd}(I, R) \leq \dim R - \ell = \dim R - \gamma_R(I, R).$$

Now, it is clear that $\gamma_R(I, R) \leq \dim R - \text{cd}(I, R)$.

On the other hand, in view of [14, Lemma 2.10],

$$\dim R - \text{cd}(I, R) \leq \dim R/I.$$

Therefore,

$$\beta_R(I, R) \leq \gamma_R(I, R) \leq \dim R - \text{cd}(I, R) \leq \dim R/I.$$

Now, assume that $\gamma_R(I, R) = \dim R/I$. Then it is clear that

$$\gamma_R(I, R) = \dim R - \text{cd}(I, R) \text{ and } \dim R/I = \dim R - \text{cd}(I, R).$$

So, by [13, Theorem 17.4],

$$\text{cd}(I, R) = \dim R - \dim R/I = \text{height } I. \quad \square$$

Corollary 2.3. *Let (R, \mathfrak{m}) be a local Cohen-Macaulay ring of dimension d and I be an ideal of R with $\beta_R(I, R) = d - 1$. Then $\text{cd}(I, R) = 1$.*

Proof. In view of Theorem 2.2, $\text{cd}(I, R) \leq \dim R - \beta_R(I, R) = 1$.

Now, if $\text{cd}(I, R) = 0$, then I is nilpotent and hence $\beta_R(I, R) = \dim R = d$, which is a contradiction. So, $\text{cd}(I, R) = 1$. \square

The following consequence of Theorem 2.2 presents a partially affirmative answer to Question 2 in an special case.

Corollary 2.4. *Let (R, \mathfrak{m}) be a local Cohen-Macaulay ring of dimension d and I be an ideal of R with $\gamma_R(I, R) = d - 1$. Then $\text{cd}(I, R) = 1$.*

Proof. In view of Theorem 2.2, $\text{cd}(I, R) \leq \dim R - \gamma_R(I, R) = 1$.

Now, if $\text{cd}(I, R) = 0$, then I is a nilpotent ideal and so

$$\gamma_R(I, R) \geq \beta_R(I, R) = \dim R = d,$$

which is a contradiction. Therefore, $\text{cd}(I, R) = 1$. \square

Acknowledgements. The authors are deeply grateful to the referee for a very careful reading of the manuscript and many valuable suggestions. Also, we would like to thank Professor Reza Naghipour for his careful reading of the first draft and many helpful suggestions.

References

- [1] K. Bahmanpour, *Cohen-Macaulay modules over Noetherian local rings*, Bull. Korean Math. Soc. **51** (2014), no. 2, 373–386. <https://doi.org/10.4134/BKMS.2014.51.2.373>
- [2] ———, *Exactness of ideal transforms and annihilators of top local cohomology modules*, J. Korean Math. Soc. **52** (2015), no. 6, 1253–1270. <https://doi.org/10.4134/JKMS.2015.52.6.1253>
- [3] K. Bahmanpour and M. S. Samani, *On the cohomological dimension of finitely generated modules*, Bull. Korean Math. Soc. **55** (2018), no. 1, 311–317. <https://doi.org/10.4134/BKMS.b161017>
- [4] M. Brodmann, *Asymptotic stability of $\text{Ass}(M/I^n M)$* , Proc. Amer. Math. Soc. **74** (1979), no. 1, 16–18. <https://doi.org/10.2307/2042097>
- [5] ———, *The asymptotic nature of the analytic spread*, Math. Proc. Cambridge Philos. Soc. **86** (1979), no. 1, 35–39. <https://doi.org/10.1017/S030500410000061X>
- [6] M. P. Brodmann and R. Y. Sharp, *Local cohomology*, second edition, Cambridge Studies in Advanced Mathematics, **136**, Cambridge University Press, Cambridge, 2013.
- [7] K. Divaani-Aazar, R. Naghipour, and M. Tousi, *Cohomological dimension of certain algebraic varieties*, Proc. Amer. Math. Soc. **130** (2002), no. 12, 3537–3544. <https://doi.org/10.1090/S0002-9939-02-06500-0>
- [8] G. Faltings, *Über lokale Kohomologiegruppen hoher Ordnung*, J. Reine Angew. Math. **313** (1980), 43–51. <https://doi.org/10.1515/crll.1980.313.43>
- [9] G. Ghasemi, K. Bahmanpour, and J. A'zami, *Upper bounds for the cohomological dimensions of finitely generated modules over a commutative Noetherian ring*, Colloq. Math. **137** (2014), no. 2, 263–270. <https://doi.org/10.4064/cm137-2-10>
- [10] R. Hartshorne, *Local Cohomology*, A seminar given by A. Grothendieck, Harvard University, Fall, 1961. Lecture Notes in Mathematics, No. 41, Springer-Verlag, Berlin, 1967.
- [11] ———, *Cohomological dimension of algebraic varieties*, Ann. of Math. (2) **88** (1968), 403–450. <https://doi.org/10.2307/1970720>
- [12] C. Huneke and G. Lyubeznik, *On the vanishing of local cohomology modules*, Invent. Math. **102** (1990), no. 1, 73–93. <https://doi.org/10.1007/BF01233420>
- [13] H. Matsumura, *Commutative Ring Theory*, translated from the Japanese by M. Reid, Cambridge Studies in Advanced Mathematics, **8**, Cambridge University Press, Cambridge, 1986.
- [14] A. A. Mehrvarz, K. Bahmanpour, and R. Naghipour, *Arithmetic rank, cohomological dimension and filter regular sequences*, J. Algebra Appl. **8** (2009), no. 6, 855–862. <https://doi.org/10.1142/S0219498809003692>
- [15] L. J. Ratliff, Jr., *On prime divisors of I^n , n large*, Michigan Math. J. **23** (1976), no. 4, 337–352 (1977). <http://projecteuclid.org/euclid.mmj/1029001769>

IRAJ BAGHERIYEH
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES
UNIVERSITY OF MOHAGHEGH ARDABILI
56199-11367, ARDABIL, IRAN
Email address: Ir_ba2004@yahoo.com

KAMAL BAHMANPOUR
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES
UNIVERSITY OF MOHAGHEGH ARDABILI
56199-11367, ARDABIL, IRAN
Email address: bahmanpour.k@gmail.com

GHADER GHASEMI
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCES
UNIVERSITY OF MOHAGHEGH ARDABIL
56199-11367, ARDABIL, IRAN
Email address: ghghasemi@gmail.com

51YUX.CZDFbR