

EXACT FORMULA FOR JACOBI-EISENSTEIN SERIES OF SQUARE FREE DISCRIMINANT LATTICE INDEX

RAN XIONG

ABSTRACT. In this paper we give an exact formula for the Fourier coefficients of the Jacobi-Eisenstein series of square free discriminant lattice index. For a special case the discriminant of lattice is prime we show that the Jacobi-Eisenstein series corresponds to a well known Eisenstein series of modular forms.

1. Introduction

Jacobi forms of lattice index, whose theory can be viewed as an extension of the theory of classical Jacobi forms, play important roles in many mathematical fields, such as the theory of orthogonal modular forms, the theory of vertex operator algebras. A classical example of Jacobi form is Jacobi-Eisenstein series. In [1], Ajouz defined Jacobi-Eisenstein series of lattice index and studied its basic properties. In [6], Mocanu gave the first formula of Fourier expansion for Jacobi-Eisenstein series of lattice index and, for the trivial case, she showed that the Fourier coefficients in fact are special values of Dirichlet L -functions up to finite Euler factors. In [7], Woitalla considered this for the lattice $\bigoplus_{1 \leq j \leq N} A_1$ where A_1 is the scalar lattice $(\mathbb{Z}, (x, y) \rightarrow 2xy)$. In this short paper we give an exact formula for Fourier coefficients of Jacobi-Eisenstein series of square free discriminant lattice index. This type lattice has occurred in several problems.

This paper is organized as follows: In Section 2 we review some basic facts for Jacobi forms of lattice index briefly. An exact formula for Jacobi-Eisenstein series of square free discriminant index is given and proved in Section 3. Finally, for the case the discriminant of lattice is prime we refer that the Jacobi-Eisenstein series corresponds to a well known Eisenstein series of elliptic modular forms.

Received April 1, 2019; Accepted August 23, 2019.

2010 *Mathematics Subject Classification.* 11F50, 11M36.

Key words and phrases. Jacobi forms of lattice index, Fourier coefficients of Jacobi-Eisenstein series.

This work is supported by the Fundamental Research Funds for the Central Universities of China (grant no. 22120180508).

2. Jacobi forms of lattice index

Throughout this paper, for a complex number t , write $e(t) := e^{2\pi it}$, $q = e(\tau)$ for τ of the complex upper half plane \mathbb{H} . For a prime p , a rational number m , denote $\nu_p(m)$ the p -adic valuation of m . We put $\Gamma = SL_2(\mathbb{Z})$ and, for a positive integer $N > 1$, set the congruence subgroup $\Gamma_0(N) := \{\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma : N \mid c\}$.

We call a pair $\underline{L} = (L, \beta)$ a positive definite even lattice if L is a \mathbb{Z} -module of finite rank $r_{\underline{L}}$, equipped with a positive definite integral quadratic form β . Denote by L^\sharp the dual lattice

$$L^\sharp := \{y \in L \otimes_{\mathbb{Z}} \mathbb{Q} : \beta(y, x) \in \mathbb{Z} \text{ for all } x \in L\}.$$

The discriminant form of \underline{L} is defined as

$$D_{\underline{L}} := (L^\sharp/L, x + L \rightarrow \beta(x) + \mathbb{Z}).$$

Note that $\text{card}(L^\sharp/L) = \det(\underline{L}) := \det(F)$ where F is the Gram matrix corresponding to a \mathbb{Z} -basis of L . For $r \in L^\sharp$, let ω_r be its order in L^\sharp/L . Define the discriminant of \underline{L} as

$$\Delta_{\underline{L}} := \begin{cases} (-1)^{\lfloor \frac{r_{\underline{L}}}{2} \rfloor} \det(\underline{L}), & r_{\underline{L}} \text{ even,} \\ (-1)^{\lfloor \frac{r_{\underline{L}}}{2} \rfloor} 2 \det(\underline{L}), & r_{\underline{L}} \text{ odd.} \end{cases}$$

It is well known that $\Delta_{\underline{L}} \equiv 0, 1 \pmod{4}$. For $a \in \mathbb{Z}$, $D \in \mathbb{Q}$ such that $\Delta_{\underline{L}} D \in \mathbb{Z}$, denote $\chi_{\underline{L}}(D, a) := \left(\frac{\Delta_{\underline{L}} D}{a}\right)$ and in particular put $\chi_{\underline{L}}(a) := \chi_{\underline{L}}(1, a)$.

Let $\underline{L} = (L, \beta)$ be a positive definite even lattice and k an integer. The space $J_{k, \underline{L}}(\Gamma)$ of Jacobi forms of weight k and index \underline{L} consists of all holomorphic functions $\phi(\tau, z)$ on $\mathbb{H} \times (L \otimes_{\mathbb{Z}} \mathbb{C})$ which satisfy

- (1) For any $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma$ and $h = (x, y) \in L \times L$, the following identities hold:

$$\phi(\tau, z) = (\phi|_{k, \underline{L}} A)(\tau, z) := (c\tau + d)^{-k} e\left(\frac{-c\beta(z)}{c\tau + d}\right) \phi\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right),$$

$$\phi(\tau, z) = (\phi|_{k, \underline{L}} h)(\tau, z) := e(\tau\beta(x) + \beta(x, z)) \phi(\tau, z + x\tau + y);$$

- (2) The Fourier expansion of ϕ is of the form

$$\phi(\tau, z) = \sum_{\substack{n \in \mathbb{Z}, r \in L^\sharp \\ \beta(r) - n \leq 0}} c_\phi(n, r) q^n \zeta_\beta^r.$$

Here and in the following, for $z \in L \otimes_{\mathbb{Z}} \mathbb{C}$ and $r \in L^\sharp$ we write $\zeta_\beta^r := e(\beta(r, z))$.

By [1, Chapter 2, Proposition 2.4.3], the quality of $c_\phi(n, r)$ depends on the value of $\beta(r) - n$ and $r \pmod{L}$. In the following we write $D := \beta(r) - n$ and $C_\phi(D, r) = c_\phi(n, r)$ as usual.

Define the Jacobi theta series $\vartheta_{\underline{L}}(\tau, z)$ as

$$\vartheta_{\underline{L}}(\tau, z) := \sum_{x \in L} q^{\beta(x)} \zeta_\beta^x.$$

For an even integer $k > \frac{r_{\underline{L}}}{2} + 2$, the Jacobi-Eisenstein series $E_{k,\underline{L}}(\tau, z)$ is defined as

$$E_{k,\underline{L}}(\tau, z) := \sum_{A \in \Gamma_{\infty} \setminus \Gamma} (\vartheta_{\underline{L}}(\tau, z)) \Big|_{k,\underline{L}} A,$$

where $\Gamma_{\infty} = \{\pm \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} : n \in \mathbb{Z}\}$. We have that $E_{k,\underline{L}}(\tau, z) \in J_{k,\underline{L}}(\Gamma)$.

3. Fourier expansion of Jacobi-Eisenstein series of square free discriminant index

From now on we assume that $\Delta_{\underline{L}}$, the discriminant of \underline{L} is square free. This implies that $r_{\underline{L}}$ is even and $D_{\underline{L}} \approx \bigoplus_{\substack{p|\Delta_{\underline{L}} \\ p \text{ prime}}} (\mathbb{Z}/p\mathbb{Z}, \frac{a_p x^2}{p})$. With notations as before the main result of this paper is stated as follows:

Theorem 3.1. *Let $\underline{L} = (L, \beta)$ be a positive definite even lattice of square free discriminant. Then for even integer k with $k > \frac{r_{\underline{L}}}{2} + 2$, the Jacobi-Eisenstein series $E_{k,\underline{L}}(\tau, z)$ of weight k and index \underline{L} has the following Fourier expansion*

$$E_{k,\underline{L}}(\tau, z) = \vartheta_{\underline{L}}(\tau, z) + \sum_{\substack{n \in \mathbb{Z}, r \in L^{\sharp} \\ \beta(r) < n}} c(n, r) q^n \zeta_{\beta}^r,$$

where the qualities of $c(n, r)$ are given by

$$c(n, r) = \frac{2(-1)^{\lceil \frac{r_{\underline{L}}}{4} \rceil}}{L(1-k + \frac{r_{\underline{L}}}{2}, \chi_{\underline{L}})} \sum_{d|\Delta_{\underline{L}}D} d^{k - \frac{r_{\underline{L}}}{2} - 1} \chi_{\underline{L}}(|\Delta_{\underline{L}}|D/d) \\ \times \prod_{\substack{p|\Delta_{\underline{L}} \\ p \nmid \omega_r}} \left(1 + p^{\nu_p(|\Delta_{\underline{L}}D|)(1-k + \frac{r_{\underline{L}}}{2})} \chi_{\underline{L}}(a_p \Delta_{\underline{L}}D / p^{\nu_p(\Delta_{\underline{L}}^2 D)}, p) \chi_{\underline{L}}(-1/p, p^{\nu_p(\Delta_{\underline{L}}D)}) \right).$$

Proof. The application $\Gamma_{\infty} A \mapsto \pm(0, 1)A$ gives a bijection

$$\Gamma_{\infty} \setminus \Gamma \rightarrow \{(c, d) \in \mathbb{Z}^2 : \gcd(c, d) = 1\} / \{\pm 1\}$$

therefore by the definition of Jacobi-Eisenstein series,

$$E_{k,\underline{L}}(\tau, z) = \vartheta_{\underline{L}}(\tau, z) + \sum_{c \geq 1} \sum_{\substack{d \in \mathbb{Z} \\ \gcd(c,d)=1}} \sum_{x \in L} \frac{e\left(\frac{a\tau+b}{c\tau+d}\beta(x) + \frac{\beta(x,z)}{c\tau+d} - \frac{c\beta(z)}{c\tau+d}\right)}{(c\tau+d)^k},$$

where a, b are chosen such that $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is in Γ . Using the identity

$$\frac{a\tau+b}{c\tau+d}\beta(x) + \frac{\beta(x,z)}{c\tau+d} - \frac{c\beta(z)}{c\tau+d} = -\frac{c}{c\tau+d}\beta\left(z - \frac{x}{c}\right) + \frac{a}{c}\beta(x)$$

we obtain

$$\sum_{x \in L} \frac{e\left(\frac{a\tau+b}{c\tau+d}\beta(x) + \frac{\beta(x,z)}{c\tau+d} - \frac{c\beta(z)}{c\tau+d}\right)}{(c\tau+d)^k}$$

$$= \sum_{c \geq 1} \frac{1}{c^k} \sum_{\substack{d \bmod c \\ \gcd(c,d)=1}} \sum_{x \in L/cL} e(a\beta(x)) F\left(\tau + \frac{d}{c}, z - \frac{x}{c}\right),$$

where

$$F(\tau, z) := \sum_{h \in \mathbb{Z}} \sum_{y \in L} \frac{e\left(-\frac{\beta(z+y)}{\tau+h}\right)}{(\tau+h)^k}.$$

Applying Poisson summation we obtain

$$F(\tau, z) = \sum_{n \in \mathbb{Z}, r \in L^\#} \omega(n, r) q^n \zeta_\beta^r$$

with

$$\begin{aligned} \omega(n, r) &= \int_{\mathbb{R} \times (L \otimes_{\mathbb{Z}} \mathbb{R})} \tau^{-k} e\left(-\frac{\beta(z)}{\tau}\right) e(-(n\tau + \beta(r, z))) d\tau dz \\ &= \int_{\Im \tau = v} \tau^{-k} e(D\tau) d\tau \int_{L \otimes_{\mathbb{Z}} \mathbb{R}} e\left(-\frac{\beta(z + r\tau)}{\tau}\right) dz. \end{aligned}$$

The inner integral equals to $\frac{\tau^{r_{\underline{L}}/2}}{i^{r_{\underline{L}}/2} \sqrt{\det(\underline{L})}}$. By [5, Page 19],

$$\omega(n, r) = \frac{(2\pi)^{k - \frac{r_{\underline{L}}}{2}} i^k |D|^{k - \frac{r_{\underline{L}}}{2} - 1}}{\sqrt{\det(\underline{L})} \Gamma(k - \frac{r_{\underline{L}}}{2})}$$

if $D < 0$ and 0 otherwise. Summing up we find

$$(1) \quad E_{k, \underline{L}}(\tau, z) = \vartheta_{\underline{L}}(\tau, z) + \sum_{\substack{n \in \mathbb{Z}, r \in L^\# \\ \beta(r) < n}} \frac{(2\pi)^{k - \frac{r_{\underline{L}}}{2}} i^k |D|^{k - \frac{r_{\underline{L}}}{2} - 1}}{\sqrt{\det(\underline{L})} \Gamma(k - \frac{r_{\underline{L}}}{2})} L(\gamma_{n,r}, k) q^n \zeta_\beta^r,$$

where $L(\gamma_{n,r}, k) := \sum_{c \geq 1} \frac{\gamma_{n,r}(c)}{c^k}$ with

$$\gamma_{n,r}(c) := \sum_{\substack{d \in \mathbb{Z} \\ \gcd(c,d)=1}} \sum_{x \in L/cL} e(d(\beta(x) + \beta(r, x) + n))$$

(For the expression of $\gamma_{n,r}(c)$ we replaced x by dx). Note that $\gamma_{n,r}(c)$ is multiplicative in c . Thus

$$L(\gamma_{n,r}, k) = \prod_p L_p(\gamma_{n,r}, k) \quad \text{with} \quad L_p(\gamma_{n,r}, k) := \sum_{\nu \geq 0} \frac{\gamma_{n,r}(p^\nu)}{p^{k\nu}}.$$

By [3, Theorem 7], [3, Theorem 11],

$$\prod_{p \nmid \Delta_{\underline{L}}} L_p(\gamma_{n,r}, k) = \frac{1}{L(\chi_{\underline{L}}, k - \frac{r_{\underline{L}}}{2})} \sum_{d \mid |\Delta_{\underline{L}} D|} d^{1-k + \frac{r_{\underline{L}}}{2}} \chi_{\underline{L}}(d).$$

Applying the functional equation for L -function we have

$$\begin{aligned}
 & \frac{(2\pi)^{k-\frac{r_L}{2}} i^k |D|^{k-\frac{r_L}{2}-1}}{\sqrt{\det(\underline{L})} \Gamma(k-\frac{r_L}{2})} \prod_{p \nmid \Delta_{\underline{L}}} L_p(\gamma_{n,r}, k) \\
 (2) \quad &= \frac{2(-1)^{\lceil \frac{r_L}{4} \rceil}}{L(\chi_{\underline{L}}, 1-k+\frac{r_L}{2})} \sum_{d \mid |\Delta_{\underline{L}} D|} d^{k-\frac{r_L}{2}-1} \chi_{\underline{L}}(|\Delta_{\underline{L}}| D/d).
 \end{aligned}$$

Now we calculate $L_p(\gamma_{n,r}, k)$ for $p \mid \Delta_{\underline{L}}$.

If prime p does not divide ω_r then

$$(3) \quad \gamma_{n,r}(p^\nu) = \sum_{\substack{d \bmod p^\nu \\ \gcd(d, p^\nu)=1}} e_{p^\nu}(-d\omega_r^2 D) \sum_{x \in L/p^\nu L} e_{p^\nu}(d(\beta(x))).$$

We have that the quadratic form β is \mathbb{Z}_p -isomorphic to the form $x_1^2 + \cdots + a'_p x_{r_L-1}^2 + pa_p x_{r_L}^2$ where $a'_p \in \mathbb{Z}$ such that $a_p a'_p = 2^{r_L} \det(\underline{L})/p$ (see [4, Chapter 11, Theorem 2]). Thus we calculate the inner sum

$$\begin{aligned}
 & \sum_{x \in L/p^\nu L} e_{p^\nu}(d(\beta(x))) \\
 &= \prod_{j=0}^{r_L-2} \left(\sum_{x_j \bmod p^\nu} e_{p^\nu}(dx_j^2) \right) \sum_{x_{r_L-1} \bmod p^\nu} e_{p^\nu}(da'_p x_{r_L-1}^2) \sum_{x_{r_L} \bmod p^\nu} e_{p^\nu}(da_p p x_{r_L}^2) \\
 &= p^{\frac{\nu(r_L-1)}{2}} \left(\frac{d}{p^\nu} \right)^{r_L-1} \left(\frac{a'_p}{p} \right) \sqrt{\left(\frac{-1}{p^\nu} \right)^{r_L-1}} \times p^{\frac{\nu+1}{2}} \left(\frac{a_p d}{p^{\nu-1}} \right) \sqrt{\left(\frac{-1}{p^{\nu-1}} \right)} \\
 &= p^{\frac{\nu r_L+1}{2}} \left(\frac{a'_p a_p}{p^\nu} \right) \left(\frac{a_p d}{p} \right) \sqrt{\left(\frac{-1}{p^\nu} \right)^{r_L-1}} \sqrt{\left(\frac{-1}{p^{\nu-1}} \right)} \\
 &= p^{\frac{\nu r_L+1}{2}} \left(\frac{\det(\underline{L})/p}{p^\nu} \right) \left(\frac{a_p d}{p} \right) \sqrt{\left(\frac{-1}{p^\nu} \right)^{r_L-1}} \sqrt{\left(\frac{-1}{p^{\nu-1}} \right)}.
 \end{aligned}$$

Moreover,

$$\begin{aligned}
 & \sqrt{\left(\frac{-1}{p^\nu} \right)^{r_L-1}} \sqrt{\left(\frac{-1}{p^{\nu-1}} \right)} \\
 &= \sqrt{\left(\frac{-1}{p} \right)^{\nu(r_L-1)}} \left(\frac{-1}{p} \right)^{\lfloor \frac{\nu}{2} \rfloor (r_L-1)} \sqrt{\left(\frac{-1}{p} \right)^{\nu-1}} \left(\frac{-1}{p} \right)^{\lfloor \frac{\nu-1}{2} \rfloor} \\
 &= \sqrt{\left(\frac{-1}{p} \right)^{\nu r_L}} \sqrt{\left(\frac{-1}{p} \right)^{-1}} \left(\frac{-1}{p} \right)^{\lfloor \frac{\nu}{2} \rfloor + \lfloor \frac{\nu-1}{2} \rfloor}
 \end{aligned}$$

$$\begin{aligned}
&= \left(\frac{(-1)^{\frac{r_L}{2}}}{p^\nu} \right) \sqrt{\left(\frac{-1}{p} \right) \left(\frac{-1}{p} \right)^{\lfloor \frac{\nu}{2} \rfloor + \lfloor \frac{\nu-1}{2} \rfloor + 1}} \\
&= \sqrt{\left(\frac{-1}{p} \right) \left(\frac{(-1)^{\frac{r_L}{2}+1}}{p^\nu} \right)}.
\end{aligned}$$

Therefore

$$\sum_{x \in L/p^\nu L} e_{p^\nu}(d\beta(x)) = p^{\frac{\nu r_L + 1}{2}} \sqrt{\left(\frac{-1}{p} \right) \left(\frac{-\Delta_L/p}{p^\nu} \right) \left(\frac{a_p d}{p} \right)}.$$

Inserting this into (3) and applying [1, Lemma 2.1.14] we find

$$(4) \quad \gamma_{n,r}(p^\nu) = \begin{cases} p^{\nu(\frac{r_L}{2}+1)} \left(\frac{-\Delta_L/p}{p^\nu} \right) \left(\frac{a_p \omega_r^2 D / p^{\nu_p(\omega_r^2 D)}}{p} \right), & p^\nu \parallel pD, \\ 0, & \text{otherwise.} \end{cases}$$

If $p \mid \omega_r$, then the quadratic polynomial $\beta(x) + \beta(r, x)$ is \mathbb{Z}_p -isomorphic to the form

$$x_1^2 + \cdots + a'_p x_{r_L-1}^2 + pa_p x_{r_L}^2 + 2(r_1 x_1 + \cdots + a'_p r_{r_L-1} x_{r_L-1} + a_p r_{r_L} x_{r_L}),$$

where $r_1, \dots, r_{r_L} \in \mathbb{Z}$ with $p \nmid r_{r_L}$. For $\nu \geq 1$ and each $u \bmod p^\nu$, the equation $px_{r_L}^2 + 2a_p r_{r_L} x_{r_L} \equiv u \bmod p^\nu$ has exact one solution in $\mathbb{Z}/p^\nu \mathbb{Z}$, which implies that

$$\sum_{x_{r_L} \bmod p^\nu} e_{p^\nu}(d(px_{r_L}^2 + 2a_p r_{r_L} x_{r_L})) = \sum_{u \bmod p^\nu} e_{p^\nu}(du) = 0.$$

Therefore $\gamma_{n,r}(p^\nu) = 0$ for $\nu \geq 1$.

By above discussions, for $p \mid \Delta_L$, if $p \mid \omega_r$, then $L_p(\gamma_{n,r}, k) = 1$ and if $p \nmid \omega_r$, then rewrite the first item of (4) in terms of the character associated with lattice we obtain

$$L_p(\gamma_{n,r}, k) = 1 + p^{\nu_p(|\Delta_L D|)(1-k+\frac{r_L}{2})} \chi_L(a_p \Delta_L D / p^{\nu_p(\Delta_L^2 D)}, p) \chi_L(-1/p, p^{\nu_p(\Delta_L D)}).$$

Inserting (2) and the formula for $L_p(\gamma_{n,r}, k)$ with $p \mid \Delta_L$ into (1) we complete the proof of Theorem 3.1. \square

4. Corresponding to Eisenstein series of modular forms

In this section we suppose that the discriminant of L is prime p thus $D_L \approx (\mathbb{Z}/p\mathbb{Z}, x \rightarrow \frac{a_p x^2}{p})$. In [1, Chapter 5] Ajouz constructed a surjective map from $J_{k,L}(\Gamma)$ to $M_{k-\frac{r_L}{2}}(p, \chi_L)$, the space of modular forms of weight $k - \frac{r_L}{2}$ on $\Gamma_0(p)$ with character χ_L . Explicitly the application

$$\Omega : J_{k,L}(\Gamma) \rightarrow M_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(p, \chi_L)$$

given by

$$\sum_{\substack{n \in \mathbb{Z}, r \in L^\sharp \\ \beta(r) \leq n}} c_\phi(n, r) q^n \zeta_\beta^r \rightarrow \sum_{n \geq 0} \sum_{\substack{x \in L^\sharp/L \\ -n/p \equiv \beta(x) \pmod{\mathbb{Z}}}} C_\phi(-n/p, x) q^n$$

is an isomorphism, where

$$M_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(p, \chi_L) = \left\{ f(\tau) \in M_{k-\frac{r_L}{2}}(p, \chi_L) : f(\tau) = \sum_{\chi_L(-n) \neq -\chi_L(a_p)} a_f(n) q^n \right\}.$$

Let the two Eisenstein series of $M_{k-\frac{r_L}{2}}(p, \chi_L)$ be

$$E_{k-\frac{r_L}{2}}^{1, \chi_L}(\tau) = 1 + \frac{2}{L(1-k+\frac{r_L}{2}, \chi_L)} \sum_{n>0} \sum_{d|n} d^{k-\frac{r_L}{2}-1} \chi_L(d) q^n,$$

$$E_{k-\frac{r_L}{2}}^{\chi_L, 1}(\tau) = \sum_{n>0} \sum_{d|n} d^{k-\frac{r_L}{2}-1} \chi_L(n/d) q^n.$$

By [2, (14)], the unique Eisenstein series of $M_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(p, \chi_L)$ is

$$E_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(\tau) := E_{k-\frac{r_L}{2}}^{1, \chi_L}(\tau) + \frac{2\chi_L(-a_p)}{L(1-k+\frac{r_L}{2}, \chi_L)} E_{k-\frac{r_L}{2}}^{\chi_L, 1}(\tau).$$

We now show that $\Omega(E_{k, \underline{L}}(\tau, z)) = E_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(\tau)$. It is deduced by the table in [1, Page 92] that $(-1)^{\lfloor \frac{r_L}{4} \rfloor} \chi_L(-1) = \chi_L(a_p)$. By this we rewrite the Fourier expansion of $E_{k, \underline{L}}(\tau, z)$ as

$$E_{k, \underline{L}}(\tau, z) = \sum_{x \in L} q^{\beta(x)} \zeta_\beta^x + \sum_{\substack{n \in \mathbb{Z}, r \in L^\sharp \\ \beta(r) < n}} c(n, r) q^n \zeta_\beta^r$$

with

$$c(n, r) = \frac{2}{L(1-k+\frac{r_L}{2}, \chi_L)} \times \begin{cases} \chi_L(-a_p) \sum_{d|-pD} d^{k-\frac{r_L}{2}-1} \chi_L(-pD/d) + \sum_{d|-pD} d^{k-\frac{r_L}{2}-1} \chi_L(d), & r \in L, \\ \chi_L(-a_p) \sum_{d|-pD} d^{k-\frac{r_L}{2}-1} \chi_L(-pD/d), & r \notin L. \end{cases}$$

The condition $D \in \mathbb{Z}$ is equivalent to $r \in L$, and for $r \notin L$, the equation $\beta(x) = -D$ has exact $1 + \chi_L(-a_p D)$ solutions in L^\sharp/L . Therefore

$$\begin{aligned} & \Omega(E_{k, \underline{L}}(\tau, z)) \\ &= 1 + \frac{2}{L(1-k+\frac{r_L}{2}, \chi_L)} \end{aligned}$$

$$\begin{aligned}
& \times \sum_{n>0, p|n} \left(\sum_{d|n} d^{k-\frac{r_L}{2}-1} \chi_L(d) + \chi_L(-a_p) \sum_{d|n} d^{k-\frac{r_L}{2}-1} \chi_L(n/d) \right) q^n \\
& + \frac{2\chi_L(-a_p)}{L(1-k+\frac{r_L}{2}, \chi_L)} \sum_{n>0, p \nmid n} (1 + \chi_L(-a_p n)) d^{k-\frac{r_L}{2}-1} \chi_L(n/d) q^n \\
& = E_{k-\frac{r_L}{2}}^{1, \chi_L}(\tau) + \frac{2\chi_L(-a_p)}{L(1-k+\frac{r_L}{2}, \chi_L)} E_{k-\frac{r_L}{2}}^{\chi_L, 1}(\tau) \\
& = E_{k-\frac{r_L}{2}}^{\chi_L(a_p)}(\tau).
\end{aligned}$$

Acknowledgement. The author would like to thank the referees for their valuable comments on this work.

References

- [1] A. Ajouz, *Hecke operators on Jacobi forms of lattice index and the relation to elliptic modular forms*, Ph.D. thesis, University of Siegen, 2015.
- [2] J. H. Bruinier and M. Bundschuh, *On Borcherds products associated with lattices of prime discriminant*, Ramanujan J. **7** (2003), no. 1-3, 49–61. <https://doi.org/10.1023/A:1026222507219>
- [3] J. H. Bruinier and M. Kuss, *Eisenstein series attached to lattices and modular forms on orthogonal groups*, Manuscripta Math. **106** (2001), no. 4, 443–459. <https://doi.org/10.1007/s229-001-8027-1>
- [4] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, third edition, Grundlehren der Mathematischen Wissenschaften, **290**, Springer-Verlag, New York, 1999. <https://doi.org/10.1007/978-1-4757-6568-7>
- [5] M. Eichler and D. Zagier, *The theory of Jacobi forms*, Progress in Mathematics, **55**, Birkhäuser Boston, Inc., Boston, MA, 1985. <https://doi.org/10.1007/978-1-4684-9162-3>
- [6] A. Mocanu, *Poincaré and Eisenstein series for Jacobi forms of lattice index*, arXiv1712.08174v2, 2018.
- [7] M. Woitalla, *Calculating the Fourier coefficients of Jacobi-Eisenstein series*, arXiv1705.04595v2, 2017.

RAN XIONG
SCHOOL OF MATHEMATICAL SCIENCES
TONGJI UNIVERSITY
SHANGHAI. 200092, P. R. CHINA
Email address: ranxiong2012@163.com